

ESOMEPRAZOLE- esomeprazole magnesium capsule, delayed release

YYBA CORP

Esomeprazole

Drug Facts

Active ingredient (in each capsule)

*Esomeprazole 20 mg (Each delayed-release capsule contains 22.3 mg esomeprazole magnesium trihydrate)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

- Do not use if you are allergic to esomeprazole
- Esomeprazole may cause severe skin reactions. Symptoms include:

-
- skin reddening
 - blisters
 - rash
-

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (**1-800-222-1222**).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use.
- Store at 20°C to 25°C (68°F to 77°F). [See USP controlled room temperature.]
- keep the carton. It contains important information.

Inactive Ingredients

FD & C blue no 1, FD & C red no 3, ferrousferrous oxide, gelatin, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, mono and di glycerides, polysorbate 80, potassium hydroxide, propylene glycol, shellac, sugar spheres (corn starch and sucrose), talc, titanium dioxide and triethyl citrate.

Questions?

Call toll-free weekdays 9 AM to 6 PM EST at **866-933-6337**.

TAMPER-EVIDENT FEATURES: Do not use if seal under bottle cap imprinted with "*SEALED for YOUR PROTECTION*" or blue band around center of each capsule is broken or missing.

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Distributed by:

Wellspring

Airmont, NY 10952, U.S.A.

R0922

PRINCIPAL DISPLAY PANEL - 42 Capsule Bottle Carton

welmate™

See new warning
information

†Compare To the
active ingredient of
Nexium® 24 HR

Heartburn Relief

Acid Reducer
Esomeprazole Magnesium Delayed-Release Capsules USP, 20 mg*

24
HOUR

- ✓Treats Frequent Heartburn
- ✓May Take 1 to 4 Days for Full Effect

#1
PHARMACIST
RECOMMENDED

42 CAPSULES

3 X 14 Count Bottles Inside
Three 14-day courses of treatment

Heartburn Relief

Acid Reducer

Esomeprazole Magnesium Delayed-Release Capsules USP, 20 mg*



Treats Frequent Heartburn



See new warning information

†Compare To the active ingredient of Nexium® 24 HR

Heartburn Relief

Acid Reducer

Esomeprazole Magnesium Delayed-Release Capsules USP, 20 mg*



- ✓ Treats Frequent Heartburn
- ✓ May Take 1 to 4 Days for Full Effect

#1 PHARMACIST RECOMMENDED

42 CAPSULES 3 X 14 Count Bottles Inside
Three 14-day courses of treatment



Tips for Managing Heartburn.

■ Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables. ■ Eat slowly and do not eat big meals. ■ Do not eat late at night or just before bedtime. ■ Do not lie flat or bend over soon after eating. ■ Raise the head of your bed. ■ Wear loose-fitting clothing around your stomach. ■ If you are overweight, lose weight. ■ If you smoke, quit smoking.

Distributed by:
WellSpring
Airmont, NY 10952, U.S.A.

0523



3 73581 00026 5

†All trademarks are property of their respective owners.
This product is not affiliated with the makers/owners of Nexium®.

Batch No.

Expiration Date:

Non Varnish Area



5241217

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active Ingredient (in each capsule) Purpose

*Esomeprazole 20 mg, Acid reducer
(Each delayed-release capsule contains 22.3 mg esomeprazole magnesium trihydrate)

Uses

■ treats frequent heartburn (occurs 2 or more days a week)

Drug Facts (continued)

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Drug Facts (continued)

- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor
- Repeated 14-Day Courses (if needed)**
- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

DO NOT INTERRUPT OR STOP THIS DRUG FOR MORE THAN 4 DAYS FOR FULL EFFECT

Warnings
Allergy alert:
 Do not use if you are allergic to esomeprazole
 Esomeprazole may cause severe skin reactions. Symptoms include:
 skin reddening blisters rash
 If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
 trouble or pain swallowing food, vomiting with blood, or bloody or black stools
 heartburn with lightheadedness, sweating or dizziness
 chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
 frequent chest pain
 These may be signs of a serious condition. See your doctor.

stop use and ask a doctor if
 your heartburn continues or worsens
 you need to take this product for more than 14 days
 you need to take more than 1 course of treatment every 4 months
 you get diarrhea
 you develop a rash or joint pain

pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 adults 18 years of age and older
 this product is to be used once a day (every 24 hours), every day for 14 days
 may take 1 to 4 days for full effect

14-Day Course of Treatment
 swallow 1 capsule with a glass of water before eating in the morning
 take every day for 14 days

read the directions and warnings before use
 Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room temperature.]
 keep the carton. It contains important information.

Inactive ingredients
 FD & C blue no 1, FD & C red no 3, ferrousferic oxide, gelatin, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, mono and di glycerides, polysorbate 80, potassium hydroxide, propylene glycol, shellac, sugar spheres (corn starch and sucrose), talc, titanium dioxide and triethyl citrate.

Questions? Call toll-free weekdays 9 AM to 6 PM EST at 866-933-6337.

TAMPER-EVIDENT FEATURES: Do not use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or blue band around center of each capsule is broken or missing.



ESOMEPRAZOLE

esomeprazole magnesium capsule, delayed release

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:73581-013 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ESOMEPRAZOLE MAGNESIUM (UNII: R6DXU4WAY9) (ESOMEPRAZOLE - UNII:N3PA6559FT) | ESOMEPRAZOLE | 20 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| FERROSFERRIC OXIDE (UNII: XM0M87F357) | |
| GELATIN, UNSPECIFIED (UNII: 2G86QN327L) | |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) | |
| HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K) | |
| MAGNESIUM STEARATE (UNII: 70097M6130) | |
| METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER (UNII: NX76LV5T8J) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| POTASSIUM HYDROXIDE (UNII: WZH3C48M4T) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| SHELLAC (UNII: 46N107B71O) | |
| TALC (UNII: 7SEV7J4R1U) | |

| | |
|---|--|
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| TRIETHYL CITRATE (UNII: 8Z96QXD6UM) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| SUCROSE (UNII: C151H8M554) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |
| FD&C RED NO. 3 (UNII: PN2ZH5LOQY) | |
| GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4) | |

Product Characteristics

| | | | |
|-----------------|---------|---------------------|----------|
| Color | PINK | Score | no score |
| Shape | CAPSULE | Size | 14mm |
| Flavor | | Imprint Code | RG;50 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:73581-013-42 | 3 in 1 CARTON | 03/08/2023 | |
| 1 | | 14 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA212866 | 03/08/2023 | |

Labeler - YYBA CORP (006339772)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------------|---------|-----------|------------------------|
| Ohm Laboratories Inc. | | 184769029 | MANUFACTURE(73581-013) |

Revised: 9/2023

YYBA CORP