

HEAD CONGESTION PLUS MUCUS PE- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated
United Natural Foods, Inc. dba UNFI

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1173-ELN-2023-0517

Drug Facts

| Active ingredients (in each caplet) | Purpose |
|--|-----------------------------|
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Guaifenesin 200 mg | Expectorant |
| Phenylephrine HCl 5 mg | Nasal decongestant |

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - minor aches and pains
 - nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

| | |
|---------------------------------------|--|
| adults and children 12 years and over | <ul style="list-style-type: none"> ▪ take 2 caplets every 4 hours ▪ do not take more than 10 caplets in 24 hours |
| children under 12 years | <ul style="list-style-type: none"> ▪ ask a doctor |

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-855-423-2630

Principle Display Panel

EQUALINE®

NDC 41163-712-02

compare to Sudafed PE® Head Congestion + Mucus active ingredients†

head congestion plus mucus PE

acetaminophen (pain reliever/fever reducer)

guaifenesin (expectorant)

phenylephrine HCl (nasal decongestant)

for relief of

- sinus pressure
- headache
- chest congestion

24 caplets

actual size

| | |
|---|---|
| Drug Facts (continued) Other information ■ store between 20°-25°C (68°-77°F) in a dry place ■ retain carton for complete product information and warnings Inactive ingredients colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, | Drug Facts (continued) Questions or comments? 1-855-423-2630 magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide |
|---|---|

| | |
|--|---|
| Drug Facts (continued) Warnings Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients Ask a doctor before use if you have ■ liver disease ■ heart disease ■ thyroid disease ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus) | Drug Facts (continued) Directions ■ do not take more than directed (see overdose warning) adults and children 12 years and over ■ take 2 caplets every 4 hours ■ do not take more than 10 caplets in 24 hours ■ ask a doctor 12 years and under Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Keep out of reach of children. If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition. ■ cough comes back or occurs with rash or headache that lasts ■ new symptoms occur ■ redness or swelling is present ■ fever gets worse or lasts more than 3 days ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ nervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if When using this product do not exceed recommended dosage Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin Drug Facts (continued) |
|--|---|



| | |
|--|--|
| Drug Facts Active ingredients (in each caplet) Acetaminophen 325 mg.....Pain reliever/fever reducer Guaifenesin 200 mg.....Expectorant Phenylephrine HCl 5 mg.....Nasal decongestant Uses ■ temporarily relieves these symptoms associated with | Drug Facts (continued) may fever or other upper respiratory allergies, and the common cold: ■ sinus congestion and pressure ■ headache ■ nasal congestion ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily reduces fever |
|--|--|

EQUALINE®

NDC 411163-712-02

compare to Sudafed PE® Head Congestion + Mucus active ingredients†

head congestion plus mucus PE

acetaminophen (pain reliever/fever reducer)
 guaifenesin (expectorant)
 phenylephrine HCl (nasal decongestant)

for relief of

- sinus pressure
- headache
- chest congestion

24 caplets



actual size

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Sudafed PE® Head Congestion + Mucus.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

Like it or let us make it right. That's our quality promise. 855-423-2630

DISTRIBUTED BY UNFI PROVIDENCE, RI 02908 USA

HEAD CONGESTION PLUS MUCUS PE

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:41163-712 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 200 mg |
| PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV) | PHENYLEPHRINE | 5 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| CROSPVIDONE (UNII: 2S7830E561) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | |
| STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TALC (UNII: 7SEV7J4R1U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | white | Score | no score |
| Shape | OVAL | Size | 19mm |
| Flavor | | Imprint Code | AAA;1173 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:41163-712-02 | 2 in 1 CARTON | 08/19/2022 | |
| 1 | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part341 | 08/19/2022 | |

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 5/2023

United Natural Foods, Inc. dba UNFI