

ASPIRIN LOW DOSE- aspirin tablet, chewable
L.N.K. International, Inc.

Sound Body 44-218A

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - have bloody or black stools
 - feel faint
 - vomit blood
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- see end flap for expiration date and lot number
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

SOUNDBODY™

†Compare to the active ingredient in Bayer® Chewable Low Dose Aspirin

LOW DOSE

Aspirin

Chewable Tablets

Pain Reliever (**NSAID**) 81 mg

NDC 50844-219-07

Aspirin Regimen**

Chew or crush tablets completely before swallowing.

Orange
Flavored

36 CHEWABLE TABLETS

Actual Size

**Talk to your doctor or other healthcare provider before using this product for your heart.

Triple Pack 3 bottles x 36 tablets each = 108 tablets

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.

50844 REV0122A21807

Manufactured for Big Lots Stores, Inc.

by **LNK INTERNATIONAL, INC.**

60 Arkay Drive, Hauppauge, NY 11788 USA

V#733000 ITEM#022721807



Aspirin

Chewable Tablets
Pain Reliever (NSAID) 81 mg

Triple Pack

3 bottles x 36 tablets each =
108 tablets

Aspirin

Chewable Tablets
Pain Reliever (NSAID) 81 mg



**Talk to your doctor or other healthcare provider before using this product for your heart.



[†]Compare to the active ingredient in Bayer® Chewable Low Dose Aspirin

LOW DOSE

NDC 50844-219-97

Aspirin

Chewable Tablets
Pain Reliever (NSAID) 81 mg

36 CHEWABLE TABLETS

Aspirin Regimen**
Chew or crush tablets completely before swallowing.

Orange Flavored



Actual Size



[†]Compare to the active ingredient in Bayer® Chewable Low Dose Aspirin

LOW DOSE

Aspirin

Chewable Tablets
Pain Reliever (NSAID) 81 mg

36 CHEWABLE TABLETS

Aspirin Regimen**
Chew or crush tablets completely before swallowing.

Orange Flavored



Actual Size



[†]Compare to the active ingredient in Bayer® Chewable Low Dose Aspirin

LOW DOSE

Aspirin

Chewable Tablets
Pain Reliever (NSAID) 81 mg

36 CHEWABLE TABLETS

Aspirin Regimen**
Chew or crush tablets completely before swallowing.

Orange Flavored



Actual Size

Triple Pack 3 bottles x 36 tablets each = 108 tablets

[†]This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.

50844 REV0122A21807
Manufactured for Big Lots Stores, Inc.
by LINK INTERNATIONAL, INC.
60 Atkey Drive, Hauppauge, NY 11788 USA
V#733000 ITEM#022721807

No print area
Lot no. & Exp. date

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



3 50844 21907 7

Drug Facts (continued)

- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information ■ see end flap for expiration date and lot number
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients corn starch, dextrates hydrated, FD&C yellow #6 aluminum lake, flavo, maltodextrin, sucralose

Questions or comments? 1-800-426-9391

B-0227-218A-07BR
REV0122A21807

Directions ■ do not take more than directed
 ■ drink a full glass of water with each dose

Learn more at www.aspirin.com

Drug Facts
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each chewable tablet)
Aspirin 81 mg (NSAID)
Pain reliever
nonsteroidal anti-inflammatory drug

Uses
for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ take more or for a longer time than directed
■ have 3 or more alcoholic drinks every day while using this product
■ have had stomach ulcers or bleeding problems

Do not use
■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you have asthma ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ have bloody or black stools ■ feel faint ■ vomit blood
■ have stomach pain that does not get better
■ ringing in the ears or a loss of hearing occurs ■ redness or swelling is present
■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days
■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control

Sound Body 44-218A

ASPIRIN LOW DOSE			
aspirin tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-219
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	
Inactive Ingredients			
Ingredient Name	Strength		
STARCH, CORN (UNII: O8232NY3SJ)			
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			

SUCRALOSE (UNII: 96K6UQ3ZD4)

Product Characteristics

Color	orange (light)	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	44;218
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-219-07	3 in 1 CARTON	01/16/2020	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/16/2020	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-219)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-219)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-219)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-219)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(50844-219)

