

**DG HEALTH ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride,
simethicone tablet
Dolgenercorp, LLC**

Dolgenercorp, LLC Anti-Diarrheal/Anti-Gas Drug Facts

Active ingredients (in each caplet)

Loperamide HCl 2 mg

Simethicone 125 mg

Purposes

Anti-diarrheal

Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25° C (68°-77° F). Protect from light.
- **do not use if printed blister unit is broken or torn**

Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin,

microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

Questions or comments?

1-888-309-9030

Package/Label Principal Display Panel

DG™ |health

Compare to the active ingredients of Imodium® Multi-Symptom Relief

Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125mg

Anti-Diarrheal/Anti-Gas

Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea plus

- Cramps & Pressure
- Bloating
- Gas

12 Caplets*

*Capsule-Shaped Tablets

Actual Caplet Size



Drug Facts (continued)
Active Ingredients Loperamide hydrochloride, simethicone, calcium phosphate, microcrystalline cellulose, and indandiolololone.
 How: polymeric, white to off-white powder.

Drug Facts
Active Ingredients (in each caplet) Loperamide HCl 2 mg, Simethicone 125 mg.
Purposes Antidiarrheal and anti-gas.
Uses relieves symptoms of diarrhea and gas.
Warnings
 Allergy alert: Do not use if you have ever had a reaction to loperamide HCl or other ingredients.
 Heart alert: Taking more than the directed amount may cause a slow heart rate.
 Do not use if you have had a heart attack, heart failure, or other heart conditions.
 Ask a doctor before use if you have a history of dizziness or fainting, a history of low blood pressure, or a history of heart rhythm problems.
 Ask a doctor or pharmacist before use if you are taking any other drugs, especially those used for heart disease, high blood pressure, or other conditions.
 When using this product, dizziness or other symptoms may occur. Do not drink alcohol or use grapefruit juice while taking this product.
 Stop use and ask a doctor if:
 ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bloating, or may be bloated
 ■ you have difficulty swallowing the caplet
Other Information
 Each caplet contains calcium 165 mg, sodium 3 mg, and iron 0.25 mg. Protect from light.
 Do not use if printed blister unit is broken or torn.

CODE AREA

OR153 VT C3

Loperamide Hydrochloride and Simethicone
 Tablets, 2 mg/125 mg

Anti-Diarrheal/Anti-Gas

Multi-Symptom Relief Caplet



Compare to the active ingredients of Imodium® Multi-Symptom Relief**

Loperamide Hydrochloride and Simethicone
 Tablets, 2 mg/125 mg

Anti-Diarrheal/Anti-Gas

Multi-Symptom Relief Caplet

Relieves symptoms of diarrhea *plus*

- Cramps & Pressure
- Bloating
- Gas



12 Caplets*

Actual Caplet Size *Capsule-Shaped Tablets

100% Satisfaction Guaranteed
 If you're not satisfied with this product for any reason, please call us so we can make you feel better. (888) 309-9030

100% Satisfaction Guaranteed
 If you're not satisfied with this product for any reason, please call us so we can make you feel better. (888) 309-9030

© 2012
 GOODLETTSVILLE, TN 37022
 OLD EAST MAIN CO. DIVISION R.D. 6
 DISTRIBUTED BY
 Multi-Symptom Relief
 Johnson & Johnson Consumer Products Inc., distributor of
 Imodium®
 Loperamide Hydrochloride and Simethicone Tablets, 2 mg/125 mg
 *This product is not for use in children under 12 years of age.

DG HEALTH ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-383
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L0R1;2;125
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-383-53	12 in 1 CARTON	10/09/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:55910-383-89	18 in 1 CARTON	10/09/2018	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

3	NDC:55910-385-62	24 in 1 CARTON	04/07/2022	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA209837		10/09/2018	

Labeler - Dolgencorp, LLC (068331990)

Revised: 4/2022

Dolgencorp, LLC