

CHILDRENS IBUPROFEN ORAL SUSPENSION DYE FREE- ibuprofen suspension
FSA Store Inc. (Caring Mill)

Childrens Ibuprofen Oral Suspension

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen, USP 100 mg (NSAID)**

**nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters.

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child :

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drug containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stool
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-

800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. if possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- **each 5 mL contains:** Sodium 2 mg
- store between 20° to 25°C (68° to 77°F)
- **do not use if printed neckband is broken or missing**

Inactive ingredients

Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments?

Call 1-888-588-1418 from 9 AM to 5 PM ET, Monday-Friday.

Principal display panel list

NDC 81522-311-11

†Compare to the active ingredient in Children's Motrin®

FOR AGES 2-11 YEARS

CHILDREN'S IBUPROFEN

100 mg per 5 mL

Oral Suspension, USP (**NSAID**)

PAIN RELIEVER/FEVER REDUCER

ORIGINAL Berry Flavor

- Dye-Free
- Alcohol-Free

Lasts up to 8 HOURS

4 FL OZ (118mL)

Important: Read all product information before using. Keep this box for important information.

This product is intended for use in children.

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Manufactured for **FSA Store Inc.**

5473 Blair Rd Suite 100

PMB 24308 Dallas, TX 75231

Drug Facts (continued)

- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- 1 mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Drug Facts (continued)

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL) *
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information

- each 5 mL contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if cap is opened or printed bottle neckband is broken or missing
- see bottom panel for lot number and expiration date

Inactive ingredients

Acetaminophen potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments? Call 1-888-688-1415 from 9 AM to 5 PM ET, Monday-Friday.

** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Manufactured for FSA Store Inc.
5473 Blair Rd Suite 100
PMB 24308 Dallas, TX 75231

caring mill™

Children's Ibuprofen
For Ages 2-11 Years

Oral Suspension (NSAID)
100mg per 5mL

NDC 81522-311-11

*Compare to the active ingredient in Children's Motrin®

caring mill™

Children's
Ibuprofen
For Ages 2-11 Years

Oral Suspension (NSAID)

100mg per 5mL

Pain Reliever/Fever Reducer

Original Berry Flavor



Dye-Free
Alcohol-Free



See New Warnings

4 FL OZ
(118mL)

Important: Read all product information before using.
Keep this box for important information.
This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)	Purpose
Ibuprofen 100 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids

36-149



8 1002592902 2

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID) *Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache ■ reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child: ■ has had stomach ulcers or bleeding

Purpose

NDC 81522-311-11

caring mill™

Children's Ibuprofen

Oral Suspension (NSAID)

100mg per 5mL

Pain Reliever/Fever Reducer - Dye-Free

4 FL OZ (118mL)

Original Berry Flavor

problems ■ takes a blood thinning (anticoagulant) or steroid drug ■ takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ takes more or for a longer time than directed. **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly.

Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. **Do not use** ■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery **Ask a doctor before use if** ■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers ■ child has not been drinking fluids ■ child has lost a lot of fluid due to vomiting or diarrhea ■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke ■ child has asthma ■ child is taking a diuretic

Ask a doctor or pharmacist before use if the child is ■ under a doctor's care for any serious condition ■ taking any other drug **When using this product** ■ take with food or milk if stomach upset occurs **Stop use and ask a doctor if** ■ child experiences any of the following signs of stomach bleeding: ■ feels faint ■ vomits blood ■ has bloody or black stools ■ has stomach pain that does not get better ■ child has symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ the child does not get



LR-223

PEEL HERE

Drug Facts (continued)

any relief within first day (24 hours) of treatment ■ fever or pain gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ this product does not contain directions or complete warnings for adult use

■ do not give more than directed ■ shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ use only enclosed dosing cup. Do not use any other dosing device. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day ■ replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

*or as directed by a doctor

Other information ■ each 5 mL contains: sodium 2 mg ■ store between 20-25°C (68-77°F) ■ do not use if printed neckband is broken or missing

Inactive ingredients Acesulfame potassium, anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, microcrystalline cellulose, polysorbate 80, propylene glycol, purified water, sodium benzoate, sucrose, and xanthan gum

Questions or comments? Call 1-888-589-1418 from 9 AM to 5 PM ET, Monday-Friday.

** This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin®.

Manufactured for FSA Store Inc.
5473 Blair Rd Suite 100
PMB 24308 Dallas, TX 75231

res

CHILDRENS IBUPROFEN ORAL SUSPENSION DYE FREE

ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81522-311
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
GLYCERIN (UNII: PDC6A3C0OX)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	BERRY (ORIGINAL BERRY)	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81522-311-11	1 in 1 CARTON	09/05/2025	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210602	09/05/2025	

Labeler - FSA Store Inc. (Caring Mill) (049283340)

Registrant - APTAPHARMA INC. (790523323)

Establishment			
Name	Address	ID/FEI	Business Operations
APTAPHARMA INC.		790523323	manufacture(81522-311)