

**ALTICOTRON- dextromethorphan hbr, guaifenesin, phenylephrine hcl solution/ drops**  
**Alternative Pharmacal Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

<b>Active Ingredients (in each 1 mL)</b>	<b>Purpose</b>
Dextromethorphan HBr..... 5 mg.....	Cough Suppressant
Guaifenesin..... 100 mg.....	Expectorant
Phenylephrine HCl..... 2.5 mg.....	Nasal Decongestant

Cough Suppressant  
Expectorant  
Nasal Decongestant

**Uses**

- non-narcotic cough suppressant which temporarily calms due to minor throat and bronchial irritation as may occur with common cold
- helps decongest sinus openings and passages; temporarily relieves sinus openings and passages; temporarily relieves sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily relieves nasal congestion due to the common cold
- helps decongest sinus openings and passages; temporarily relieves sinus congestion and pressure

**Warnings**

A persistent cough may be sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor.

**DO NOT USE**

If a child is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug that contains MAOI; ask your doctor or pharmacist before taking this product.

**Ask a doctor before use if the child has:**

- Heart disease
- High blood pressure
- Thyroid diseases
- Diabetes
- Cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or where cough is accompanied by excessive phlegm (mucus)

**When using this product do not exceed recommended dosage.**

**Stop use and ask a doctor before use if:**

- your child gets nervous, dizzy, or sleepless

- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**ⓘ If pregnant or breast-feeding,** ⓘ ask a health profession before use.

**ⓘ Keep out of reach of children.** ⓘ In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- Do not use more than 6 doses in any 24 hour period
- Repeat every 4 hours
- Measure with dosage device provided. Do not use with any other device

Children 6 years to under 12 years of age	2 ml
Children 2 years to under 6 years of age	Ask a doctor
Children under 2 years of age	Ask a doctor

**ⓘ Other Information**

ⓘ Store at room temperature 15<sup>00</sup> - 30<sup>00</sup>°C (59<sup>00</sup>-86<sup>00</sup>°F)

**ⓘ Tamper Evident Feature:** ⓘ Do not use if seal is torn, broken or missing. Avoid excessive heat or humidity.

**ⓘ Inactive ingredients ⓘ**

citric acid, grape flavor, glycerin, methylparaben, polyethylene glycol, propylparaben, purified water, sodium citrate, and sucralose

**ⓘ Questions or comments?** ⓘ 1-786-507-1566

**Drug Facts**

**Active ingredients (in each 1 mL)** **Purpose**  
 Dextromethorphan HBr,..... 5 mg .....Cough Suppressant  
 Guaifenesin, ..... 100 mg .....Expectorant  
 Phenylephrine HCl,..... 2.5 mg .....Nasal Decongestant

**Uses** ■ non-narcotic cough suppressant which temporarily calms due to minor throat and bronchial irritation as may occur with common cold ■ helps decongest sinus openings and passages; temporarily relieves sinus openings and passages; temporarily relieves sinus congestion and pressure ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive ■ temporarily relieves nasal congestion due to the common cold ■ helps decongest sinus openings and passages; temporarily relieves sinus congestion and pressure

**Warnings** ■ a persistent cough may be a sign of a serious condition. If cough persists for more than 1 week, tends to recur, or is accompanied by fever, rash, or persistent headache, consult a doctor ■ **Do not use** in a child who is taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has** ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ cough that lasts or is chronic such as occurs with asthma, chronic bronchitis, or where cough is accompanied by excessive phlegm (mucus)

**When using this product do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ your child gets nervous, dizzy, or sleepless ■ symptoms do not get better within 7 days or are accompanied by fever ■ cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Lot #:

Rev. 9/14

Exp. Date:

NDC 53163-107-02

# Alticotron

## PED Drops



- Cough Suppressant
- Expectorant
- Nasal Decongestant

Contains the same active ingredients as Broncotron® PED Drops

Manufactured For:  
 Alternative Pharmaceutical Corp.  
 Miami, FL 33166  
 www.alternativepharmaceutical.com

Grape Flavor

Sugar FREE  
 Alcohol FREE  
 Dye FREE Gluten FREE

2 fl. oz. (60 mL)

**Drug Facts (continued)**

**Keep out of the reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away.

**Directions:** ■ Do not use more than 6 doses in any 24 hour period ■ Repeat every 4 hours ■ Measure with the dosage device provided. Do not use with any other device.

Age	Dose
Children 6 to under 12 years of age	2 mL
Children 2 to under 6 years of age	Ask a doctor
Children under 2 years of age	Ask a doctor

**Other information** ■ Tamper Evident Feature: **Do not use if cap seal is torn, broken or missing.** ■ Store at room temperature 15-30°C (59-86°F) ■ Avoid excessive heat and humidity

**Inactive ingredients:** Citric acid, grape flavor, glycerin, methylparaben, polyethylene glycol, propylparaben, purified water, sodium citrate, sucralose.

**Questions or comments?** 1-786-507-1566

**Manufactured For:** Alternative Pharmaceutical Corp., Miami, FL 33166. www.alternativepharmaceutical.com

**ALTICOTRON**

dextromethorphan hbr, guaifenesin, phenylephrine hcl solution/ drops

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53163-107
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 1 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 1 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYETHYLENE GLYCOLS</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53163-107-02	60 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	03/01/2015	

**Labeler** - Alternative Pharmacal Corporation (078528214)

Revised: 6/2015

Alternative Pharmacal Corporation