

SODIUM BICARBONATE- sodium bicarbonate injection, solution
Medical Purchasing Solutions, LLC

Sodium Bicarbonate Injection, USP

**FOR CORRECTION OF METABOLIC ACIDOSIS AND OTHER CONDITIONS
REQUIRING SYSTEMIC ALKALINIZATION.**

R_xonly

DESCRIPTION

Sodium Bicarbonate Injection, USP is a sterile, nonpyrogenic, hypertonic solution of sodium bicarbonate (NaHCO_3) in water for injection for administration by the intravenous route as an electrolyte replenisher and systemic alkaliizer.

Solutions are offered in concentrations of 4.2%, 7.5% and 8.4%. See table in **HOW SUPPLIED** section for contents and characteristics.

The solutions contain no bacteriostat, antimicrobial agent or added buffer and are intended only for use as a single-dose injection, the approximate pH of the solutions is 8. When smaller doses are required, the unused portion should be discarded.

Sodium bicarbonate, 84 mg is equal to one milliequivalent each of Na^+ and HCO_3^- . Sodium Bicarbonate, USP is chemically designated NaHCO_3 , a white crystalline powder soluble in water.

Water for Injection, USP is chemically designated H_2O .

CLINICAL PHARMACOLOGY

Intravenous sodium bicarbonate therapy increases plasma bicarbonate, buffers excess hydrogen ion concentration, raises blood pH and reverses the clinical manifestations of acidosis.

Sodium bicarbonate in water dissociates to provide sodium (Na^+) and bicarbonate (HCO_3^-) ions. Sodium (Na^+) is the principal cation of the extracellular fluid and plays a large part in the therapy of fluid and electrolyte disturbances. Bicarbonate (HCO_3^-) is a normal constituent of body fluids and the normal plasma level ranges from 24 to 31 mEq/liter. Plasma concentration is regulated by the kidney through acidification of the urine when there is a deficit or by alkalization of the urine when there is an excess. Bicarbonate anion is considered "labile" since at a proper concentration of hydrogen ion (H^+) it may be converted to carbonic acid (H_2CO_3) and thence to its volatile form, carbon dioxide (CO_2) excreted by the lung. Normally a ratio of 1:20 (carbonic acid: bicarbonate) is present in the extracellular fluid. In a healthy adult with normal kidney function, practically all the glomerular filtered bicarbonate ion is reabsorbed; less than 1% is excreted in the urine.

INDICATIONS AND USAGE

Sodium Bicarbonate Injection, USP is indicated in the treatment of metabolic acidosis which may occur in severe renal disease, uncontrolled diabetes, circulatory insufficiency due to shock or severe dehydration, extracorporeal circulation of blood, cardiac arrest and severe primary lactic acidosis. Sodium bicarbonate is further indicated in the treatment of certain drug intoxications, including barbiturates (where dissociation of the barbiturate-protein complex is desired), in poisoning by salicylates or methyl alcohol and in hemolytic reactions requiring alkalization of the urine to diminish nephrotoxicity of hemoglobin and its breakdown products. Sodium bicarbonate also is indicated in severe diarrhea which is often accompanied by a significant loss of bicarbonate.

Treatment of metabolic acidosis should, if possible, be superimposed on measures designed to control the basic cause of the acidosis - e.g., insulin in uncomplicated diabetes, blood volume restoration in shock. But since an appreciable time interval may elapse before all of the ancillary effects are brought about, bicarbonate therapy is indicated to minimize risks inherent to the acidosis itself.

Vigorous bicarbonate therapy is required in any form of metabolic acidosis where a rapid increase in plasma total CO₂ content is crucial - e.g., cardiac arrest, circulatory insufficiency due to shock or severe dehydration, and in severe primary lactic acidosis or severe diabetic acidosis.

CONTRAINDICATIONS

Sodium Bicarbonate Injection, USP is contraindicated in patients who are losing chloride by vomiting or from continuous gastrointestinal suction, and in patients receiving diuretics known to produce a hypochloremic alkalosis.

WARNINGS

Solutions containing sodium ions should be used with great care, if at all, in patients with congestive heart failure, severe renal insufficiency and in clinical states in which there exists edema with sodium retention.

In patients with diminished renal function, administration of solutions containing sodium ions may result in sodium retention.

The intravenous administration of these solutions can cause fluid and/or solute overloading resulting in dilution of serum electrolyte concentrations, overhydration, congested states or pulmonary edema.

Extravascular infiltration should be avoided, see **ADVERSE REACTIONS**.

PRECAUTIONS

General

Do not use unless solution is clear and the container or seal is intact. Discard unused portion.

The potentially large loads of sodium given with bicarbonate require that caution be exercised in the use of sodium bicarbonate in patients with congestive heart failure or

other edematous or sodium-retaining states, as well as in patients with oliguria or anuria. See table in **HOW SUPPLIED** section for amounts of sodium present in solutions.

Caution must be exercised in the administration of parenteral fluids, especially those containing sodium ions, to patients receiving corticosteroids or corticotropin.

Potassium depletion may predispose to metabolic alkalosis and coexistent hypocalcemia may be associated with carpopedal spasm as the plasma pH rises. These dangers can be minimized if such electrolyte imbalances are appropriately treated prior to or concomitantly with bicarbonate infusion.

Laboratory Tests

The aim of all bicarbonate therapy is to produce a substantial correction of the low total CO₂ content and blood pH, but the risks of overdosage and alkalosis should be avoided. Hence, repeated fractional doses and periodic monitoring by appropriate laboratory tests are recommended to minimize the possibility of overdosage.

Drug Interactions

Additives may be incompatible; norepinephrine and dobutamine are incompatible with sodium bicarbonate solution.

The addition of sodium bicarbonate to parenteral solutions containing calcium should be avoided, except where compatibility has been previously established. Precipitation or haze may result from sodium bicarbonate-calcium admixtures. **NOTE:** Do not use the injection if it contains precipitate.

Additives may be incompatible. Consult with pharmacist, if available. When introducing additives, consult with pharmacist, if available. When introducing additives, use aseptic technique, mix thoroughly and do not store.

Pregnancy: Teratogenic Effects

Pregnancy Category C

Animal reproduction studies have not been conducted with sodium bicarbonate. It is also not known whether sodium bicarbonate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium bicarbonate should be given to a pregnant woman only if clearly needed.

Pediatric

Rapid injection (10 mL/min) of hypertonic Sodium Bicarbonate Injection, USP solutions into neonates and children under two years of age may produce hypernatremia, a decrease in cerebrospinal fluid pressure and possible intracranial hemorrhage. The rate of administration in such patients should therefore be limited to no more than 8 mEq/kg/day. A 4.2% solution may be preferred for such slow administration. In emergencies such as cardiac arrest, the risk of rapid infusion must be weighed against the potential for fatality due to acidosis.

Geriatric

Clinical studies of Sodium Bicarbonate Injection, USP did not include sufficient numbers

of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

ADVERSE REACTIONS

Overly aggressive therapy with Sodium Bicarbonate Injection, USP can result in metabolic alkalosis (associated with muscular twitchings, irritability and tetany) and hypernatremia.

Inadvertent extravasation of intravenously administered hypertonic solutions of sodium bicarbonate have been reported to cause chemical cellulitis because of their alkalinity, with tissue necrosis, ulceration or sloughing at the site of infiltration. Prompt elevation of the part, warmth and local injection of lidocaine or hyaluronidase are recommended to reduce the likelihood of tissue sloughing from extravasated I.V. solutions.

OVERDOSAGE

Should alkalosis result, the bicarbonate should be stopped and the patient managed according to the degree of alkalosis present. 0.9% sodium chloride injection intravenous may be given; potassium chloride also may be indicated if there is hypokalemia. Severe alkalosis may be accompanied by hyperirritability or tetany and these symptoms may be controlled by calcium gluconate. An acidifying agent such as ammonium chloride may also be indicated in severe alkalosis. See **WARNINGS** and **PRECAUTIONS**.

DOSAGE AND ADMINISTRATION

Sodium Bicarbonate Injection, USP is administered by the intravenous route.

In cardiac arrest, a rapid intravenous dose of one to two 50 mL vials (44.6 to 100 mEq) may be given initially and continued at a rate of 50 mL (44.6 to 50 mEq) every 5 to 10 minutes if necessary (as indicated by arterial pH and blood gas monitoring) to reverse the acidosis. Caution should be observed in emergencies where very rapid infusion of large quantities of bicarbonate is indicated. Bicarbonate solutions are hypertonic and may produce an undesirable rise in plasma sodium concentration in the process of correcting the metabolic acidosis. In cardiac arrest, however, the risks from acidosis exceed those of hypernatremia.

In less urgent forms of metabolic acidosis, Sodium Bicarbonate Injection, USP may be added to other intravenous fluids. The amount of bicarbonate to be given to older children and adults over a four-to-eight-hour period is approximately 2 to 5 mEq/kg of body weight - depending upon the severity of the acidosis as judged by the lowering of total CO₂ content, blood pH and clinical condition of the patient. In metabolic acidosis associated with shock, therapy should be monitored by measuring blood gases, plasma osmolarity, arterial blood lactate, hemodynamics and cardiac rhythm. Bicarbonate therapy should always be planned in a stepwise fashion since the degree of response from a given dose is not precisely predictable. Initially an infusion of 2 to 5 mEq/kg body

weight over a period of 4 to 8 hours will produce a measurable improvement in the abnormal acid-base status of the blood. The next step of therapy is dependent upon the clinical response of the patient. If severe symptoms have abated, then the frequency of administration and the size of the dose may be reduced.

In general, it is unwise to attempt full correction of a low total CO₂ content during the first 24 hours of therapy, since this may be accompanied by an unrecognized alkalosis because of a delay in the readjustment of ventilation to normal. Owing to this lag, the achievement of total CO₂ content of about 20 mEq/liter at the end of the first day of therapy will usually be associated with a normal blood pH. Further modification of the acidosis to completely normal values usually occurs in the presence of normal kidney function when and if the cause of the acidosis can be controlled. Values for total CO₂ which are brought to normal or above normal within the first day of therapy are very likely to be associated with grossly alkaline values for blood pH, with ensuing undesired side effects.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. See **PRECAUTIONS**.

HOW SUPPLIED

Sodium Bicarbonate Injection, USP is supplied in the following dosage forms:

NDC Code:	Conc %	mg/mL (NaHCO ₃)	mEq/mL (Na ⁺)	mEq/mL (HCO ₃ ⁻)	mEq/Container size (mL)	mOsmol
51754-5001-4 51754-5001-5	8.4	84	1	1	50/50	2/mL
51754-5011-4	8.4	84	1	1	10/10	2/mL
51754-5002-5	7.5	75	0.9	0.9	44.6/50	1.79/mL
51754-5012-4	4.2	42	0.5	0.5	5/10	1/mL

The 50 mL Sodium Bicarbonate Injection vials are supplied as either 20 vials per carton, or 25 vials per carton, with a single package insert and the 10 mL Sodium Bicarbonate Injection vials are supplied as 25 vials per carton with a single package insert.

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. Protect from freezing. Store at 20°C to 25°C (68°F to 77°F). [See USP Controlled Room Temperature.]

Revised: August 2022

Manufactured by:



Exela Pharma Sciences, LLC
Lenoir, NC 28645

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL- Vial Label

Rx Only

8.4% Sodium Bicarbonate Injection, USP

50 mEq/50 mL

(1 mEq/mL)

For Intravenous Use Only. Discard Unused Portion.

50 mL Single Dose Vial

Each mL contains sodium bicarbonate, 84 mg, 2 mOsmol/mL (calc.), pH 7.0 to 8.5.

Usual dosage: See insert.
Sterile, nonpyrogenic. Discard unused portion.

Store at 20°C to 25°C (68°F to 77°F).
[See USP Controlled Room Temperature]

EXELA
PHARMA SCIENCES
Manufactured and Distributed by:
Exela Pharma Sciences, LLC,
Lenoir, NC 28645 USA

Rx Only
8.4% Sodium Bicarbonate Injection, USP
50 mEq/50 mL
(1 mEq/mL)
For Intravenous Use Only
50 mL Single Dose Vial

(01) 0 0351754 50011 8
LOT
EXP

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL- Outer Package

NDC 71872-7310-1

Rx Only

8.4% Sodium Bicarbonate Injection, USP

50 mEq/50 mL

(1 mEq/mL)

For Intravenous Use Only

1 x 50 mL Single Dose Vial



Rx only



(01) 0 0871872 73101 0
(17) 000101
(10) XXXX-XXXXXX
(21) 7310A000000

8.4% Sodium Bicarbonate Injection, USP 50 mEq/50 mL (1 mEq/mL) 50mL SDV

Qty: 1 vial Lot# XXXX-XXXXXX Exp: 01/01/1900

For Intravenous Use Only

Discard Unused Portion

Each mL contains sodium bicarbonate, 84 mg, 2 mOsmol/mL (calc.),
pH 7.0 to 8.5.

Usual dosage: see insert. Sterile, nonpyrogenic.

Contains no bacteriostat.

Use aseptic technique. Remove cover from Fliptop vial and cleanse
stopper with antiseptic. Use only if clear and seal is intact and undamaged.

Store at 20°C to 25°C (68°F to 77°F).

[See USP Controlled Room Temperature]

50 mL single dose vials. Do not use the injection if it contains precipitate.

Discard unused portion. Do not resterilize.

Manufactured and Distributed by: Exela Pharma Sciences, LLC
Lenoir, NC 28645 USA
Manufacturer NDC: 51754-5001-04

Repackaged

& Distributed By:



Medical Purchasing Solutions

Scottsdale, AZ 85260

www.medicalpurchasingsolutions.com

SODIUM BICARBONATE

sodium bicarbonate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71872-7310(NDC:51754-5001)
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	84 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:71872-7310-1	1 in 1 BAG	09/20/2023	
1		50 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211091	08/01/2019	

Labeler - Medical Purchasing Solutions, LLC (601458529)

Establishment

Name	Address	ID/FEI	Business Operations
Medical Purchasing Solutions, LLC		601458529	repack(71872-7310)

Revised: 11/2023

Medical Purchasing Solutions, LLC