

VIRCIN ADVANCED WART TREATMENT- salicylic acid cream
Pure Source, LLC

VIRCIN Advanced wart treatment

Active Ingredient

Salicylic acid 17%; in collodion; contains alcohol

Purpose

Wart Remover

Uses:

For the removal of plantar warts on the bottom of the foot. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness and the interruption of the footprint pattern. For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance of the surface.

Warnings

on irritated skin, infected or reddened areas, moles, birthmarks, warts with hair growing from them, genital warts, warts on the face or mucous membrane. diabetes or poor blood circulation. Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes. Stop use and ask a doctor if discomfort persists. Cap tube tightly and store at room temperature away from heat. **For external use only. Do not use. Ask a doctor before use if you have. Avoid inhaling vapors. Flammable. Keep away from fire and flame.**

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions:

Wash affected area, may soak in warm water for 5 minutes. Dry area thoroughly. By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

Inactive Ingredients:

BHT, Ethyl Pyruvate, Glycerin, Povidone Iodine, Propylene Glycol, Transcutol, Triethanolamine

Package Labeling:



Drug Facts

Active ingredient: Salicylic acid 17%; in colloidon.....Wart remover

Purpose

Uses: For the removal of plantar warts on the bottom of the foot. The plantar wart is recognized by its location only on the bottom of the foot, its tenderness and the interruption of the footprint pattern. For the removal of common warts. The common wart is easily recognized by the rough "cauliflower-like" appearance of the surface.

Warnings: For external use only. Do not use on irritated skin, infected or reddened areas, moles, birthmarks, warts with hair growing from them, genital warts, warts on the face or mucous membrane. **Ask a doctor before use if you have diabetes or poor blood circulation.** Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes. **Avoid inhaling vapors.** Stop use and ask a doctor if discomfort persists. **Flammable. Keep away from fire and flame.** Cap tube tightly and store at room temperature away from heat. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions: Wash affected area, may soak in warm water for 5 minutes. Dry area thoroughly. By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

Inactive ingredients: Butylated Hydroxytoluene (BHT), Ethyl Pyruvate, Glycerin, Hydroxyethylcellulose, Isopropyl Alcohol, Methylsulfonylmethane (MSM), Povidone Iodine, Propylene Glycol, Retinyl Palmitate (Vitamin A), Triethanolamine, Urea

DISPENSE ONLY BY PHYSICIANS



VIRCIN

Advanced wart treatment

Net Weight 0.5 oz / 14 gm



DIRECTIONS FOR USE: Wash affected area, may soak in warm water for 5 minutes. Dry area thoroughly. By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

Distributed by Pedicis Research LLC, Radnor, PA 19087. Made in the USA
Questions: 1-800-748-6539

Patent Pending

DIRECTIONS FOR USE: By squeezing tube gently, apply one drop at a time to sufficiently cover each wart. Let dry. Repeat this procedure twice daily as needed (until wart is removed) for up to 12 weeks or as directed by a doctor.

ACTIVE INGREDIENT: Salicylic acid 17%; in collodion; contains alcohol. **INACTIVE INGREDIENTS:** See Carton

WARNINGS: For external use only. Do not use on irritated skin, infected or reddened areas, moles, birthmarks, warts with hair growing from them, genital warts, warts on the face or mucous membrane.


Ask a doctor before use if you have diabetes or poor blood circulation. Avoid contact with eyes. If product gets into the eye, flush with water for 15 minutes.

Flammable. Keep away from fire and flame. Cap tube tightly and store at room temperature away from heat. **Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center.

Distributed by Pedicis Research LLC, Radnor, PA 19087. Made in the USA

Questions: 1-800-748-6539

Patent Pending



VIRCIN ADVANCED WART TREATMENT

salicylic acid cream

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:65121-397
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	17 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
ETHYL PYRUVATE (UNII: 03O98E01OB)	
GLYCERIN (UNII: PDC6A3C0OX)	
POVIDONE-IODINE (UNII: 85H0HZU99M)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

TROLAMINE (UNII: 903K93S3TK)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65121-397-15	1 in 1 CARTON	02/10/2017	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	06/12/2014	

Labeler - Pure Source, LLC (080354456)

Establishment

Name	Address	ID/FEI	Business Operations
Pure Source, LLC		080354456	manufacture(65121-397)

Revised: 12/2024

Pure Source, LLC