

**FOSTER AND THRIVE CHILDRENS PAIN AND FEVER-  
acetaminophen suspension  
Strategic Sourcing Services LLC**

-----  
**McKesson Children's Pain & Fever Drug Facts**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily:

- reduces fever
- relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has**

liver disease

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin

**When using this product**

**do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed (see overdose warning)**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child’s dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)*</b>
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL

60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

### Other information

- **each 5 mL contains:** sodium 3 mg
- store at 20-25°C (68-77°F)
- **do not use if printed neckband is broken or missing**

### Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

### Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

### Principal Display Panel

COMPARE TO CHILDREN'S TYLENOL® ORAL SUSPENSION ACTIVE INGREDIENT

Foster & Thrive™

Children's Pain & Fever

Acetaminophen 160 mg per 5 mL

PAIN RELIEVER/FEVER REDUCER

Oral Suspension

Fast, Effective Relief Of Children's Fever And Pain


- Paraben Free
- Aspirin Free
- Ibuprofen Free
- Alcohol Free
- Gluten Free

Ages 2 to 11

BUBBLE GUM FLAVOR

DOSING CUP INCLUDED

4 FL OZ (118 mL)



**Drug Facts**  
**Active ingredient** *Purpose*  
**(in each 5 mL)**  
Acetaminophen 160 mg.....Pain reliever/fever reducer

**Uses** temporarily:  
■ reduces fever  
■ relieves minor aches and pains due to:  
■ the common cold ■ flu ■ headache  
■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if your child has ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if your child has liver disease**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**  
■ pain gets worse or lasts more than 5 days  
■ fever gets worse or lasts more than 3 days  
■ new symptoms occur  
■ redness or swelling is present  
These could be signs of a serious condition.

**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms. ▶

**Drug Facts (continued)**  
**Directions**  
■ this product does not contain directions or complete warnings for adult use  
■ do not give more than directed (see overdose warning)  
■ shake well before using  
■ mL = milliliter  
■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.  
■ remove the child protective cap and squeeze your child's dose into the dosing cup  
■ repeat dose every 4 hours while symptoms last  
■ do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*or as directed by a doctor  
**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information**  
■ each 5 mL contains: sodium 3 mg  
■ store at 20-25°C (68-77°F)  
■ do not use if printed neckband is broken or missing

**Inactive ingredients**  
anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C red #40, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

**Questions or comments?** Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

©2023 McKesson Corporation  
Distributed by: McKesson Corp.,  
via Strategic Sourcing Services LLC.  
Memphis, TN 38141  
Money Back Guarantee  
www.fosterandthrive.com

REV 06/23

Foster & Thrive™  
**Children's**  
**Pain & Fever**  
Acetaminophen  
160 mg per 5 mL  
PAIN RELIEVER/  
FEVER REDUCER

NDC 70677-1142-1 COMPARE TO CHILDREN'S  
TYLENOL® ORAL SUSPENSION  
ACTIVE INGREDIENT\*\*

Foster & Thrive™  
**Children's**  
**Pain & Fever**  
Acetaminophen  
160 mg per 5 mL  
PAIN RELIEVER/  
FEVER REDUCER

Oral Suspension  
Fast, Effective Relief Of  
Children's Fever And Pain

- Paraben Free
- Aspirin Free
- Ibuprofen Free
- Alcohol Free
- Gluten Free

 Ages 2 to 11  
**BUBBLE GUM  
FLAVOR**  
DOSING CUP INCLUDED

**4 FL OZ (118 mL)**

**DIRECTIONS:**  
**STEP 1.** Shake well before using  


**STEP 2.** Remove child protective cap  


**STEP 3.** Squeeze your child's dose into the dosing cup  


**IMPORTANT:**  
Please see dosing chart or ask a doctor for your child's correct dose.

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

\*\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.



0 10939 95803 7

10520 KM C1

## FOSTER AND THRIVE CHILDRENS PAIN AND FEVER

acetaminophen suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70677-1142
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>CALCIUM SULFATE, UNSPECIFIED FORM</b> (UNII: WAT0DDB505)	
<b>CARRAGEENAN</b> (UNII: 5C69YCD2YJ)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HIGH FRUCTOSE CORN SYRUP</b> (UNII: XY6UN3QB6S)	
<b>HYDROXYETHYL CELLULOSE, UNSPECIFIED</b> (UNII: T4V6TWG28D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SODIUM PHOSPHATE, TRIBASIC</b> (UNII: A752Q30A6X)	

**Product Characteristics**

<b>Color</b>	RED (opaque)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1142-1	1 in 1 CARTON	10/20/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/20/2023	

