



CHLORZOAZONE- chlorzoxazone tablet **Proficient Rx LP**

Chlorzoxazone Tablets, USP

DESCRIPTION

Chlorzoxazone USP is a centrally acting skeletal muscle relaxant, available as tablets of 500 mg for oral administration. Its chemical name is 5-Chloro-2-benzoxazolinone, and its structural formula is:



Chlorzoxazone, USP is a white or practically white, practically odorless, crystalline powder.

Chlorzoxazone is slightly soluble in water; sparingly soluble in alcohol, in isopropyl alcohol, and in methanol; soluble in solutions of alkali hydroxides and ammonia.

Chlorzoxazone tablets contain the inactive ingredients -Lactose monohydrate USNF, Microcrystalline cellulose USNF, FD&C Yellow 6/Sunset Yellow FCF Aluminium lake, D&C Red 27/Phloxine Aluminium lake, Hypromellose USP (a)

Binder solution

Hypromellose USP (b),

Polysorbate 80 USNF,

Ethanol (96%v/v).

Extragranular

Pregelatinized Starch USNF,

Sodium starch glycolate USNF,

Magnesium stearate USNF.

FDA approved dissolution method differs from that of the USP.

CLINICAL PHARMACOLOGY

Chlorzoxazone is a centrally-acting agent for painful musculoskeletal conditions. Data available from animal experiments as well as human study indicate that chlorzoxazone acts primarily at the level of the spinal cord and subcortical areas of the brain where it inhibits multisynaptic reflex arcs involved in producing and maintaining skeletal muscle spasm of varied etiology. The clinical result is a reduction of the skeletal muscle spasm with relief of pain and increased mobility of the involved muscles. Blood levels of chlorzoxazone can be detected in people during the first 30 minutes and peak levels may be reached, in the majority of the subjects, in about 1 to 2 hours after oral administration of chlorzoxazone.

Chlorzoxazone is rapidly metabolized and is excreted in the urine, primarily in a conjugated form as the glucuronide. Less than one percent of a dose of chlorzoxazone is excreted unchanged in the urine in 24 hours.

INDICATIONS

Chlorzoxazone is indicated as an adjunct to rest, physical therapy, and other measures for the relief of discomfort associated with acute, painful musculoskeletal conditions. The mode of action of this drug has not been clearly identified, but may be related to its sedative properties. Chlorzoxazone does not directly relax tense skeletal muscles in man.

CONTRAINDICATIONS

Chlorzoxazone is contraindicated in patients with known intolerance to the drug.

WARNINGS

Serious (including fatal) hepatocellular toxicity has been reported rarely in patients receiving chlorzoxazone. The mechanism is unknown but appears to be idiosyncratic and unpredictable. Factors predisposing patients to this rare event are not known. Patients should be instructed to report early signs and/or symptoms of hepatotoxicity such as fever, rash, anorexia, nausea, vomiting, fatigue, right upper quadrant pain, dark urine, or jaundice. Chlorzoxazone should be discontinued immediately and a physician consulted if any of these signs or symptoms develop. Chlorzoxazone use should also be discontinued if a patient develops abnormal liver enzymes (e.g., AST, ALT, alkaline phosphatase and bilirubin.)

The concomitant use of alcohol or other central nervous system depressants may have an additive effect.

Usage in Pregnancy

The safe use of chlorzoxazone has not been established with respect to the possible adverse effects upon fetal development. Therefore, it should be used in women of childbearing potential only when, in the judgment of the physician, the potential benefits outweigh the possible risks.

PRECAUTIONS

Chlorzoxazone should be used with caution in patients with known allergies or with a history of allergic reactions to drugs. If a sensitivity reaction occurs such as urticaria, redness, or itching of the skin, the drug should be stopped.

If any symptoms suggestive of liver dysfunction are observed, the drug should be discontinued.

ADVERSE REACTIONS

Chlorzoxazone containing products are usually well tolerated. It is possible in rare instances that chlorzoxazone may have been associated with gastrointestinal bleeding. Drowsiness, dizziness, lightheadedness, malaise, or over-stimulation may be noted by an occasional patient. Rarely, allergic type skin rashes, petechiae, or ecchymoses may develop during treatment. Angioneurotic edema or anaphylactic reactions are extremely rare. There is no evidence that the drug will cause renal damage.

Rarely, a patient may note discoloration of the urine resulting from a phenolic metabolite of chlorzoxazone. This finding is of no known clinical significance.

To report SUSPECTED ADVERSE EVENTS, contact Aurobindo Pharma USA, Inc. at 1-866-850-2876 or FDA at 1-800- FDA-1088 or <http://www.fda.gov/> for voluntary reporting of adverse reactions.

OVERDOSAGE

Symptoms

Initially, gastrointestinal disturbances such as nausea, vomiting, or diarrhea together with drowsiness, dizziness, lightheadedness or headache may occur. Early in the course there may be malaise or sluggishness followed by marked loss of muscle tone, making voluntary movement impossible. The deep tendon reflexes may be decreased or absent. The sensorium remains intact, and there is no peripheral loss of sensation. Respiratory depression may occur with rapid, irregular respiration and intercostal and substernal retraction. The blood pressure is lowered, but shock has not been observed.

Treatment

Gastric lavage or induction of emesis should be carried out, followed by administration of activated charcoal. Thereafter, treatment is entirely supportive. If respirations are depressed, oxygen and artificial respiration should be employed and a patent airway assured by use of an oropharyngeal airway or endotracheal tube. Hypotension may be counteracted by use of dextran, plasma, concentrated albumin or a vasopressor agent such as norepinephrine. Cholinergic drugs or analeptic drugs are of no value and should not be used.

DOSAGE AND ADMINISTRATION

Usual Adult Dosage

One tablet three or four times daily. If adequate response is not obtained with this dose, it may be increased to one and one-half tablets (750 mg) three or four times daily. As

improvement occurs dosage can usually be reduced.

HOW SUPPLIED

Chlorzoxazone tablets, USP 500 mg are available as Light orange to orange round shaped, flat hexagonal, beveled debossed with “N” on one side of break line and “500” on other side of break line and plain on other side.

Bottles of 20 NDC 71205-711-20

Bottles of 30 NDC 71205-711-30

Bottles of 60 NDC 71205-711-60

Bottles of 90 NDC 71205-711-90

Store at controlled room temperature 15°- 30°C (59°-86° F).

Dispense contents with a child-resistant closure (as required) and in a tight container as defined in the USP/NF.

Keep out of the reach of children.

Distributed by:

Aurobindo Pharma USA, Inc.

279 Princeton-Hightstown Road

East Windsor, NJ 08520

Manufactured by:

Aurobindo Pharma Limited

Hyderabad-500 032, India

Repackaged by:

Proficient Rx LP

Thousand Oaks, CA 91320

Revised: 11/2021

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 500 mg (30 Tablets Bottle)

NDC 71205-711-30

Rx only

Chlorzoxazone

Tablets, USP

500 mg

30 Tablets



Scan Here



NDC 71205-711-30

Packaged By: Proficient Rx LP
Thousand Oaks, CA 91320

RX Only

Chlorzoxazone 500mg
#30 Tablets SN# MASTER
Lot #:00000 Exp:00/00/00
NDC 71205-711-30

Chlorzoxazone 500mg
#30 Tablets SN# MASTER
Lot #:00000 Exp:00/00/00
NDC 71205-711-30

Chlorzoxazone 500mg
#30 Tablets SN# MASTER
Lot #:00000 Exp:00/00/00
NDC 71205-711-30



GTIN: 00371205711306
SN# MASTER
Exp. 00/00/00
Lot #:00000



Chlorzoxazone 500mg

#30 Tablets

Each tablet contains: Chlorzoxazone, USP 500 mg

Light orange to orange round shaped, flat hexagonal, beveled debossed with "N" on one side of break line and "500" on other side of break line and plain on other side

Product ID: QC071130

Mfr. By: Aurobindo Pharma Limited Hyderabad-500 032, India

Store at controlled room temperature 15° to 30°C (59° to 86° F).

Keep medication out of the reach of children

CHLORZOAZONE

chlorzoxazone tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71205-711(NDC:59651-306)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORZOAZONE (UNII: H0DE420U8G) (CHLORZOAZONE - UNII:H0DE420U8G)	CHLORZOAZONE	500 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
ALCOHOL (UNII: 3K9958V90M)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	ORANGE (Light orange to orange)	Score	2 pieces
Shape	ROUND (flat, hexagonal)	Size	13mm
Flavor		Imprint Code	N;500
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71205-711-20	20 in 1 BOTTLE; Type 0: Not a Combination Product	03/12/2025	
2	NDC:71205-711-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/03/2022	
3	NDC:71205-711-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	11/03/2022	
4	NDC:71205-711-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	11/03/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA089853	08/03/2020	

Labeler - Proficient Rx LP (079196022)

Establishment

Name	Address	ID/FEI	Business Operations
Proficient Rx LP		079196022	REPACK(71205-711) , RELABEL(71205-711)

Revised: 3/2025

Proficient Rx LP