

**ADVENTURE MEDICAL KITS 1-2 PERSON FIRST AID - benzalkonium chloride, aspirin, ibuprofen, isopropyl alcohol, benzocaine, sd alcohol, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, acetaminophen, diphenhydramine chloride
Tender Corp dba Adventure Medical Kits**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ADVENTURE MEDICAL KITS 1-2 PERSON [1.0] FIRST AID KIT

Active Ingredient:Bacitracin Zinc 400 units

Neomycin Sulfate 5mg (equivalent to 3.5 mg Neomycin base)

Polymyxin B Sulfate 5000 units

Uses: To help prevent infection in:
minor cuts; scrapes; burns

Warnings:

For external use only.

Do not use: in eyes; over large areas of the body;

If allergic to any of the ingredients; for more than one week unless directed by a physician.

Stop use and consult a doctor:

if the condition persists or gets worse; a rash or other allergic reaction develops

Keep out of reach of children.

If ingested, contact a Poison

Control Center right away.

Directions: clean affected area; apply small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily; may be covered with a sterile bandage

Other information:

Store at room temperature.

Genuine Triple Antibiotic

First Aid Ointment

To Help Prevent Infection

Each Gram Contains:

Bacitracin Zinc 400 units

Neomycin Sulfate 5 mg

(equivalent to 3.5 mg

Neomycin base)

Polymyxin B Sulfate 5000 units

Net Wt. 0.5g ; (1/64 oz)

Manufactured in CHINA for

GENUINE FIRST AID.

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: shock, facial swelling, asthma (wheezing) rash, skin reddening, blisters, hives. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug, take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have a high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor before use if you are:
taking any other drug containing NSAID (prescription or nonprescription); taking aspirin for heart attack or stroke, because Ibuprofen may decrease this benefit of aspirin; taking any other drug

When using this product:
take with food or milk if stomach upset occurs

Stop use and ask a doctor If:
you experience any of the following signs of stomach bleeding; feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
do not use more than directed; the smallest effective dose should be used; do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information: Store at controlled room temperature; avoid excessive heat 40 degree Celsius (104 degree Fahrenheit); tamper evident sealed packets; do not use any opened or torn packets

Inactive ingredients: cellulose, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, polydextrose, polyethylene glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

IBUPROFEN 2 Tablets

IBUPROFEN 2 Tablets

Active Ingredient (in each tablet)	Purpose
Aspirin (NSAID*) 325 mg.....	Pain Reliever / fever reducer

*nonsteroidal anti-inflammatory drug

Uses Temporarily relieves minor aches and pains associated with:
headache ; muscular aches ; minor arthritis pain ; backache ; common cold ; toothache ; menstrual cramps
; Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock, If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product; take more or for a longer time than directed

Do not use: if you have ever had an allergic reaction to any other pain reliever/ fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis, or kidney disease; you are taking a diuretic

Ask a doctor or pharmacist before use if you are:

under a doctor's care for any serious condition; taking any other drug

When using this product: take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

feel faint; vomit blood; have bloody or black stools; have stomach pain that does not get better; pain gets worse or lasts more than 10 days; fever gets worse or lasts more than 3 days; you have difficulty swallowing; if ringing in the ears or loss of hearing occurs; redness or swelling is present in the painful areas; any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not use more than directed
the smallest effective dose should be used
drink a full glass of water with each dose
do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); avoid excessive heat and humidity; tamper evident sealed packets;

Do not use any opened or torn packets

Inactive Ingredients: hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

ASPIRIN 2 Tablets

Active Ingredient (in each tablet)	Purpose
Acetaminophen 500 mg.....	Pain Reliever / fever reducer

Purpose: Pain reliever, fever reducer

Uses for the temporary relief of minor aches and pains associated with headache ; muscular aches ; minor arthritis pain ; toothache ; common cold ; menstrual cramps ; for the reduction of fever

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 8 tablets in 24 hours, which is the maximum daily amount; with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using this product.

Do not use: with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

for more than 10 days for pain unless directed by a doctor
for more than 3 days for fever unless directed by a doctor

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop using and ask a doctor if
symptoms do not improve
new symptoms occur
pain or fever persists or gets worse
redness or swelling is present

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if

you do not notice any signs or symptoms

If pregnant or breast-feeding, ask a health professional before use.

Adults and children: (12 years and older)

take 2 tablets every 4 to 6 hours as needed. Do not take more than 8 tablets in 24 hours.

Children under 12 years:

do not give to children under 12 years of age.

Store at 59 - 86 degree Fahrenheit (15 - 30 degree Celsius); Tamper-evident sealed packets. do not use any opened or torn packets Mfd. for MEDIQUE PRODUCTS, FORT MYERS, FL 33967

Inactive Ingredients: Cellulose*, corn starch*, crospovidone*, hydroxypropyl cellulose*, hypromellose*, magnesium stearate*, microcrystalline cellulose*, mineral oil*, opadry clear*, polyethylene glycol*, polyvinylpyrrolidone*, povidone*, pregelatinized starch*, propylene glycol*, silicon dioxide*, sodium carboxymethylcellulose*, sodium starch glycolate*, starch 1500*, stearic acid, talc*, titanium dioxide*, triacetin*.

Active Ingredient (in each tablet)	Purpose
Diphenhydramine Hydrochloride 25mg.....	Antihistimine

Purpose: Antihistimine

Uses Temporarily relieves the following symptoms associated with hay fever or other upper respiratory allergies:

runny nose, sneezing, itching of the nose or throat, itchy, watery eyes

Store at room temperature. Tamper-evident sealed packets. do not use any opened or torn packets Mfd. for MEDIQUE PRODUCTS, FORT MYERS, FL 33967

Warnings

Ask a doctor before use

if you have:

a breathing problem such as emphysema or chronic bronchitis

glaucoma

difficulty in urination due to enlargement of the prostate gland

if you are:

taking any drugs for asthma

taking sedatives or tranquilizers

When using this product

Drowsiness may occur

avoid alcoholic beverages

alcohol, sedatives, and tranquilizers may increase the drowsiness effect

use caution when driving a motor vehicle or operating machinery

excitability may occur, especially in children

Do not exceed recommended dosage. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or poison control center immediately. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.

Adults and children: (12 years and older)

take 1 capsule every 4 to 6 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years:

do not give to children under 12 years of age unless directed by a doctor.

Inactive Ingredients: DandC Red 28, FDandC Blue 1, FDandC Red 40, gelatin, starch

Active Ingredient: Purpose

Benzalkonium Chloride 0.40%..... First Aid Antiseptic

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warning: For external use only.

Keep out of reach of children: If swallowed, get medical help or contact a Poison Control Center right away.

Stop use if unusual redness, swelling or other symptoms occur. Consult a physician immediately.

Do not use in the eyes or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive Ingredient: Purified water

LOT/EXP: Made in CHINA

20130301

Antiseptic Towelette

Genuine First Aid LLC, Clearwater FL 33755

www.GenuineFirstAid.com

1/pouch

GENUINE FIRST AID



GENUINE FIRST AID®



ISO 9001:2015



Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com



REORDER AST-001

DRUG FACTS - Antiseptic Towelette

Active Ingredient: Benzalkonium Chloride 0.40%..First Aid Antiseptic

Purpose:

Use: For Professional and Hospital use. Helps prevent infection. Antiseptic cleansing of face, hands and body without soap and water.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If unusual redness, swelling or other symptoms occur, consult a physician immediately.

Do not use: In the eyes, or over large areas of the body.

Directions: Tear open packet, unfold towelette and use to cleanse desired skin area. Discard towelette appropriately after single use.

Inactive ingredient: Purified water.

--- TEAR HERE ---

Made in CHINA



GENUINE FIRST AID®



ISO 9001:2010



Genuine First Aid LLC, Clearwater FL 33755
www.GenuineFirstAid.com

**Insect Sting
Relief Pad**

**Toallitas para
Picaduras de Insectos**

REORDER ISRP-001

DRUG FACTS - Insect Sting Relief Pad

Active Ingredient: Benzocaine, 6% w/v Topical Anesthetic

Purpose: Topical Anesthetic

SD alcohol, 60% w/v Antiseptic

Use: For the temporary relief of pain and itching associated with minor burns, scrapes and insect bites.

Warnings: For external use only. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Flammable - keep away from fire or flame. Avoid contact with eyes. If this happens, rinse thoroughly with water.

Do not use: In eyes, on broken skin, deep puncture wounds. If unusual redness, swelling, irritation or other symptoms occur, consult a physician immediately.

--- TEAR HERE ---

Made in CHINA

Reorder TAO-001

Genuine Triple Antibiotic

**First Aid Ointment
To Help Prevent Infection**

Each Gram Contains:
Bacitracin Zinc 400 units
Neomycin Sulfate 5 mg
(equivalent to 3.5 mg
Neomycin base)
Polymyxin B Sulfate 5000 units
Net Wt 0.5g · (1/64 oz)



Manufactured in CHINA for
GENUINE FIRST AID®

Drug Facts

Uses: To help prevent infection
in • minor cuts • scrapes • burns

Warnings:

For external use only

Do not use • in eyes • over large
areas of the body • if allergic to
any of the ingredients • for more
than one week unless directed
by a physician

Stop use and consult a doctor

• if the condition persists or gets
worse • a rash or other allergic
reaction develops

Keep out of reach of children.

If ingested, contact a Poison
Control Center right away.

Directions: • clean affected area
• apply small amount of this product
(an amount equal to the surface
area of the tip of a finger) on the
area 1 to 3 times daily • may be
covered with a sterile bandage

Other information:

Store at room temperature

LOT 600609 2015/01

IBUPROFEN 2 Tablets

Active ingredient (in each tablet) Purpose
Ibuprofen USP (NSAID*) 200mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:
■ the common cold ■ headache ■ toothache ■ muscular aches
■ backache ■ minor pain of arthritis ■ menstrual cramps
temporarily reduces fever

Warnings **Allergy alert:** Ibuprofen may cause a severe
allergic reaction, especially in people allergic to aspirin. Symptoms
may include: ■ shock ■ facial swelling ■ asthma (wheezing)
■ rash ■ skin reddening ■ blisters ■ hives If an allergic reaction
occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal
anti-inflammatory drug (NSAID), which may cause severe stomach
bleeding. The chance is higher if you: ■ are age 60 or older
■ have had stomach ulcers or bleeding problems ■ take a blood
thinner (anticoagulant) or steroid drug ■ take other drugs containing
NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more
alcoholic drinks every day while using this product ■ take more or
for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other
pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ stomach bleeding warning applies to
you ■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or
kidney disease ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking any
other drug containing an NSAID (prescription or nonprescription) ▶

Warnings (continued)

■ taking aspirin for heart attack or stroke, because ibuprofen may
decrease this benefit of aspirin ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the
following signs of stomach bleeding: ■ feel faint ■ vomit blood
■ have bloody or black stools ■ have stomach pain that does not
get better ■ pain gets worse or lasts more than 10 days ■ fever
gets worse or lasts more than 3 days ■ redness or swelling is present
in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It
is especially important not to use ibuprofen during the last 3 months of
pregnancy unless definitely directed to do so by a doctor because it may
cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical
help or contact a Poison Control Center right away.

Directions ■ do not use more than directed ■ the smallest
effective dose should be used ■ do not take longer than 10 days,
unless directed by a doctor

Adults and Children (12 years and older): Take 1 tablet every 4 to 6
hours while symptoms persist. If pain or fever does not respond to 1
tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours,
unless directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information ■ store at controlled room temperature ■ avoid
excessive heat 40° C (104° F) ■ tamper evident sealed packets ■ do not
use any opened or torn packets

Inactive ingredients cellulose, corn starch, fumed silica gel,
hypromellose, lactose, magnesium stearate, polydextrose, polyethylene
glycol, povidone, silica, sodium starch glycolate, stearic acid, titanium
dioxide, triacetin

Distributed by GENUINE FIRST AID
600 Cleveland Str Suite 400, Clearwater, FL 33755

ASPIRIN

2 Tablets

Active ingredient (in each tablet)

Purpose

Aspirin (NSAID*) 325 mg Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- backache
- common cold
- toothache
- menstrual cramps

Temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ skin reddening ■ facial swelling ■ rash ■ asthma (wheezing) ■ blisters ■ shock

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinner (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/ fever reducer ■ right before or after heart surgery ■ if you are taking a prescription drug for gout, diabetes or arthritis

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic

Warnings (continued)

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not use more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- do not take longer than 10 days, unless directed by a doctor

Adults and children: (12 years and older) Take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Other information

- store at 59°-86°F (15°-30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients hypromellose, polyethylene glycol, propylene glycol, corn starch

Distributed by **GENUINE FIRST AID**
600 Cleveland Str Suite 400, Clearwater, FL 33755

Warnings (continued)**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor.
- for more than 3 days for fever unless directed by a doctor.

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

If pregnant or breast-feeding, ask a health professional before use.

Directions ■ do not use more than directed

Adults and children: (12 years and older)

Take 2 tablets every 4 to 6 hours as needed. Do not take more than 8 tablets in 24 hours.

Children under 12 years:

Do not give to children under 12 years of age.

SEE CARTON FOR COMPLETE PRODUCT INFORMATION

Lot 1857
Exp. 01-2013

Médique[®]

2 Tablets

Made in
USA

Extra Strength[®]

APAP

Active Ingredient (in each tablet) Purpose
Acetaminophen 500 mg. Pain reliever/fever reducer

Uses

For the temporary relief of minor aches and pains associated with

- headache
- muscular aches
- minor arthritis pain
- toothache
- common cold
- menstrual cramps

For the reduction of fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Store at 59°-86°F (15°-30°C) • Tamper-evident sealed packets • Do not use any opened or torn packets
Mfg. for **MÉDIQUE PRODUCTS** • Fort Myers, FL 33967

Medique[®]

NDG# 47682-182-46

Made in
USA

Diphen

1 Capsule

Active Ingredient

Purpose

(In each capsule)

Diphenhydramine

Hydrochloride 25mgAntihistamine

Uses:

Temporarily relieves the following symptoms associated with hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes

Store at room temperature • Tamper-Evident Sealed
Packets • Do Not Use any Opened or Torn Packets

Mfd. for **MEDIQUE PRODUCTS**
Fort Myers, FL 33967

Warnings

Ask a Doctor Before Use

If you have:

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

If you are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using this product:

- drowsiness may occur
- avoid alcohol beverages
- alcohol, sedatives and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children.

Do not exceed recommended dosage. Keep this and all drugs out of the reach of children. In case of accidental overdose, contact a physician or poison control center immediately. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.

Directions: Do not use more than directed

Adults and children: (12 years and older) Take 1 capsule every 4 to 6 hours as needed. Do not take more than 12 capsules in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age unless directed by a doctor.

Lot 3951 Exp. 11-2011



1-2 PEOPLE

FIRST
[1.0] AID

MEDICAL SUPPLIES & SURVIVAL TOOLS

As
Med
21
855



ADVENTURE MEDICAL KITS 1-2 PERSON FIRST AID

benzalkonium chloride, aspirin, ibuprofen, isopropyl alcohol, benzocaine, sd alcohol, bacitracin zinc, neomycin sulfate, polymyxin b sulfate, acetaminophen, diphenhydramine chloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:44224-2000
---------------------	----------------	---------------------------	----------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:44224-2000-1	1 in 1 KIT		

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 TUBE	1 g
Part 2	2 PACKAGE	1 mL
Part 3	4 PACKET	8
Part 4	2 PACKET	2
Part 5	3 PACKET	6
Part 6	2 PACKAGE	4

Part 7 10 PACKAGE

8 mL

Part 1 of 7

GENUINE TRIPLE ANTIBIOTIC

bacitracin zinc,neomycin sulfate,polymyxin b sulfate ointment

Product Information

Item Code (Source) NDC:52124-0003

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN ZINC	400 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	5 mg in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B SULFATE	5000 [iU] in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0003-1	.5 g in 1 TUBE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	08/26/2010	

Part 2 of 7

INSECT STING RELIEF PAD

benzocaine,alcohol swab

Product Information

Item Code (Source) NDC:52124-0008

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	6 mL in 100 mL
ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	60 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0008-1	0.5 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part348	08/26/2010	

Part 3 of 7**IBUPROFEN**

ibuprofen tablet

Product Information

Item Code (Source)	NDC:52124-0009
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE (UNII: J2B2A4N98G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (WHITE)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;352
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0009-1	2 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	08/26/2010	

Part 4 of 7

MEDIQUE DIPHEN

diphenhydramine hydrochloride capsule

Product Information

Item Code (Source)	NDC:47682-182
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
GELATIN (UNII: 2G86QN327L)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	

FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	pink (PINK) , white (WHITE)	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	CPC;835
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-182-46	1 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/25/2010	

Part 5 of 7

MEDIQUE APAP EXTRA STRENGTH

acetaminophen tablet, film coated

Product Information

Item Code (Source)	NDC:47682-175
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
CROSPVIDONE (UNII: 68401960MK)	
HYDROXYPROPYL CELLULOSE (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	

POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	white (WHITE)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	AZ;235
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-175-46	2 in 1 PACKET		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/26/2010	

Part 6 of 7

ASPIRIN

aspirin tablet

Product Information

Item Code (Source)	NDC:52124-0011
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	

POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

STARCH, CORN (UNII: O8232NY3SJ)

Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52124-0011-1	2 in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	08/26/2010	

Part 7 of 7

ANTISEPTIC TOWELETTE

benzalkonium chloride swab

Product Information

Item Code (Source)	NDC:52124-0001
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.40 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:52124-0001-1	0.8 mL in 1 PACKAGE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part333E	09/09/2010		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part333B	08/26/2010		

Labeler - Tender Corp dba Adventure Medical Kits (064437304)

Registrant - GFA Production (Xiamen) Co., Ltd. (421256261)

Establishment			
Name	Address	ID/FEI	Business Operations
GFA Production (Xiamen) Co., Ltd.		421256261	manufacture

Revised: 1/2011

Tender Corp dba Adventure Medical Kits