

## **CHILDRENS PAIN AND FEVER- acetaminophen suspension WALGREENS**

-----  
**Walgreens Acetaminophen 160 mg Liquid Suspension - Children's Cherry Flavor**

### ***Drug Facts***

#### ***Active ingredient (in each 5 mL)***

Acetaminophen 160 mg

#### ***Purpose***

Pain reliever/fever reducer

#### ***Uses*** temporarily:

reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

#### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### ***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if your child has** liver disease

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

**Do not take more than directed.**

See overdose warning

- **this product does not contain directions or complete warnings for adult use.**
- **shake well before using**
- mL = milliliter
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

---

Weight (lbs)	Age (yr)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

---

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

### ***Other information***

- **each 5 mL contains:** sodium 2 mg
- store between 20-25° (68-77°F)

## Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum

**Questions or comments? Call 1-877-290-4008**

**Drug Facts (continued)**

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL


\*or as directed by a doctor  
**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other Information** ■ each 5 mL contains sodium 2 mg ■ store between 20°-25° (68°-77°F)  
**Inactive ingredients** anhydrous citric acid, FD&C red no. 40, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum


**Questions or comments?** Call 1-877-290-4008

**IMPORTANT:**  
 Please see dosing chart or ask a doctor for your child's correct dose. **DOSAGE CUP PROVIDED**


**STEP 1.**  
Shake well before using




**STEP 2.**  
Remove child protective cap



**STEP 3.**  
Squeeze your child's dose into the dosing cup



Walgreens's Pharmacist Recommended.  
 Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.



how2recycleinto  
PLASTIC BOTTLE

Walgreens

Children's  
Pain & Fever


ACETAMINOPHEN 160 mg PER 5 mL

PAIN RELIEVER / FEVER REDUCER  
ORAL SUSPENSION

Aspirin Free
Alcohol Free

Ibuprofen Free

AGES 2-11 YEARS



Cherry  
flavor

4 FL OZ (118 mL)

TAMPER EVIDENT: DO NOT USE IF SHRINK BAND SEAL AROUND THE BOTTLE NECK IS BROKEN OR MISSING

Drug Facts

**Active ingredient (in each 5 mL)**  
 Acetaminophen 160 mg Pain reliever/fever reducer

**Uses** temporarily:  
 ■ relieves fever  
 ■ relieves minor aches and pains, due to:  
 ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product**

**Ask a doctor before use if your child has liver disease**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin when using this product. Do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if:** ■ pain gets worse or lasts more than 3 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present ■ these could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** Cap of overdose gel medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** Do not take more than directed. See overdose warning


**before using** ■ this product does not contain directions or complete warnings for adult use. ■ shake well ■ mL = milliliter ■ Find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ Remove the child protective cap and squeeze your child's dose into the dosing cup ■ repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours

Drug Facts (continued under label)

DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015  
 100% SATISFACTION GUARANTEEED  
 walgreens.com ©2023 Walgreen Co.  
 MADE IN INDIA  
 ITEM 776624 W00000-0000-0  
 758L 0224  
 NDC 0363-7014-05  
 W3ORG0324-F

LOT:

EXP:





## CHILDRENS PAIN AND FEVER

acetaminophen suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-7014
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) ACETAMINOPHEN 160 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCROSE (UNII: C151H8M554)	

### Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-7014-05	1 in 1 CARTON	04/10/2024	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/10/2024	

**Labeler** - WALGREENS (008965063)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 4/2024

WALGREENS