

DYE FREE INFANTS PAIN AND FEVER- acetaminophen suspension WALGREENS

Walgreens Acetaminophen 160mg/5 ml suspension - Infants Cherry Dye Free

Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses temporarily:

reduces fever

relieves minor aches and pains due to:

- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product, do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

this product does not contain directions or complete warnings for adult use. shake well before using

mL = milliliter

find right dose on chart below. If possible, use weight to dose; otherwise, use age.

push air out of syringe. Insert syringe tip into the bottle opening

flip bottle upside down. Pull yellow part of syringe to correct dose

dispense liquid slowly into child's mouth, toward inner cheek

repeat dose every 4 hours while symptoms last

do not give more than 5 times in 24 hours

replace cap tightly to maintain child resistance

Weight (lbs)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

*or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device

Other information

store between 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, sucrose, xanthan gum

Questions or comments? Call **1-877-290-4008**

Walgreens

Dye-Free

sin Tinturas Infants' Pain & Fever Dolor & Fiebre en Bebés

ACETAMINOPHEN 160 mg PER 5 mL ORAL SUSPENSION
ACETAMINOFÉN 160 mg POR CADA 5 ml SUSPENSIÓN ORAL
PAIN RELIEVER / FEVER REDUCER
PARA ALIVIAR EL DOLOR / REDUCIR LA FIEBRE

- USE ONLY WITH ENCLOSED SYRINGE
- Use únicamente la jeringa adjunta

AGES 2-3 YEARS

Para 2 a 3 años de edad



Cherry flavor
sabor a Cereza

2 FL OZ (60 mL)

TAMPER EVIDENT: DO NOT USE IF SHRINKRAMP SEAL AROUND THE BOTTLE NECK IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg Pain reliever/fever reducer

Uses temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, runny nose, or vomiting, consult a doctor promptly.
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
When using this product, do not exceed recommended dose (see overdose warning)
Stop use and ask a doctor if ■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed. See overdose warning
■ this product does not contain directions or complete warnings for adult use. ■ shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ push air out of syringe. Insert syringe tip into the bottle opening. ■ tip bottle upside down. Pull yellow part of syringe to correct dose. ■ dispense liquid slowly into child's mouth, toward inner cheek. ■ repeat dose every 4 hours while symptoms last ■ do not give more than 5 times in 24 hours ■ replace cap lightly to maintain child resistance

Drug Facts (continued under label)

DISTRIBUTED BY: WALGREEN CO., DISTRIBUTIDO POR: WALGREEN CO., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2023 Walgreen Co. MADE IN INDIA
ITEM 776788 W00000-0000-0 W3ORG0324-F 738L 0224 WDC 03693-7015-04

LOT:



EXP.:

Drug Facts (continued)

DOSING CHART

Weight (lbs)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL

*Or as directed by a doctor

Important: Use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device

Other information ■ store between 20-25°C (68-77°F)

Inactive ingredients ■ arylpyrazole sodium, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and croscarmellose sodium, propylparaben, purified water, sodium benzoate, sorbitol solution, sucralose, xanthan gum

Questions or comments? Call 1-877-296-4008

IMPORTANT: Please see dosing chart or ask a doctor for your child's correct dose.

STEP 1.
See Directions for the correct dose.
Shake well before using.

STEP 2.
Push air out of the syringe.

STEP 3.
Insert syringe tip into bottle opening then flip bottle upside down.

STEP 4.
Pull yellow part of syringe to the 1st dose line then push product back into bottle.

STEP 5.
Pull yellow part of syringe until it reaches and stays at the correct dose.

STEP 6.
Dispense liquid slowly into child's mouth.



Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.



DYE FREE INFANTS PAIN AND FEVER

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-7015
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
SUCROSE (UNII: C151H8M554)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
GLYCERIN (UNII: PDC6A3C0OX)	

Product Characteristics

Color	white (white to off white)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-7015-04	1 in 1 BOX	04/10/2024	
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/10/2024	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(0363-7015)

Revised: 4/2024

WALGREENS