

LOPERAMIDE HYDROCHLORIDE- loperamide hcl solution
Preferred Pharmaceuticals Inc.

Major Pharmaceuticals Loperamide Hydrochloride Oral Solution Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.
- mL = milliliter

adults and children 12 years and over	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours
children 9-11 years (60-95 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours
children 6-8 years (48-59 lbs)	15 mL after the first loose stool; 7.5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each 30 mL contains:** sodium 15 mg
- store between 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

Relabeled By: Preferred Pharmaceuticals Inc.

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
XANTHAN GUM (UNII: TTV12P4NEE)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	GREEN (opaque, viscous)	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-7842-1	120 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/19/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091292	01/19/2021	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-7842)

Revised: 7/2023

Preferred Pharmaceuticals Inc.