

**VANACOF XP COUGH / CHEST CONGESTION- dextromethorphan hbr,
guaifenesin solution
GM Pharmaceuticals, INC**

VANACOF XP Cough / Chest Congestion

Active Ingredients (in each 15 mL (1TBSP))

Dextromethorphan HBr 18 mg

Guaifenesin 396 mg

Purpose

Cough Suppressant

Expectorant

Uses

- ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- ■ temporarily relieves:

■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

■ the intensity of coughing

■ the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- ■ a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- ■ a cough that occurs with too much phlegm (mucus)

When using this product

do not use more than directed

Stop use and ask a doctor if

■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- ■ do not take more than 6 doses in any 24-hour period
- ■ dose as follows or as directed by a doctor ■ mL= milliliter

adults and children 12 years of age and over:	15 mL (1 TBSP) every 4 hours, not to exceed 90 mL (6 TBSP) per 24 hours
children 6 to under 12 years of age:	7.5 mL (1/2 TBSP) every 4 hours, not to exceed 45 mL (3 TBSP) per 24 hours
children under 6 years:	consult a doctor.

Other information

■ each 15 mL (1 TBSP) contains: Sodium 6 mg

■ store at 20° to 30°C (68° to 86°F)

Inactive ingredients

citric acid, flavors, glycerin, l-menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-888-535-0305 9 a.m. - 5 p.m. CST

Tamper evident by foil seal under cap. Do no use if foil seal is broken or missing.

Distributed by: GM Pharmaceuticals, Inc. Fort Worth, TX 76118

NDC 58809-187-08

VANACOF® XP

COUGH / CHEST CONGESTION

Each 15 mL (1 TBSP) contains:

Dextromethorphan HBr.....18 mg

Guaifenesin396 mg

Cough Suppressant ■ Expectorant
 Raspberry-Mint Flavor
 Alcohol Free / Sugar Free / Gluten Free / Dye Free

NDC 58809-187-08

VANACOF[®] XP

COUGH / CHEST CONGESTION

Each 15 mL (1 TBSP) contains:
 Dextromethorphan HBr 18 mg
 Guaifenesin 396 mg

Cough Suppressant • Expectorant

Raspberry-Mint Flavor

Alcohol Free / Sugar Free / Gluten Free / Dye Free

GM Pharmaceuticals, Inc.

8 fl oz (237 mL)

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Drug Facts

Active ingredients (in each 15 mL (1TBSP))

Dextromethorphan HBr 18 mg.....Cough Suppressant
 Guaifenesin 396 mg.....Expectorant

Purpose

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Drug Facts (continued)

Stop use and ask a doctor if

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- dose as follows or as directed by a doctor

adults and children 12 years of age and over:	15 mL (1 TBSP) every 4 hours, not to exceed 90 mL (6 TBSP) per 24 hours
children 6 to under 12 years of age:	7.5 mL (1/2 TBSP) every 4 hours, not to exceed 45 mL (3 TBSP) per 24 hours
children under 6 years:	consult a doctor.

Other information

- each 15 mL (1 TBSP) contains: Sodium 6 mg
- store at 20° to 30°C (68° to 86°F)

Inactive ingredients citric acid, flavors, glycerin, l-menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments? 1-888-535-0305 9 a.m. - 5 p.m. CST

Distributed by: GM Pharmaceuticals, Inc. Fort Worth, TX 76118



VANACOF XP COUGH / CHEST CONGESTION			
dextromethorphan hbr, guaifenesin solution			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58809-187
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)		GUAIFENESIN	396 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE	18 mg in 15 mL
Inactive Ingredients			
Ingredient Name			Strength
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
MENTHOL (UNII: L7T10EIP3A)			
XANTHAN GUM (UNII: TTV12P4NEE)			
SORBITOL (UNII: 506T60A25R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
GLYCERIN (UNII: PDC6A3COOX)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0KO0R)

Product Characteristics

Color		Score	
Shape		Size	
Flavor	RASPBERRY, MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58809-187-08	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/06/2023	

Labeler - GM Pharmaceuticals, INC (793000860)

Revised: 1/2026

GM Pharmaceuticals, INC