

**TABCIN ACTIVE- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled OPMX LLC.**

-----  
**Tabcin Active**

**Drug Facts**

<b>Active ingredients (in each Softgel)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

Temporarily relieves common cold/flu symptoms:

- Nasal congestion
- Cough due to minor throat & bronchial irritation
- Sore throat
- Headache
- Minor aches & pains
- Fever

**Warnings**

**Liver warning**

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include.

Skin reddening blisters • rash • If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
- If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- Liver disease
- Heart disease
- Diabetes
- High blood pressure
- Trouble urinating due to enlarged prostate gland
- Cough that occurs with too much phlegm (mucus)
- Persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema.

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

do not use more than directed.

**Stop use and ask a doctor if**

- You get nervous, dizzy or sleepless.
- Symptoms get worse or last more than 5 days (children) or 7 days (adults).
- Fever gets worse or lasts more than 3 days.
- Redness or swelling is present.
- New symptoms occur.
- Cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**Keep out of reach of children.**

**Keep out of reach of children.**

**If pregnant or breast-feeding,**

ask a health professional before use.

**Overdose warning**

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well for children even if you do not notice any signs or symptoms.

**Directions**

- Take only as directed.
- Do not exceed 4 doses per 24 hrs.

Adults & children 12 yrs & over	2 softgels with water every 4 hrs
Children 4 to under 12 yrs	Ask a doctor
Children under 4 yrs	Do not use

### **Other information**

Store at room temperature

### **Inactive ingredients**

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special, titanium dioxide.

### **Questions?**

Call to free 619-600-5632 Monday through Friday 9AM - 5PM EST

### **PRINCIPAL DISPLAY PANEL**

NDC 69729-152-10

tabcin Active

Non-Drowsy

COUGH & FLU / TOS Y GRIPE

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

- Pain Reliever/ Analgésico
- Fever Reducer/ Reduce la Fiebre
- Cough Suppressant / Supresor de la Tos
- Nasal Decongestant / Descongestivo Nasal

10 Softgels / 10 Cápsulas blandas de gel



# tabcin ACTIVE

# tabcin ACTIVE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

Non-Drowsy

**10** Softgels\*  
10 Cápsulas Blandas de Gel \*

**COUGH & FLU TOS Y GRIPE**  
**Multi-Symptom Relief**  
Pain Reliever / Analgésico  
Fever Reducer / Reduce la Fiebre  
Cough Suppressant / Supresor de la Tos  
Nasal Decongestant / Descongestivo Nasal

# tabcin ACTIVE

NDC 69729-152-10

Lot:  
Exp:  
REV. 11/2023 ITEM # INV030  
8 50043 1 16906 2

**Drug Facts (continued)**  
occurs with too much phlegm (mucus). • Persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema. Taking the blood thinning drug warfarin.  
**Stop use and ask a doctor if**  
• You get nervous, dizzy or sleepless. • Symptoms get worse or last more than 5 days (children) or 7 days (adults). • Fever gets worse or lasts more than 3 days. • Cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.  
**Keep out of reach of children.**  
**If pregnant or breast-feeding, ask a health professional before use.** **Overdose warning:** Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions**  
• Take only as directed. • Do not exceed 4 doses per 24 hours.  
Adults & children 12 yrs & over: 2 softgels with water every 4 hrs.  
Children 4 to under 12 yrs: Ask a doctor.  
Children under 4 yrs: Do not use.  
**Other information** • Store at room temperature.  
**Inactive ingredients:** FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide.  
**Questions?**  
Call toll free 619-600-5632, Monday through Friday 9AM - 5PM EST.

**Drug Facts**  
**Active ingredients (in each Softgel)**  
Acetaminophen 325 mg, Pain reliever/fever reducer  
Dextromethorphan HBr 10mg, Cough suppressant  
Phenylephrine HCl 5 mg, Nasal decongestant  
**Uses**  
Temporarily relieves common cold/flu symptoms:  
• Nasal congestion • Cough due to minor throat & bronchial irritation • Sore throat • Headache • Minor aches and pains • Fever  
**Warnings**  
This product contains acetaminophen. Severe liver damage may occur if you take:  
• More than 4 doses in 24 hrs. which is the maximum daily amount for this product. • With other drugs containing acetaminophen. • 3 or more alcoholic drinks daily while using this product.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin redness • Blisters • Rash  
**Severe throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, muscle aches, or if a skin reaction occurs, stop use and seek medical help right away.  
**Do not use**  
• With any other drug containing acetaminophen (prescription or nonprescription). • If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription non-narcotic cough suppressant (Mx) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the Mx) drug, if you do not know if your prescription drug contains an Mx) drug, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have**  
• Liver disease • Heart disease • Diabetes • High blood pressure • Trouble urinating due to enlarged prostate gland • Cough that

**TAMPER EVIDENT:**  
Do not use if package is damaged, tampered with, torn, broken or shows any sign of tampering.

Exclusively distributed by:  
**OPMX**  
Chula Vista, CA 91910  
Phone: 619-600-5632

Made for:  
**inovx**  
FARMACEUTIQUE

Manufactured in GMP Registered Facility in the USA

**TABCIN ACTIVE**  
acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69729-152
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	AP01
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-152-10	1 in 1 CARTON	11/28/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/28/2023	

**Labeler** - OPMX LLC. (029918743)

**Establishment**

Name	Address	ID/FEI	Business Operations
SOFTECH PHARMA PRIVATE LIMITED		677111277	manufacture(69729-152) , label(69729-152) , pack(69729-152)

Revised: 8/2025

OPMX LLC.