

TABCIN ACTIVE- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled OPMX LLC.

Tabcin Active

Drug Facts

Active ingredients (in each Softgel)	Purpose
Acetaminophen 325 mg	Pain reliever/Fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

Temporarily relieves common cold/flu symptoms:

- Nasal congestion
- Cough due to minor throat & bronchial irritation
- Sore throat
- Headache
- Minor aches & pains
- Fever

Warnings

Liver warning

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hrs, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include.

Skin reddening blisters • rash • If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
- If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- Liver disease
- Heart disease
- Diabetes
- High blood pressure
- Trouble urinating due to enlarged prostate gland
- Cough that occurs with too much phlegm (mucus)
- Persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product,

do not use more than directed.

Stop use and ask a doctor if

- You get nervous, dizzy or sleepless.
- Symptoms get worse or last more than 5 days (children) or 7 days (adults).
- Fever gets worse or lasts more than 3 days.
- Redness or swelling is present.
- New symptoms occur.
- Cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Keep out of reach of children.

If pregnant or breast-feeding,

ask a health professional before use.

Overdose warning

Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well for children even if you do not notice any signs or symptoms.

Directions

- Take only as directed.
- Do not exceed 4 doses per 24 hrs.

Adults & children 12 yrs & over	2 softgels with water every 4 hrs
Children 4 to under 12 yrs	Ask a doctor
Children under 4 yrs	Do not use

Other information

Store at room temperature

Inactive ingredients

FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol special, titanium dioxide.

Questions?

Call to free 619-600-5632 Monday through Friday 9AM - 5PM EST

PRINCIPAL DISPLAY PANEL

NDC 69729-152-10

tabcin Active

Non-Drowsy

COUGH & FLU / TOS Y GRIPE

Multi-Symptom Relief

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

- Pain Reliever/ Analgésico
- Fever Reducer/ Reduce la Fiebre
- Cough Suppressant / Supresor de la Tos
- Nasal Decongestant / Descongestivo Nasal

10 Softgels / 10 Cápsulas blandas de gel

INFORMACION FARMACOLOGICA EN ESPAÑOL - IMPRESA AL INTERIOR DE ESTA CAJA

Drug Facts (continued)

occurs with too much phlegm (mucus). • Persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis or emphysema. Taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- You get nervous, dizzy or sleepless. • Symptoms get worse or last more than 5 days (children) or 7 days (adults). • Fever gets worse or lasts more than 3 days. • Cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than directed can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- Take only as directed. • Do not exceed 4 doses per 24 hours.

Adults & children 12 yrs & over: 2 softgels with water every 4 hrs. Ask a doctor.

Children 4 to under 12 yrs: 1 do not use.

Children under 4 yrs: 1 do not use.

Other information • Store at room temperature.

Inactive ingredients: FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol, titanium dioxide.

Questions?

Call toll free 619-600-5632, Monday through Friday 9AM - 5PM EST.

Drug Facts

Active ingredients (in each Softgel)

Acetaminophen 325 mg, Dextromethorphan HBr 10mg, Phenylephrine HCl 5 mg, Pain reliever/fever reducer, Cough suppressant, Nasal decongestant.

Uses

- Temporarily relieves common cold/flu symptoms.
- Nasal congestion. • Cough due to minor throat & bronchial irritation. • Sore throat. • Headache. • Minor aches and pains. • Fever.

Warnings

- **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: • More than 4 doses in 24 hrs. which is the maximum daily amount for this product. • With other drugs containing acetaminophen. • 3 or more alcoholic drinks daily while using this product.
- **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin redness • Blisters • Rash
- **Severe throat warning:** If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, muscle aches, or if a skin reaction occurs, stop use and seek medical help right away.

Do not use

- With any other drug containing acetaminophen (prescription or nonprescription). • If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are now taking a prescription non-narcotic opioid analgesic (MMA) (certain drugs for depression, anxiety or emotional conditions) or Parkinson's disease), or for 2 weeks after stopping the MMA) drug, if you do not know if your prescription drug contains an MMA), ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- Liver disease • Heart disease • Diabetes • High blood pressure

• Trouble urinating due to enlarged prostate gland • Cough that

Manufactured in GMP Registered Facility in the USA

Made for:

inovx
FARMACEUTIQUE

Exclusively distributed by:

OPMX
Chula Vista, CA 91910
Phone: 619-600-5632

TAMPER EVIDENT:
Do not use if package is damaged, tampered with, torn, broken or shows any sign of tampering.

tabcin ACTIVE

NDC 69729-152-10

tabcin ACTIVE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl

Non-Drowsy

COUGH & FLU TOS Y GRIPE

Multi-Symptom Relief

Pain Reliever / Analgésico
Fever Reducer / Reduce la Fiebre
Cough Suppressant / Supresor de la Tos
Nasal Decongestant / Descongestivo Nasal

10 Softgels*
10 Cápsulas Blandas de Gel *

tabcin ACTIVE

Lot: _____
Exp: _____

REV. 11/2023 ITEM # INV030
8 50043 1 16906 2

TABCIN ACTIVE
acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69729-152
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	AP01
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69729-152-10	1 in 1 CARTON	11/28/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	11/28/2023	

Labeler - OPMX LLC. (029918743)

Establishment

Name	Address	ID/FEI	Business Operations
SOFTECH PHARMA PRIVATE LIMITED		677111277	manufacture(69729-152) , label(69729-152) , pack(69729-152)

Revised: 4/2024

OPMX LLC.