

WONG TO YICK EXTERNAL ANALGESIC- camphor menthol methyl salicylate oil
Wicsun Trading Inc

WTYextrnlanglscOil

Active Ingredients

Camphor 10%

Menthol 16%

Methyl Salicylate 50%

Purpose

External Analgesic

Uses

Temporarily relieves minor aches and pains of muscles and joints due to :

- simple backache
- arthritis
- strains
- bruises
- sprains

Warnings

Do not use

- on wounds
- on irritated or damaged skin
- otherwise than as directed

When using this product

- avoid contact with the eyes or mucous membranes
- do not bandage tightly

Stop use and ask a doctor if

- condition worsens,
- symptoms persist for more than 7 days
- symptoms clear up and occur again within a few days
- excessive irritation of the skin develops
- nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs

when using for pain of arthritis

- pain persists for more than 10 days
- redness is present
- in conditions affecting children under 12 years of age

Keep out of reach of children

to avoid accidental poisoning. If swallowed, get medical help or contact Poison Control Center right away

Directions

- adults and children above 3 years of age: apply 3 to 4 drops of medicated oil to the affected area 1 to 2 times daily. After applying the product, a warm pad may be placed on the affected area for 10 minutes and then massaged as shown in the illustrations.
- children under 3 years of age: consult your physician.
- refer to the above warnings; use otherwise than as directed may be dangerous.

Other Information

- store at 16 to 30°C (61 to 86°F)
- preserve in tight container

Inactive Ingredients

Lavender oil

Package Label

WONG TO YICK WOOD LOCK OINTMENT LIMITED
 黃道益活絡油有限公司
 製造商 Manufacturer
 KWAH CHUNG, N.T., HONG KONG
 Tel: (852) 2409 0920 Fax: (852) 2407 2505
 批號 Lot No.:
 失效日期 Expiry Date:
 香港 製造 MADE IN HONG KONG

黃道益活絡油
 有限公司出品



SINCE 1968 創製

黃道益活絡油 (配方II)
WONG TO YICK WOOD LOCK MEDICATED OIL

Drug Facts 藥品資料

Active ingredients 有效成分 **Purpose 用途**
 Camphor 10% 樟腦.....External analgesic外用鎮痛
 Menthol 16% 薄荷腦.....External analgesic外用鎮痛
 Methyl Salicylate 50% 水楊酸甲酯.....External analgesic外用鎮痛

Uses 作用
 Temporarily relieves minor aches and pains of muscles and joints due to:
 暫時緩解因下列症狀引起的肌肉和關節輕微疼痛：
 ■ simple backache 普通背痛 ■ arthritis 關節炎 ■ strains 損傷
 ■ bruises 瘀傷 ■ sprains 扭傷

Warnings 告誡 **For external use only 只供外用**
Do not use 在下列情況下請勿使用
 ■ on wounds 傷口
 ■ on irritated or damaged skin 過敏或破損的皮膚
 ■ otherwise than as directed 說明以外的用途

When using this product 在使用本品時
 ■ avoid contact with the eyes or mucous membranes 避免接觸眼睛或黏膜
 ■ do not bandage tightly 切勿用綁帶包紮過緊

Drug Facts (continued) 藥品資料 (續)
Stop use and ask a doctor if 列出下列情況應停止使用本品並請
 醫師注意：
 ■ symptoms worsen 情況惡化
 ■ symptoms persist for more than 7 days 症狀持續超過七天以上
 ■ symptoms clear up and occur again within a few days 症狀消失幾天後又再出現
 ■ excessive irritation of the skin develops 皮膚出現過敏
 ■ nausea, vomiting, abdominal discomfort, diarrhea, or skin rash occurs 噁心、嘔吐、腹部不適、腹瀉、或出現皮疹

Drug Facts (continued) 藥品資料 (續)
When using for pain of arthritis 用於關節炎的止痛時：
 ■ pain persists for more than 10 days 疼痛持續超過10天
 ■ in conditions affecting children under 12 years of age 影響12歲以下的兒童之情況
Keep out of reach of children to avoid accidental poisoning. If swallowed, get medical help or contact a Poison Control Center right away. 保存在兒童接觸不到的地方以免中毒。如果吞下，請立即致電或聯絡毒物控制中心。

黃道益
 活絡油 (配方II)
 HKC-17520 NDC 57597-043-01

WONG TO YICK
 WOOD LOCK MEDICATED OIL
 EXTERNAL ANALGESIC 外用鎮痛

舒筋活絡
 跌打止痛

1.7 FL OZ (50ml)
 1.7液體盎司 (50毫升)

WOOD LOCK

美國總經銷 USA Distributor
 維新貿易公司
WICSUN TRADING INC
 22 STEUBEN STREET, BROOKLYN, NY 11205, USA
 Tel: (718) 852-6844 Fax: (718) 852-6854



US-1706UB50-V4

Drug Facts (continued) 藥品資料 (續)
 ■ refer to the above warnings; use otherwise than as directed may be dangerous. 請參閱上述告誡；若不依照說明使用可能會導致危險。
Other Information 其他資訊
 ■ store at 16 to 30°C (61 to 86°F) 儲存溫度16至30°C (61至86°F)
 ■ preserve in tight container 保持容器蓋緊
Inactive Ingredients 無效成分
 ■ Lavender oil 薰衣草油
Questions Or Comments? 有關或建議?
 ■ (718) 852-6844, M-F 9am to 6pm 星期一至五，上午9時至下午6時

Drug Facts (continued) 藥品資料 (續)
Directions 用法
 ■ adults and children above 3 years of age: apply 3 to 4 drops of medicated oil to the affected area 1 to 2 times daily. After applying the product, a warm pad may be placed on the affected area for 10 minutes and then massaged as shown in the illustrations. 成人和3歲以上的兒童：塗在患處，每天塗一至兩次，每次三至四滴。塗上藥物後，可將熱敷10分鐘，並依照圖示。
 ■ children under 3 years of age: consult your physician. 3歲以下的兒童：應請教醫生。

WONG TO YICK EXTERNAL ANALGESIC			
camphor menthol methyl salicylate oil			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66897-001
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of	Strength

Ingredient Name	Strength	Strength
CAMPHOR (SYNTHETIC) (UNII: 5TJD82A1ET) (CAMPHOR (SYNTHETIC) - UNII:5TJD82A1ET)	CAMPHOR (SYNTHETIC)	1 mg in 10 mL
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	1.6 mg in 10 mL
METHYL SALICYLATE (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
LAVENDER OIL (UNII: ZBP1YXW0H8)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66897-001-50	50 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	11/15/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	11/15/2011	

Labeler - Wicsun Trading Inc (960476174)

Establishment

Name	Address	ID/FEI	Business Operations
Wong To Yick Wood Lock Ointment Ltd		664995651	manufacture(66897-001)

Revised: 10/2024

Wicsun Trading Inc