

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**Cardinal Health (Leader) 70000**

-----  
**Drug Facts**

**Active ingredient (in each capsule)**

**Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\***

**(present as the free acid and potassium salt)**

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever/fever reducer**

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - minor pain of arthritis
  - toothache
  - backache
  - the common cold
  - menstrual cramps
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- **each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

### **Inactive ingredients**

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

### **Questions or comments?**

**Call 1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

#### **Ibuprofen**

Capsules, 200 mg

Pain Reliever/Fever Reducer (**NSAID**)

SOFTGELS

(Liquid Filled Capsules)

COMPARE TO ADVIL® LIQUI-GELS® active ingredient\*\*

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

\*\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil® Liqui-Gels®.

DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

### **Product Label**

Exp. Date:

Lot No.:

Rev# 03/12  
PLD-B7E F0003787



CIN 53197 REV 2/17

NDC 70000-0197-2

# Ibuprofen

Capsules, 200 mg  
Pain Reliever / Fever Reducer  
(NSAID)

40 SOFTGELS<sup>®</sup>  
(Liquid Filled Capsules)



LEADER<sup>®</sup>

LEADER<sup>®</sup>

COMPARE TO  
ADVIL<sup>®</sup> LIQUI-GELS<sup>®</sup>  
active ingredient\*\*

100% Money Back Guarantee

© 2017 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

DISTRIBUTED BY CARDINAL HEALTH  
EUBANK DRV 4387  
www.mylider.com  
1-800-201-4387



Essential to Care<sup>®</sup> Since 2011

All LEADER<sup>™</sup> Brand Products Have A  
100% Money Back Guarantee

Return to place of purchase if not satisfied.

**Drug Facts (continued)**

**Directions**

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
  - take 1 capsule every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 capsule, 2 capsules may be used
  - do not exceed 6 capsules in 24 hours, unless directed by a doctor
  - children under 12 years: ask a doctor

**Other information**

- each capsule contains: potassium 20 mg
- read all warnings and directions before use. Keep carton.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

**Inactive ingredients** FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

**Questions or comments?**  
Call 1-877-753-3635 Monday-Friday 9AM-5PM EST

This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Advil<sup>®</sup> Liqui-Gels<sup>®</sup>.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Drug Facts (continued)**

**Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## LEADER Ibuprofen Capsules 200mg

### IBUPROFEN

ibuprofen capsule, liquid filled

#### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:70000-0197

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg	
<b>Inactive Ingredients</b>				
Ingredient Name		Strength		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
GELATIN (UNII: 2G86QN327L)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)				
WATER (UNII: 059QF0KO0R)				
SORBITOL (UNII: 506T60A25R)				
SORBITAN (UNII: 6O92ICV9RU)				
<b>Product Characteristics</b>				
Color	BLUE	Score	no score	
Shape	CAPSULE	Size	19mm	
Flavor		Imprint Code	IB200	
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0197-1	1 in 1 BOX	11/30/2016	12/31/2022
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:70000-0197-2	1 in 1 BOX	11/30/2016	12/31/2022
2		40 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA078682	11/30/2016	12/31/2022	

**Labeler** - Cardinal Health (Leader) 70000 (097537435)

Revised: 11/2019

Cardinal Health (Leader) 70000