

**DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid**  
**P & L Development, LLC**

-----

**Drug Facts**

**Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purposes**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - fever
  - nasal congestion
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if**

you are taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occurs
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs, with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause

liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see overdose warning)**
- do not exceed 4 doses in any 24 hour-period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL=milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **When using Day Time and Night Time products, carefully read each label to ensure correct dosing**

### Other information

- **each 15 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate

### Inactive ingredients

citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

### Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu\*

#### day time

#### Cold & Flu relief

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

pain reliever/fever reducer

cough suppressant

nasal decongestion

Relieves:

- aches
- fever
- sore throat
- cough
- nasal congestion

alcohol free

non-drowsy

antihistamine-free

for ages 6 years and over

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

Manufactured by:

PL Developments

11865 S. Alameda St

Lynwood, CA 90262

**Package Label**

Compare to the active ingredients in Vicks® DayQuil® Cold & Flu\*  
NDC 49580-0466-4

**ready incase**

**\$1.25**

**day time cold & flu relief**

**Acetaminophen 325 mg**

**Dextromethorphan HBr 10 mg**

**Phenylephrine HCl 5 mg**

pain reliever/fever reducer  
cough suppressant  
nasal decongestant

relieves:  
• aches  
• fever  
• sore throat  
• cough  
• nasal congestion

alcohol-free  
non-drowsy  
antihistamine-free  
for ages 6 years and over

**4 fl oz (118 mL)**

**Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg..... Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg..... Cough suppressant  
Phenylephrine HCl 5 mg..... Nasal decongestant

**Purposes**

**Uses** ■ temporarily relieves common cold and flu symptoms  
■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion ■ fever ■ cough due to minor throat and bronchial irritation

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if: ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount ■ child takes more than 4 doses (15 mL each) in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a

**Drug Facts (continued)**

drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**  
■ liver disease ■ heart disease ■ high blood pressure  
■ thyroid disease ■ diabetes  
■ trouble urinating due to an enlarged prostate gland  
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema  
■ cough that occurs with too much phlegm (mucus)

**Drug Facts (continued under label)** **PEEL HERE** →

**Drug Facts (continued)**

**Ask a doctor or pharmacist before use** if you are taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleeplessness occur ■ pain, nasal congestion, or cough gets worse, or lasts more than 5 days (children) or 7 days (adult) ■ fever gets worse, or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**if pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ do not take more than directed (see ▶)

**Drug Facts (continued)**

**Overdose warning** ■ do not take more than 4 doses in any 24-hour period ■ measure only with dosing cup provided. Do not use any other dosing device.  
■ mL = milliliter ■ keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information** ■ each 15 mL contains: sodium 12 mg ■ store between 20-25°C (68-77°F). Do not refrigerate.

**inactive ingredients** citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

**Questions or comments?** Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Drug Facts (continued)**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

Manufactured by: PL Developments 11865 S. Alameda St Lynwood, CA 90262

FLD-8380D LB009547

**PARENTS:** Learn about your medicine today. www.EasyMedicine.com

3 49580 46640 1

## READYinCASE Day Time Cold & Flu Relief

### DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride liquid

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49580-0466
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-0466-6	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2015	01/30/2027
2	NDC:49580-0466-1	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2015	01/30/2027
3	NDC:49580-0466-2	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2015	01/30/2027
4	NDC:49580-0466-4	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/30/2015	01/30/2027

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2015	01/30/2027

**Labeler** - P & L Development, LLC (101896231)

Revised: 7/2025

P & L Development, LLC