

**DAYTIME SEVERE AND NIGHTTIME SEVERE COLD AND FLU- daytime severe - acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl, nighttime severe-acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl**  
**TARGET CORPORATION**

-----  
**Target Daytime Severe and Nighttime Severe Cold and Flu Berry**

**Nighttime Severe Cold & Flu**  
***Active ingredients (in each 30 mL)***

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

***Purpose***

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

***Uses***

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 98 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients**

citric acid monohydrate, FD&C blue no.1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

**Questions or comments?**

Call **1-800-910-6874**

**Daytime Severe Cold & Flu****Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

### ***Purpose***

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

### ***Uses***

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even

if you do not notice any signs or symptoms.

## Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

## Other information

- each 15 mL contains: sodium 56 mg
- store at no greater than 25°C and do not refrigerate

## Inactive ingredients

citric acid monohydrate, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

## Questions or comments?

Call 1-800-910-6874

## Principal Display Panel

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu and NyQuil® Severe Cold & Flu Berry Liquid\*

**Maximum Strength**

**Daytime  
Nighttime  
Severe Cold & Flu**

**Acetaminophen** (Pain Reliever / Fever Reducer)  
**Dextromethorphan HBr** (Cough Suppressant)  
**Guafenesin** (Expectorant)  
**Phenylephrine HCl** (Nasal Decongestant)

- Headache, fever, sore throat, minor aches and pains
- Cough
- Chest congestion, thin and loosens mucus
- Nasal congestion, sinus pressure
- Alcohol free

**Acetaminophen** (Pain Reliever / Fever Reducer)  
**Dextromethorphan HBr** (Cough Suppressant)  
**Doxylamine succinate** (Antihistamine)  
**Phenylephrine HCl** (Nasal Decongestant)

- Headache, fever, sore throat, minor aches and pains
- Cough
- Sneezing, runny nose
- Nasal congestion, sinus pressure
- Nighttime relief
- Alcohol free

**up&up**  
Berry Flavor

12 FL OZ (354 mL)  
TOTAL 24 FL OZ (708 mL) - 2 x 12 FL OZ

12 FL OZ (354 mL)

**PARENTS:**  
Learn about rare medicine uses  
www.StopMedicineAbuse.org

**Satisfaction guaranteed - Love it or your money back.**

Dist. by Target Corporation  
Minneapolis, MN 55403  
Made in India  
TM & ©2024  
Target Brands, Inc.

NIGHTTIME SEVERE COLD & FLU		DAYTIME SEVERE COLD & FLU	
<b>Drug Facts</b>		<b>Drug Facts</b>	
<b>Active ingredients (in each 30 mL)</b>	<b>Purpose</b>	<b>Active ingredients (in each 15 mL)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/fever reducer	Acetaminophen 162.5 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant	Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine	Sucralose 250 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant	Dextromethorphan HBr 10 mg	Nasal decongestant
<b>Uses</b> Temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ chest congestion & pressure ■ cough due to minor throat & bronchial irritation ■ cough to help you sleep ■ minor aches & pains ■ headache ■ fever ■ sore throat ■ runny nose & sneezing ■ reduces swelling of nasal passages ■ temporarily restores nose breathing through the nose ■ temporarily restores nose breathing through the nose ■ promotes nasal and/or sinus drainage		<b>Uses</b> Temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ chest congestion & pressure ■ cough due to minor throat & bronchial irritation ■ minor aches & pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores nose breathing through the nose ■ temporarily restores nose breathing through the nose ■ promotes nasal and/or sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passages if betterment occurs and make coughs more productive.	
<b>Warnings</b> Line warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses (12 mL each) in 24 hrs, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 2 or more alcoholic drinks every day while using this product Always alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness/itching ■ rash ■ hives ■ skin blisters/sores. Stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, noise, or vomiting, contact a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with dry mouth/plasma (spasms) ■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble swallowing due to enlarged prostate gland ■ sodium-restricted diet Ask a doctor or pharmacist before use if you are ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin When using this product, do not use more than directed ■ excitability may occur, especially in children ■ masked depression may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ drowsiness, dizziness, and headache may increase drowsiness Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse ■ lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ rash or hives appear ■ symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If present or brand-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Check medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		<b>Warnings</b> Line warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 2 or more alcoholic drinks every day while using this product Always alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness/itching ■ rash ■ hives ■ skin blisters/sores. Stop use and seek medical help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, noise, or vomiting, contact a doctor promptly. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble swallowing due to enlarged prostate gland ■ cough that occurs with dry mouth/plasma (spasms) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ sodium-restricted diet Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product, do not use more than directed. Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days (adults) ■ fever gets worse or lasts more than 3 days (children) ■ rash or hives appear ■ symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. If present or brand-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Check medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b> Take only as directed. ■ only use the dose cup provided. ■ do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 30 mL every 4 hrs children 6 to under 12 yrs 15 mL every 4 hrs children 4 to under 6 yrs ask a doctor children under 4 yrs do not use		<b>Directions</b> Take only as directed. ■ only use the dose cup provided. ■ do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 15 mL every 4 hrs children 6 to under 12 yrs 15 mL every 4 hrs children 4 to under 6 yrs ask a doctor children under 4 yrs do not use	
<b>Other information</b> each 30 mL contains: sodium 56 mg ■ store at no greater than 25°C and do not refrigerate		<b>Other information</b> each 15 mL contains: sodium 56 mg ■ store at no greater than 25°C and do not refrigerate	
<b>Inactive ingredients</b> citric acid monohydrate, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum		<b>Inactive ingredients</b> citric acid monohydrate, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum	
<b>Questions or comments?</b> Call 1-800-910-6874		<b>Questions or comments?</b> Call 1-800-910-6874	

**TAMPER EVIDENT: DO NOT USE IF PRINTED SHIRKIBAND SEAL AROUND THE RECIPIER IS BROKEN OR MISSING.**

This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademark Vicks® DayQuil® Severe Cold & Flu Berry Liquid.

7706 0524

124 140070 800 NDC 11673-824-02

124 140070 800 NDC 11673-824-02

Print/Varnish Omit Area

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu Berry Liquid\*

### Nighttime Severe Cold & Flu

**Acetaminophen**  
(Pain Reliever / Fever Reducer)  
**Dextromethorphan HBr**  
(Cough Suppressant)  
**Doxylamine Succinate**  
(Antihistamine)  
**Phenylephrine HCl**  
(Nasal Decongestant)

- Headache, fever, sore throat, minor aches and pains
- Cough
- Sneezing, runny nose
- Nasal congestion, sinus pressure
- Nighttime relief

Alcohol free



**12 FL OZ (354 mL)**

Berry Flavor

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING  
PEEL BACK FOR DRUG FACTS

**Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- runny nose & sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen

094.14.0070 R00 C-002262-01-088-0000  
Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in India  
TM & ©2024 Target Brands, Inc.  
NDC 11673-828-01

**PARENTS:** Learn about medication safety at [www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

774L 0624

**NOT FOR RESALE**

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_ Print/Varnish Omit Area

**Drug Facts (continued)**

- 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
  - skin redness
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**Directions**

- take only as directed
- only use the dose cap provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 4 hrs
- children 6 to under 12 yrs 15 mL every 4 hrs
- children 4 to under 6 yrs ask a doctor
- children under 4 yrs do not use

**Other information**

- each 30 mL contains: sodium 98 mg
- do not exceed 4 doses per 24 hrs
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** citric acid monohydrate, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

**Questions or comments?** Call 1-800-910-6874

\*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® DayQuil® Severe Cold & Flu Berry Liquid.

Compare to the active ingredients in Vicks® DayQuil® Severe Cold & Flu Liquid\*

### Daytime Relief Severe Cold & Flu

**Acetaminophen**  
(Pain Reliever / Fever Reducer)  
**Dextromethorphan HBr**  
(Cough Suppressant)  
**Guaifenesin**  
(Expectorant)  
**Phenylephrine HCl**  
(Nasal Decongestant)

- Headache, fever, sore throat, minor aches and pains
- Cough
- Chest congestion, thins and loosens mucus
- Nasal congestion, sinus pressure

Alcohol free



**12 FL OZ (354 mL)**

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING  
PEEL BACK FOR DRUG FACTS

**Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

094.14.0070 R00 C-002262-01-088-0000  
Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in India  
TM & ©2024 Target Brands, Inc.  
NDC 11673-830-01

**PARENTS:** Learn about medication safety at [www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

774L 0624

**NOT FOR RESALE**

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_ Print/Varnish Omit Area

**Drug Facts (continued)**

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**When using this product, do not use more than directed.**

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**Directions**

- take only as directed
- only use the dose cap provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 4 hrs
- children 6 to under 12 yrs 15 mL every 4 hrs
- children 4 to under 6 yrs ask a doctor
- children under 4 yrs do not use

**Other information**

- each 15 mL contains: sodium 56 mg
- do not exceed 4 doses per 24 hrs
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** citric acid monohydrate, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

**Questions or comments?** Call 1-800-910-6874

\*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® DayQuil® Severe Cold & Flu Liquid.

# DAYTIME SEVERE AND NIGHTTIME SEVERE COLD AND FLU

daytime severe - acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl, nighttime severe-acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-824

Packaging			
#	Item Code	Package Description	Marketing Start Date
1	NDC:11673-824-02	1 in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2024

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity

<b>Part 1</b>	1 BOTTLE	355 mL
<b>Part 2</b>	1 BOTTLE	355 mL

## Part 1 of 2

### NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl liquid

#### Product Information

<b>Item Code (Source)</b>	NDC:11673-828
<b>Route of Administration</b>	ORAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

#### Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	

#### Product Characteristics

<b>Color</b>	red (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	



**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

**Part 2 of 2****DAYTIME SEVERE COLD AND FLU**

acetaminophen,dextromethorphan hbr, guaifenesin, phenylephrine hcl liquid

**Product Information**

Item Code (Source)	NDC:11673-830
Route of Administration	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SUCROSE</b> (UNII: C151H8M554)	

<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	

### Product Characteristics

<b>Color</b>	orange (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/01/2024	

**Labeler** - TARGET CORPORATION (006961700)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(11673-824)