

ALKA-SELTZER PLUS SEVERE COUGH MUCUS AND CONGESTION LIQUID GELS-
alka-seltzer plus severe cough mucus and congestion capsule, liquid filled
Bayer HealthCare LLC.

Alka-seltzer Plus Severe Cough Mucus & Congestion LG

Drug Facts

Active ingredients (in each capsule)

Active ingredients (in each capsule)

Acetaminophen 250 mg
Dextromethorphan hydrobromide 10 mg
Guaifenesin 200 mg
Phenylephrine hydrochloride 5 mg

Purposes

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms due to a cold or flu:
 - nasal congestion · sinus congestion and pressure
 - minor aches and pains · headache
 - cough · sore throat
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough with excessive phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts.
- These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 12 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use


Other information

- store at 15-25°C (59-77°F)

Inactive ingredients

FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol-sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-03 69 (Mon-Fri 9AM - 5PM EST)



SEVERE

Alka-Seltzer PLUS

Cough, Mucus & Congestion

ACETAMINOPHEN / Pain Reliever-Fever Reducer

Dextromethorphan HBr / Cough Suppressant
Guaifenesin / Expectorant
Phenylephrine HCl / Nasal Decongestant

- ▶ Nasal & Chest Congestion
- ▶ Headache
- ▶ Sore Throat
- ▶ Mucus
- ▶ Fever



20 LIQUID GELS (Liquid Filled Capsules)

3476

3476

1/2



LOT & EXP CODE AREA

LOT/EXP

Bayer

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Do not use if blister is cut or broken.

DOES NOT CONTAIN ASPIRIN

Questions or comments? 1-800-986-0369
(Mon-Fri 9AM - 5PM EST)

Inactive Ingredients FD&C red #40, gelatin, glycerin, polyethylene glycol, polydioxane, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol-sorbitan solution, titanium dioxide

Other Information
● store at 15-25°C (59-77°F)
● do not take more than the recommended dose

Directions
● adults and children 12 years and over, take 2 capsules with water every 4 hours. Do not exceed 12 capsules in 24 hours
● children under 12 years: do not use or as directed by a doctor.

Drug Facts (continued)
● do not take more than the recommended dose
● adults and children 12 years and over, take 2 capsules with water every 4 hours. Do not exceed 12 capsules in 24 hours
● children under 12 years: do not use or as directed by a doctor.

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
When using this product do not exceed recommended dosage
Stop use and ask a doctor if
● pain, cough, or nasal congestion gets worse or lasts more than 7 days
● fever gets worse or lasts more than 3 days
● redness or swelling is present
● new symptoms occur
● cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
● nervousness, dizziness, or sleeplessness occurs
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Made in Spain
© 2017 Bayer.
Dist. by: Bayer Healthcare LLC
Whippany, NJ 07981
Bayer, the Bayer Cross and Alka-Seltzer Plus are registered trademarks of Bayer.

ALKA-SELTZER PLUS SEVERE COUGH MUCUS AND CONGESTION LIQUID GELS

alka-seltzer plus severe cough mucus and congestion capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-0505
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SORBITOL (UNII: 506T60A25R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SHELLAC (UNII: 46N107B71O)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE (UNII: FZ989GH94E)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
WATER (UNII: 059QF0KO0R)	
GELATIN (UNII: 2G86QN327L)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	AS;M
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0505-20	2 in 1 CARTON	01/07/2015	
1		20 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/07/2015	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2023

Bayer HealthCare LLC.