

**LINCOMIX 20- lincomycin hydrochloride granule**  
**Zoetis Inc.**

-----  
**Lincomix 20**

**DESCRIPTION**

**LINCOMIX® 20**

(lincomycin)  
Type A Medicated Article  
20 grams/lb

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers.

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*.

For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*.

**Each pound contains:**

**Active Drug Ingredient**

Lincomycin .....20 grams  
(as lincomycin hydrochloride agricultural grade)

**Inactive ingredients**

Soybean hulls, #20 grind; light liquid paraffin, Ph.Eur.

**IMPORTANT:**

**Must be thoroughly mixed in feeds before use.**

**Store open bag in dry place to prevent caking.**

**Store at room temperature.**

Approved by FDA under NADA # 097-505

**Caution**

**Caution: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.**

**Broilers**

**BROILERS**

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to lincomycin in broilers.

**DIRECTIONS FOR USE**

**For the control of necrotic enteritis:**

LINCOMIX 20, 20 grams/lb, should be mixed into the complete feed supplied to broiler chickens so that the final feed contains 2 grams of lincomycin per ton of feed.

**MIXING DIRECTIONS**

**Type C Medicated Feeds**

---

<b>Intermediate Premix</b> Amount of LINCOMIX 20 per 1000 lb (454 kg) of Feed Ingredients	<b>Complete Feed</b> Amount of Intermediate Premix to Provide 2 Grams of lincomycin per ton of Type C Medicated Feed <i>lincomycin per ton of feed</i> <i>2 grams</i>
50 lbs	2 lb
10 lbs	10 lbs
5 lbs	20 lb

---

**WARNING:** When using LINCOMIX 20 in approved combinations with other drugs, follow the required withdrawal times for those drugs. No drug withdrawal period is required before slaughter of birds fed LINCOMIX 20 at the approved concentration (2 grams lincomycin per ton of feed).

**CAUTION:** Not for use in layers, breeders, or turkeys. (For additional **Cautions**, see below.)

**Swine**

**SWINE**

For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*. For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*

**DIRECTIONS FOR USE**

**For the treatment of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:**

Feed 100 grams of lincomycin per ton of complete feed as the sole ration for three weeks, or until signs of disease (watery, mucoid or bloody stools) disappear.

**For the treatment and control of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:**

Feed 100 grams of lincomycin per ton of complete feed as the sole rations for three weeks, or until signs of disease (watery, mucoid or bloody stools) disappear, followed by 40 grams of lincomycin per ton.

**For the control of swine dysentery and porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:** Feed 40 grams of lincomycin per ton of complete feed as the sole ration. For use in animals or on premises with a history of swine dysentery, but where symptoms have not yet occurred.

**For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*:** Feed 100-200 grams of lincomycin per ton of complete feed as the sole ration for 21

days.

## MIXING DIRECTIONS

### Type C Medicated Feeds

**For treatment of swine dysentery, and the control of porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:**

To make complete feed containing 100 grams of lincomycin, add 5 lbs of LINCOMIX 20 per ton.

**For control of swine dysentery and porcine proliferative enteropathies (ileitis) caused by *Lawsonia intracellularis*:**

To make complete feed containing 40 grams of lincomycin, add 2 lbs of LINCOMIX 20 per ton.

**For reduction in the severity of the effects of respiratory disease associated with *Mycoplasma hyopneumoniae*:**

To make complete feed containing 100-200 grams of lincomycin, add 5-10 lbs of LINCOMIX 20 per ton.

Additional mixing directions to make complete feed containing 40, 100 or 200 grams of lincomycin per ton are provided below.

## MIXING DIRECTIONS

<b>Intermediate Premix</b> Amount of LINCOMIX 20 per 1000 lb (454 kg) of Feed Ingredients	<b>Complete Feed</b> Amount of Intermediate Premix to Provide Desired Grams of lincomycin per ton of Type C Medicated Feed
	<b>lincomycin per ton of feed</b>
125 lbs	<b>40 grams    100 grams    200 grams</b>
100 lbs	16 lbs      40 lbs      80 lbs
50 lbs	20 lbs      50 lbs      100 lbs
	40 lbs      100 lbs      200 lbs

**WARNING:** When using LINCOMIX 20 in approved combinations with other drugs, follow the required withdrawal times for those drugs. No drug withdrawal period is required before slaughter of swine fed LINCOMIX 20 at approved concentrations (40, 100 or 100-200 grams lincomycin per ton of feed).

### NOT FOR HUMAN USE

**CAUTION:** Occasionally, swine fed lincomycin may within the first two days after the onset of treatment develop diarrhea and/or swelling of the anus. On rare occasions, some pigs may show reddening of the skin and irritable behavior. These conditions have been self-correcting within five to eight days without discontinuing the lincomycin treatment. The effects of lincomycin on swine reproductive performance, pregnancy, and lactation have not been determined.

Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

Good Manufacturing Practices Should Be Observed in Preparing Feeds Containing LINCOMIX 20.

This Includes Appropriate Clean-Out Procedures to Avoid Cross-Contamination.

**Restricted Drug (California) – Use Only as Directed**



<b>Product Type</b>	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:54771-1439	
<b>Route of Administration</b>	ORAL			
<b>Active Ingredient/Active Moiety</b>				
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
	LINCOMYCIN HYDROCHLORIDE (UNII: M6T05Z2B68) (LINCOMYCIN - UNII:BOD072YW0F)	LINCOMYCIN	20 g in 0.453 kg	
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:54771-1439-1	22.7 kg in 1 BAG		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
NADA	NADA097505	06/30/1976		

**Labeler** - Zoetis Inc. (828851555)

Revised: 11/2019

Zoetis Inc.