

FOSTER AND THRIVE ACID REDUCER- famotidine tablet, film coated
Strategic Sourcing Services LLC

McKesson Acid Reducer Drug Facts

Active ingredient (in each tablet)

Famotidine 20 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

Questions or comments?

Call **833-358-6431** Monday to Friday 9:00am to 7:00pm EST

Principal Display Panel

COMPARE TO MAXIMUM STRENGTH PEPCID® AC ACTIVE INGREDIENT

Foster & Thrive™

MAXIMUM STRENGTH

Acid Reducer

Famotidine Tablets 20 mg

ACID REDUCER

Gluten Free

Just One Tablet Prevents & Relieves Heartburn Due to Acid Indigestion

ACTUAL SIZE

SEE NEW WARNINGS

50 TABLETS

ACTUAL SIZE



©2023 McNeil Corporation
Distributed by: McNeil Corp.
via Strategic Sourcing Services LLC,
Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com
REF: 05/23

NDC 70677-1101-2

COMPARE TO MAXIMUM STRENGTH
PEPCID® AC ACTIVE INGREDIENT*

Foster & Thrive™ MAXIMUM STRENGTH Acid Reducer Famotidine Tablets 20 mg ACID REDUCER

Gluten Free

Just One Tablet Prevents & Relieves
Heartburn Due to Acid Indigestion



ACTUAL SIZE

SEE NEW WARNINGS

50 TABLETS

19471 KM C1

0 10939 95643 9

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Tips to Reduce Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bed time
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking


Drug Facts	Purpose
Active ingredient (in each tablet) Famotidine 20 mg..... Acid reducer	
Uses <ul style="list-style-type: none"> ■ relieve heartburn associated with acid indigestion and sour stomach ■ prevent heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages 	
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use <ul style="list-style-type: none"> ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. There may be signs of a serious condition. See your doctor. ■ with other acid reducers Ask a doctor before use if you have <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be sign of a more serious condition. 	

Drug Facts (continued)
<ul style="list-style-type: none"> ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck, or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs. Stop use and ask a doctor if <ul style="list-style-type: none"> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Drug Facts (continued)
Directions Adults and children 12 years and over: <ul style="list-style-type: none"> ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn

Drug Facts (continued)
<ul style="list-style-type: none"> ■ store at 20°-25 °C (68°-77 °F) ■ protect from moisture Inactive ingredients canuba wax, colloidal silicon dioxide, croscarmellose sodium, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, talc, titanium dioxide

Foster & Thrive™
MAXIMUM STRENGTH
Acid Reducer
Famotidine Tablets 20 mg
ACID REDUCER



FOSTER AND THRIVE ACID REDUCER

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1101
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	L194
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1101-2	1 in 1 CARTON	08/31/2023	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:70677-1101-1	25 in 1 CARTON	11/01/2023	

2

1 in 1 BLISTER PACK; Type 0: Not a Combination Product

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077351	08/31/2023	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 11/2023

Strategic Sourcing Services LLC