

**GOOD SENSE NIGHT TIME- acetaminophen, dextromethorphan hbr,
doxylamine succinate solution
L. Perrigo Company**

Perrigo Night Time Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

| | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 6 hrs |
| children 4 to under 12 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- **each 30 mL contains:** sodium 38 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions?

1-800-719-9260

Package/Label Principal Display Panel

GOODSENSE®

Multi-Symptom Relief

Pain Reliever, Fever Reducer

Cough Suppressant

Antihistamine

Night Time

Cold & Flu

Acetaminophen

Dextromethorphan HBr

Doxylamine Succinate

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough

Original Flavor

Compare to active ingredients of Vicks[®] NyQuil[®] Cold & Flu

12 FL OZ (354 mL)

ALCOHOL 10%

NDC 0113-0335-40

GOODSENSE®

Multi-Symptom Relief

Pain Reliever, Fever Reducer
Cough Suppressant
Antihistamine

Night Time Cold & Flu

Acetaminophen
Dextromethorphan HBr
Doxylamine Succinate

- Headache, Fever, Sore Throat, Minor Aches & Pains
- Sneezing, Runny Nose
- Cough

Original Flavor

Compare to active ingredients of
Vicks® NyQuil® Cold & Flu

12 FL OZ (354 mL) ALCOHOL 10%

: 33540 C2 F5

Distributed By

Perrigo®

Allergan, MI 49010

Empty & Replace Cap



PLASTIC BOTTLE



PLASTIC CUP

how2recycle.org

Gluten Free

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED
NECKBAND IS BROKEN
OR MISSING

N
0113-0335-40
7



Drug Facts

| Active ingredients (in each 30 mL) | Purpose |
|------------------------------------|-----------------------------|
| Acetaminophen 650 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 30 mg | Cough suppressant |
| Doxylamine succinate 12.5 mg | Antihistamine |

Uses temporarily relieves common cold/flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

PEEL BACK AT
CORNER FOR MORE
INFORMATION

: 33540 C2 B1

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ liver disease ■ glaucoma
■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or

ADHESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)

■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.


If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed – see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 30 mL every 6 hrs
12 yrs & over

ADHESIVE AREA
NO VARNISH • NO TYPE



GOOD SENSE NIGHT TIME

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0113-0335 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|---------------------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 650 mg in 30 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 30 mg in 30 mL |
| DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 12.5 mg in 30 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ALCOHOL (UNII: 3K9958V90M) | |
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S) | |
| FD&C YELLOW NO. 6 (UNII: H77VEI93A8) | |
| HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| WATER (UNII: 059QF0KO0R) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) | |

Product Characteristics

| | | | |
|-----------------|---------------------------------------|---------------------|--|
| Color | GREEN (clear, bright green) | Score | |
| Shape | | Size | |
| Flavor | FRUIT (anise / cooling menthol aroma) | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0113-0335-30 | 177 mL in 1 BOTTLE; Type 0: Not a Combination Product | 08/19/2011 | 03/17/2014 |
| 2 | NDC:0113-0335-38 | 296 mL in 1 BOTTLE; Type 0: Not a Combination Product | 09/20/2011 | 01/17/2014 |
| 3 | NDC:0113-0335-34 | 237 mL in 1 BOTTLE; Type 0: Not a Combination Product | 07/20/2012 | |
| 4 | NDC:0113-0335-40 | 354 mL in 1 BOTTLE; Type 0: Not a Combination Product | 08/20/2012 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 08/19/2011 | |

Labeler - L. Perrigo Company (006013346)

Revised: 10/2024

L. Perrigo Company