

**ALKA-SELTZER- buffered aspirin tablet, effervescent
Lil' Drug Store Products, Inc.**

Alka-Seltzer ®

Pouch Header/PDP

Do not use if pouch is torn or open.

Alka-Seltzer ®

Buffered aspirin (NSAID)/Pain reliever/fever reducer

ORIGINAL FLAVOR

[Bayer logo]

Fast Relief

2 EFFERVESCENT TABLETS

Active ingredient

Active ingredient (in each tablet)

Buffered aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Purpose

Pain reliever/fever reducer

Uses

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscle pain
- backache
- toothache
- menstrual pain
- colds
- minor pain of arthritis
- temporarily reduces fever

Warnings

Warnings

Reye's syndrome

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Stomach bleeding warning: This product contains an NSAID, which

Do not use

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- diabetes
- gout
- arthritis

Stop use and ask a doctor

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Directions Fully dissolve 2 tablets in 4 ounces of water before taking. Adults and children 12 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 8 tablets in 24 hours. Adults 60 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 4 tablets in 24 hours. Children under 12 years: consult a doctor.

Other information

Other information

- **each tablet contains:** sodium 568 mg
- store at room temperature. Avoid excessive heat.
- Alka-Seltzer Original Flavor in water contains principally the analgesic sodium acetylsalicylate

Inactive ingredients

Inactive ingredients anhydrous citric acid, sodium bicarbonate

Questions?

Questions? 1-800-986-0369 Dist. By Bayer Healthcare LLC, Whippany, NJ 07981

Alka-Seltzer® Gravity Sleeve (unit dose) - PDP/Label

[Bayer logo]

Alka-

Seltzer®

Buffered aspirin (NSAID)/

Pain reliever/fever reducer

ORIGINAL FLAVOR

15 PACKETS OF

2 EFFERVESCENT TABLETS

[UPC]

Do not use • if you are allergic to aspirin or any other pain reliever/fever reducer • if you have ever had an allergic reaction to this product or any of its ingredients **Ask a doctor before use** if • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet **Ask a doctor or pharmacist before use** if you are taking a prescription drug for • diabetes • gout • arthritis **Stop use and ask a doctor** if • an allergic reaction occurs (seek medical help right away) • you experience any of the following signs of stomach bleeding (feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better) • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • ringing in the ears or a loss of hearing occurs **If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** Fully dissolve 2 tablets in 4 ounces of water before taking. Adults and children 12 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 8 tablets in 24 hours. Adults 60 years and over: take 2 tablets every 4 hours, or as directed by a doctor. Do not exceed 4 tablets in 24 hours. Children under 12 years: consult a doctor **Other information** • each tablet contains: sodium 568 mg • Store at room temperature. Avoid excessive heat. • Alka-Seltzer Original Flavor in water contains principally the analgesic sodium acetylsalicylate. **Inactive ingredients** anhydrous citric acid, sodium bicarbonate **Questions? 1-800-986-0369**
Dist. By: Bayer Healthcare LLC, Whippany, NJ 07981
88052145 **MADE IN MEXICO**

Do not use if pouch is torn or open.

Alka-Seltzer®

Buffered aspirin (NSAID)/Pain reliever/fever reducer
ORIGINAL FLAVOR



Fast Relief 2 EFFERVESCENT TABLETS

Active ingredient (in each tablet)

Buffered aspirin 325 mg (NSAID)*.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Purpose

Uses • temporarily relieves minor aches and pains due to:
• headache • muscle pain • backache • toothache • menstrual pain • colds • minor pain of arthritis • temporarily reduces fever

Warnings **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

88052161

X25FCP EXP: DEC/2025 6



ALKA-SELTZER

buffered aspirin tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5206
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg	
Inactive Ingredients				
Ingredient Name			Strength	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	25mm	
Flavor	CITRUS	Imprint Code	ALKA;SELTZER	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5206-0	15 in 1 BOX, UNIT-DOSE	06/01/2023	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	06/01/2023		

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2023

Lil' Drug Store Products, Inc.