

IBUPROFEN- ibuprofen capsule, liquid filled BJWC

Drug Facts

Active ingredient (in each capsule)

Solubilized ibuprofen equal to

200 mg ibuprofen (NSAID) †
(present as the free acid and potassium salt)

†nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause delivery problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: **potassium 20 mg**
- read all warnings and directions before use
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

Inactive ingredients

FD&C Blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan and sorbitol

Questions or comments?

call toll free **1-800-934-1204**

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH “SEALED for YOUR PROTECTION” IS BROKEN OR MISSING.

*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Advil[®] LIQUI-GELS[®].

Manufactured for:
Bionpharma Inc.

600 Alexander Road,
Princeton, NJ 08540

Distributed by:
BJ's Wholesale Club
25 Research Drive
Westborough, MA 01581
Questions? 1-800-934-1204

R0422

BJ # 228163

L0000603

300's Bottle Label

Compare to the active ingredient in Advil® LIQUI-GELS®*

berkley jensen®

IBUPROFEN CAPSULES, 200mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

Actual Size

300 SOFTGELS**

**LIQUID FILLED CAPSULES

Compare to the active ingredient in Advil® LIQUI-GELS®

berkley jensen

IBUPROFEN
CAPSULES, 200mg

PAIN RELIEVER/FEVER REDUCER (NSAID)

Actual Size
300
SOFTGELS**

**LIQUID FILLED CAPSULES

DO NOT USE IF TAMPER-EVIDENT SEAL UNDER BOTTLE CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

Drug Facts

Active ingredient (in each capsule)
Sodium ibuprofen (equivalent to 200 mg ibuprofen (NSAID))... Pain reliever/fever reducer (for relief as the free acid and potassium salt).
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
headache toothache backache menstrual cramps the common cold muscular aches minor pain of arthritis temporarily reduces fever


Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives facial swelling asthma/wheezing shock skin redness rash blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are aged 60 or older have had stomach ulcers or bleeding problems take alcohol (drinking, participating) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) take 3 or more alcoholic drinks every day while using this product take more or for longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risks are higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery

Manufactured for:
Berkley Jensen, Inc.
600 Alexander Road
Princeton, NJ 08540
Distributed by:
BJ's Wholesale Club
25 Research Drive
Westborough, MA 01581
Questions? 1-800-934-1204
R0422

BJ # 228163 L0000603
8867010814915
Exp. Date:
Lot No.:

FEEL HERE



IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-212
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	blue	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	IB1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:68391-212-83	300 in 1 BOTTLE; Type 0: Not a Combination Product	08/14/2019	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	08/14/2019	

Labeler - BJWC (159082692)

Registrant - Bionpharma Inc. (079637826)

Establishment			
Name	Address	ID/FEI	Business Operations
Patheon Softgels Inc.		002193829	manufacture(68391-212)

Revised: 12/2022

BJWC