

**ACID REDUCER - omeprazole tablet, delayed release**  
**Winco Foods, LLC.**

-----

**Tips for Managing Heartburn**

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

***Drug Facts***

***Active ingredient (in each tablet)***

Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium)

***Purpose***

Acid reducer

***Use***

- treats frequent heartburn (occurs ***2 or more*** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

***Warnings***

**Allergy alert:**

Do not use if you are allergic to omeprazole

**Do not use if you have:**

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have:**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

**Ask a doctor or pharmacist before use if you are taking:**

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- digoxin (heart medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

**Stop use and ask a doctor if:**

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

- swallow whole. Do not chew or crush tablets.

### **Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

### **Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77° F) and protect from moisture

### **Inactive ingredients**

crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide.

### **Questions?**

call **800-824-1706**

Monday - Friday 9am - 4pm MST

Distributed by:

**WINCO FOODS LLC**

BOISE, ID 83704

**MADE IN INDIA**

Code: TS/DRUGS/22/2009

**PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (14 Tablet Bottle)**

**WinCo®**

**FOODS**

**Omeprazole**

**Delayed-Release**

**OMEPRAZOLE 20 mg/**

**ACID REDUCER**

**14 TABLETS**

# Top Ply

 <b>Omeprazole Delayed-Release</b> OMEPRAZOLE 20 mg/ ACID REDUCER 14 TABLETS	<b>Tips for Managing Heartburn</b> <ul style="list-style-type: none"><li>• Do not lie flat or bend over after eating</li><li>• Do not wear tight-fitting clothing around the stomach</li><li>• Do not eat before bedtime</li><li>• Raise the head of your bed</li><li>• Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables</li><li>• Eat slowly and avoid big meals</li><li>• If overweight, lose weight</li><li>• Quit smoking</li></ul>	<b>TAMPER EVIDENT - Do not use if seal over bottle opening is broken or missing.</b>  LM-3193 WF1806 CODE: TS/DRUGS/22/2009	Lift Here
	<b>DISTRIBUTED BY:</b> WINCO FOODS, LLC, BOISE, ID 83704 <b>MADE IN INDIA</b>		

## Top Ply (Page #1)

<b>Active ingredient (in each tablet)</b> Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium)	<b>Purpose</b> Acid reducer
<b>Use</b> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	
<b>Warnings Allergy alert:</b> Do not use if you are allergic to omeprazole. Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor. Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain	
<b>Ask a doctor or pharmacist before use if you are taking:</b> ■ warfarin, clopidogrel or cimetidine (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus or mycophenolate mofetil (immune system medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methotrexate (arthritis medicine) <b>Stop use and ask a doctor if:</b> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days	

## Back of Top Ply (Page #2)

## Base (Page #3)

■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain ■ pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours **14-Day Course of Treatment** ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. **Repeated 14-Day Courses (if needed)** ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition. **Other information** ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20° to 25°C (68° to 77°F) and protect from moisture **Inactive ingredients** crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hydroxypropyl methylcellulose, magnesium stearate, methacrylic acid copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polyorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearoyl fumarate, sugar spheres (which contains liquid glucose, starch (maize) and sucrose), talc, titanium dioxide, triethyl citrate and yellow iron oxide. **Questions? Call 800-624-1706 Monday - Friday 9am - 4pm MST**

## PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg Container Carton Label

Compare to the active ingredient  
in PRILOSEC OTC®

WinCo®

FOODS

Omeprazole

Delayed-Release

Omeprazole 20 mg/

ACID REDUCER

Treats Frequent Heartburn

Occurring 2 Or More Days A Week

14 TABLETS

ONE 14-DAY COURSE

OF TREATMENT



## ACID REDUCER

omeprazole tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67091-316
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OMEPRAZOLE MAGNESIUM</b> (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CROSPROVIDONE (35 .MU.M)</b> (UNII: 40UAA97IT9)	
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)	
<b>HYDROXYPROPYL CELLULOSE (90000 WAMW)</b> (UNII: UKE75GEA7F)	
<b>HYPROMELLOSE 2910 (5 MPA.S)</b> (UNII: R75537T0T4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A</b> (UNII: NX76LV5T8J)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 4000</b> (UNII: 4R4HFI6D95)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>SODIUM STEARYL FUMARATE</b> (UNII: 7CV7WJK4UI)	
<b>DEXTROSE, UNSPECIFIED FORM</b> (UNII: IY9XDZ35W2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>HYDROXYPROPYL CELLULOSE (45000 WAMW)</b> (UNII: 8VAB711C5E)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	

### Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	RECTANGLE (Oblong)	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	Z;69
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67091-316-14	1 in 1 CARTON	06/06/2018	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206877	06/06/2018	

**Labeler** - Winco Foods, LLC. (056098817)

**Registrant** - Aurohealth LLC (078728447)

**Establishment**

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(67091-316) , MANUFACTURE(67091-316)

Revised: 6/2024

Winco Foods, LLC.