

**ASPIRIN NSAID- aspirin tablet, film coated**  
**Valu Merchandisers Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Best Choice 44-157**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

temporarily relieves:

- headache
- pain and fever of colds
- toothache
- menstrual pain
- muscle pain
- minor pain of arthritis

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### ***Directions***

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments?**

**1-800-426-9391**

***Principal display panel***

**Best Choice®**

COMPARE TO ACTIVE INGREDIENT IN GENUINE BAYER® ASPIRIN†

**Aspirin (NSAID)**

Aspirin (NSAID)

Pain Reliever & Fever Reducer

Caffeine and Sodium Free

**100 TABLETS**

325 mg Each

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS  
BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Genuine Bayer® Aspirin.

50844 REV0318B15712

**PROUDLY DISTRIBUTED BY:  
VALU MERCHANDISERS, CO.  
5000 KANSAS AVE  
KANSAS CITY, KS 66106**

Best Choice®

100% Guaranteed

[www.bestchoicebrand.com](http://www.bestchoicebrand.com)

No print/No varnish  
Lot & Exp date

0 70038 1 20154  
BEST CHOICE  
9

COMPARE TO ACTIVE INGREDIENT IN GENUINE BAYER® ASPIRIN<sup>1</sup>

# Aspirin (NSAID)

Aspirin (NSAID)  
Pain Reliever & Fever Reducer

Caffeine and Sodium Free

Best  
Choice®

Best  
Choice®

Actual Size

**100 TABLETS**  
325 mg Each

B-2115-157-12  
REV0318815712

**Drug Facts (continued)**

it may cause problems in the unborn child or complications during delivery. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients** corn starch, hypromellose, polyethylene glycol, propylene glycol

**Questions or comments? 1-800-426-9391**

<sup>1</sup>This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Genuine Bayer® Aspirin.  
50844 REV0318815712

**Best Choice®**  
100% Guaranteed  
[www.bestchoicebrand.com](http://www.bestchoicebrand.com)

**PROUDLY DISTRIBUTED BY:**  
VALU MERCHANTISERS, CO.  
5000 KANSAS AVE  
KANSAS CITY, KS 66106

**Drug Facts (continued)**

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because**

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID)<sup>\*</sup> ..... Pain reliever/fever reducer

<sup>\*</sup>nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves:

- headache
- pain and fever of colds
- toothache
- menstrual pain
- muscle pain
- minor pain of arthritis

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior, with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Best Choice 44-157

## ASPIRIN NSAID

aspirin tablet, film coated

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:63941-157

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg	
<b>Inactive Ingredients</b>				
Ingredient Name			Strength	
STARCH, CORN (UNII: O8232NY3SJ)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
<b>Product Characteristics</b>				
Color	WHITE	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;157;ASPIRIN	
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-157-12	1 in 1 CARTON	02/10/1996	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:63941-157-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
3	NDC:63941-157-17	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/10/1996	
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH NOT FINAL	part343	02/10/1996		

**Labeler** - Valu Merchandisers Company (868703513)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(63941-157)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(63941-157)

### Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Revised: 8/2019

Valu Merchandisers Company