

**ASPIRIN- aspirin tablet**  
**ST. MARY'S MEDICAL PARK PHARMACY**

-----  
**Geri Care 44-249**

***Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever/fever reducer

***Uses***

temporarily relieves:

- pain and fever of colds
- headache
- muscle pain
- minor pain of arthritis
- toothache
- menstrual pain

***Warnings***

**Reye's syndrome:**Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

**Stomach bleeding warning:**This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - feel faint
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

### Other information

- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

### Inactive ingredient

corn starch

### Questions or comments?

1-800-540-3765

### Principal Display Panel

<p>NDC 60760-930-24  <b>ASPIRIN</b>                  (NSAID)                  PAIN RELIEVER/                  FEVER REDUCER  <b>TABLETS</b>  <b>325mg</b>                  QTY: 24                  LOT# XXXXXXXX                  EXP XX-XX                  RX# ???????                  MANUFACTURED BY:                  Geni-Care Pharmaceutical Corp.                  Brooklyn, NY 11204</p>	<p><b>ASPIRIN</b>                  (NSAID) TABLETS                  Pain Reliever/Fever Reducer  <b>325mg</b>                  QTY: 24                  RX# ???????                  NDC 60760-930-24                  LOT# XXXXXXXX                  EXP XX-XX</p>	<p><b>ASPIRIN</b>                  (NSAID) TABLETS                  Pain Reliever/Fever Reducer  <b>325mg</b>                  QTY: 24                  RX# ???????                  NDC 60760-930-24                  LOT# XXXXXXXX                  EXP XX-XX</p>	<p><b>ASPIRIN</b>                  (NSAID) TABLETS                  Pain Reliever/Fever Reducer  <b>325mg</b>                  QTY: 24                  RX# ???????                  NDC 60760-930-24                  LOT# XXXXXXXX                  EXP XX-XX</p>
<p><b>TAKE AS DIRECTED</b></p>			
<p>Rx only STORE AT CONTROLLED ROOM TEMPERATURE 15°- 30° C (59°-86°F)</p>			

<b>ASPIRIN</b>			
aspirin tablet			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:60760-930(NDC:57896-901)
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>Inactive Ingredients</b>			
	<b>Ingredient Name</b>		<b>Strength</b>
	STARCH, CORN (UNII: O8232NY3S)		

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;249;ASPIRIN
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:60760-930-24	24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/17/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/04/1993	

**Labeler** - ST. MARY'S MEDICAL PARK PHARMACY (063050751)

## Establishment

Name	Address	ID/FEI	Business Operations
ST. MARY'S MEDICAL PARK PHARMACY		063050751	relabel(60760-930) , repack(60760-930)

Revised: 12/2023

ST. MARY'S MEDICAL PARK PHARMACY