

**CONTAC COLD FLU DAY NIGHT- acetaminophen, phenylephrine hydrochloride, chlorpheniramine maleate**  
**Vespyr Brands, Inc.**

-----  
**CONTAC Cold + Flu Day Night**

***Drug Facts (Day)***

***Active ingredients(in each caplet) (Day)***

Acetaminophen 500 mg  
Phenylephrine hydrochloride 5 mg

***Purposes (Day)***

Pain reliever/fever reducer  
Nasal decongestant

***Uses (Day)***

Temporarily relieves common cold/flu symptoms:

- nasal congestion • headache
- minor aches and pains • stuffy nose
- sore throat • fever
- sinus congestion and pressure

***Warnings (Day)***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hrs, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, ask a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug

contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease • heart disease
- diabetes • high blood pressure
- thyroid • trouble urinating due to disease an enlarged prostate gland

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- you get nervous, dizzy, or sleepless
- new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions (Day)***

- **do not take more than directed (see overdose warning).**
- adults and children 12 yrs of age and over: take 2 caplets every 6 hrs as needed.
- do not take more than 8 caplets in 24 hrs.
- children under 12 yrs of age: ask a doctor.

***Other information (Day)***

store at 20°-25°C (68°-77°F)

***Inactive ingredients (Day)***

corn starch, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, potassium sorbate, povidone, sodium lauryl sulfate, stearic acid, talc.

***Questions or comments? (Day)***

call toll-free 1-855-874-0970 (English/Spanish) weekdays

***Drug Facts (Night)***

## **Active ingredients (in each caplet) (Night)**

Acetaminophen 500 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine hydrochloride 5 mg

## **Purposes (Night)**

Pain reliever/fever reducer  
Antihistamine  
Nasal decongestant

## **Uses (Night)**

Temporarily relieves common cold/flu symptoms:

- nasal congestion • sneezing • headache
- runny nose • fever • sore throat
- minor aches and pains

## **Warnings (Night)**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, occurs with or is followed by fever, headache, rash, nausea, or vomiting, ask a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **Ask a doctor before use if you have**

- liver disease • heart disease
- diabetes • high blood pressure
- thyroid disease • glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

## **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- do not exceed recommended dosage
- drowsiness may occur • avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness effect
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- you get nervous, dizzy, or sleepless • new symptoms occur

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions (Night)**

- **do not take more than directed (see overdose warning).**
- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- do not take more than 8 caplets in 24 hours.
- children under 12 years of age: ask a doctor.

**Other information (Night)**

store at 20°-25°C (68°-77°F)

**Inactive ingredients (Night)**

corn starch, D&C yellow No. 10 aluminum lake, FD&C yellow No. 6 aluminum lake, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, potassium sorbate, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide.

**Questions or comments? (Night)**

call toll-free 1-855-874-0970 (English/Spanish) M - F, 8am -5pm EST

NDC 71179-025-28

CONTAC ®

COLD + FLU

DAY

Acetaminophen -

Pain reliever/Fever reducer

Phenylephrine hydrochloride -

Nasal decongestant

**Head & Body Aches**

**Fever • Sore Throat**

**Nasal Congestion**

**Sinus Pressure**

16 CAPLETS

NIGHT

Acetaminophen -

Pain reliever/Fever reducer

Chlorpheniramine maleate -

Antihistamine

Phenylephrine hydrochloride -

Nasal decongestant

**Head & Body Aches**

**Fever • Sore Throat**

**Nasal Congestion**

**Sinus Pressure**

**Runny Nose • Sneezing**

12 CAPLETS

Maximum Strength

Multi-Symptom

DISTRIBUTED BY:

**Vespyr Brands<sup>®</sup>, Inc.**

East Windsor NJ 08520

©2022 Vespyr Brands<sup>®</sup>, Inc.

All rights reserved

CONTAC<sup>®</sup> is a registered trademark of Vespyr Brands<sup>®</sup>, Inc.

**www.contac.com**

# CONTAC® COLD+FLU

DAY 

NIGHT 

NIGHT

DAY 

**Acetaminophen - Pain reliever/Fever reducer**  
**Phenylephrine hydrochloride - Nasal decongestant**

**Acetaminophen - Pain reliever/Fever reducer**  
**Chlorpheniramine maleate - Antihistamine**  
**Phenylephrine hydrochloride - Nasal decongestant**

Head & Body Aches  
Fever • Sore Throat  
Nasal Congestion  
Sinus Pressure

Head & Body Aches  
Fever • Sore Throat  
Nasal Congestion  
Sinus Pressure  
Runny Nose • Sneezing

Maximum Strength  
Multi-Symptom



16 CAPLETS

12 CAPLETS

DAY 

NIGHT 

# CONTAC® COLD+FLU

Non-Drowsy Relief

Restful Relief



DISTRIBUTED BY:  
**Veepyr Brands\*, Inc.**  
East Windsor NJ 08520  
© 2022 Veepyr Brands\*, Inc.  
All rights reserved.  
CONTAC® is a registered trademark  
of Veepyr Brands\*, Inc.  
[www.contactac.com](http://www.contactac.com) UC402528-06-06

Made in Canada

**Drug Facts (continued)**

**When using this product:**

- do not exceed recommended dose
- do not take alcohol drinks
- do not take grapefruit juice
- avoid activities that require alertness or coordination when driving, operating machinery or using power tools
- avoid alcohol when driving, operating machinery or using power tools
- avoid activities that require alertness or coordination when driving, operating machinery or using power tools

**Drug Facts (continued)**

**Other information:**

- If you do not know if you are pregnant or breastfeeding, ask your doctor before using this product.
- If you are pregnant or breastfeeding, ask your doctor before using this product.
- If you are pregnant or breastfeeding, ask your doctor before using this product.

**Drug Facts (continued)**

**Directions:**

- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- do not take more than directed (see overdose warning).
- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- child under 12 years of age: ask a doctor.

**Other information:**

- Ask a doctor before use if you have any of the following conditions:
  - heart disease
  - high blood pressure
  - liver disease
  - kidney disease
  - asthma
  - diabetes
  - glaucoma
  - thyroid disease
  - prostate disease
  - stomach or intestinal problems
  - bleeding problems such as hemiparesis or chronic hemorrhoids
  - a bleeding problem such as a bleeding problem
  - a bleeding problem such as a bleeding problem
- Ask a doctor before use if you are taking any of the following:
  - aspirin
  - NSAIDs (such as ibuprofen, naproxen, etc.)
  - blood thinners (such as warfarin, clopidogrel, etc.)
  - other cold, flu, or allergy medicines
  - other medicines that contain acetaminophen
  - other medicines that contain phenylephrine
  - other medicines that contain chlorpheniramine

**Questions or comments?**

Call toll free 1-855-674-0970 (English/Spanish) M-F, 9am-5pm EST

**Read outer carton for complete directions and warnings.**

**Drug Facts (continued)**

**Directions:**

- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- do not take more than directed (see overdose warning).
- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- child under 12 years of age: ask a doctor.

**Other information:**

- Ask a doctor before use if you have any of the following conditions:
  - heart disease
  - high blood pressure
  - liver disease
  - kidney disease
  - asthma
  - diabetes
  - glaucoma
  - thyroid disease
  - prostate disease
  - stomach or intestinal problems
  - bleeding problems such as hemiparesis or chronic hemorrhoids
  - a bleeding problem
  - a bleeding problem
- Ask a doctor before use if you are taking any of the following:
  - aspirin
  - NSAIDs (such as ibuprofen, naproxen, etc.)
  - blood thinners (such as warfarin, clopidogrel, etc.)
  - other cold, flu, or allergy medicines
  - other medicines that contain acetaminophen
  - other medicines that contain phenylephrine
  - other medicines that contain chlorpheniramine

**Questions or comments?**

Call toll free 1-855-674-0970 (English/Spanish)

**Read outer carton for complete directions and warnings.**

**Drug Facts (continued)**

**Directions:**

- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- do not take more than directed (see overdose warning).
- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- child under 12 years of age: ask a doctor.

**Other information:**

- Ask a doctor before use if you have any of the following conditions:
  - heart disease
  - high blood pressure
  - liver disease
  - kidney disease
  - asthma
  - diabetes
  - glaucoma
  - thyroid disease
  - prostate disease
  - stomach or intestinal problems
  - bleeding problems such as hemiparesis or chronic hemorrhoids
  - a bleeding problem
  - a bleeding problem
- Ask a doctor before use if you are taking any of the following:
  - aspirin
  - NSAIDs (such as ibuprofen, naproxen, etc.)
  - blood thinners (such as warfarin, clopidogrel, etc.)
  - other cold, flu, or allergy medicines
  - other medicines that contain acetaminophen
  - other medicines that contain phenylephrine
  - other medicines that contain chlorpheniramine

**Questions or comments?**

Call toll free 1-855-674-0970 (English/Spanish)

**Read outer carton for complete directions and warnings.**

**Drug Facts (continued)**

**Directions:**

- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- do not take more than directed (see overdose warning).
- adults and children 12 years of age and over: take 2 caplets every 6 hours as needed.
- child under 12 years of age: ask a doctor.

**Other information:**

- Ask a doctor before use if you have any of the following conditions:
  - heart disease
  - high blood pressure
  - liver disease
  - kidney disease
  - asthma
  - diabetes
  - glaucoma
  - thyroid disease
  - prostate disease
  - stomach or intestinal problems
  - bleeding problems such as hemiparesis or chronic hemorrhoids
  - a bleeding problem
  - a bleeding problem
- Ask a doctor before use if you are taking any of the following:
  - aspirin
  - NSAIDs (such as ibuprofen, naproxen, etc.)
  - blood thinners (such as warfarin, clopidogrel, etc.)
  - other cold, flu, or allergy medicines
  - other medicines that contain acetaminophen
  - other medicines that contain phenylephrine
  - other medicines that contain chlorpheniramine

**Questions or comments?**

Call toll free 1-855-674-0970 (English/Spanish)

**Read outer carton for complete directions and warnings.**

Non-Drowsy Relief

# CONTAC® COLD+FLU

Restful Relief

PUMP-BUFFERED PAKAGING FEATURE: This product is packaged in a sealed blister pack. Do not use if blister or printed foil is broken.



## CONTAC COLD FLU DAY NIGHT

acetaminophen, phenylephrine hydrochloride, chlorpheniramine maleate kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71179-025
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71179-025-28	1 in 1 BOX; Type 0: Not a Combination Product	12/01/2023	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	0 BLISTER PACK	2
Part 2	0 BLISTER PACK	1

## Part 1 of 2

### CONTAC COLD FLU DAY

phenylephrine hydrochloride, acetaminophen tablet

### Product Information

<b>Item Code (Source)</b>	NDC:71179-024
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
TALC (UNII: 7SEV7J4R1U)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

## Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	sun
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2023	

## Part 2 of 2

### CONTAC COLD FLU NIGHT

acetaminophen, chlorpheniramine maleat, phenylephrine hydrochloride tablet

## Product Information

Item Code (Source)	NDC:71179-026
Route of Administration	ORAL

## Active Ingredient/Active Moiety



Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	moon
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2023	

