## REMIXDIL HAIR REGROWTH TREATMENT FOR WOMEN- minoxidil 2% solution Rida LLC

-----

Minoxidil 2% w/v

#### **USE**

Use to regrow hair on the scalp

Hair regrowth treatment

For external use only.

Flamable: Keep away from fire of flame.

#### Do not use if:

- Your degree of hair loss is different than shown on the side of this carton, because this product may not work for you
- You have no family history of hair loss
- Your hair loss is sudden and/or patchy
- · Your hair loss is associated with childbirth
- You do not know the reason for your hair loss
- You are under 18 years of age. Do not use on babies or children
- Your scalp is red, inflamed, infected, irritated, or painful
- You use other medicines on the scalp

Ask a doctor before use if you have heart disease

#### When using this product

- Do not apply on other parts of the body
- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water
- Some people have expereinced changes in hair color and/or texture
- It takes time to regrow hair. Results may occur at 2 months with twice a day usage.
   For some men, you may need to use this product for at least 4 months before you see results.
- The amount of hair regrowth is different for each person. This product will not work for everyone.

#### Stop use and ask a doctor if

- Chest pain, rapid heart beat, faintness, or dissiness occurs. Sudden, unexplained weight gain occurs
- Your hands or feet swell
- Scalp irritation or redness occurs
- You get unwanted facial hair growth
- You do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

- -

#### **DIRECTIONS**

- apply 1mL with dropper 2 times a day directly onto the scalp in the hair loss area
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

#### OTHER INFORMATION

- See hair loss pictures on the side of this carton
- Before use, read all information on carton and enclosed leaflet
- Keep the carton. It contains important information
- In clincial studies of mostly white women aged 18-45 years with moderate degrees of hair loss, the following responses to 2% minoxidil topical solution was reported: 19% of women reported moderate hair regrowth after using 2% minoxidil topical solution for 8 months (19% had moderate regrowth; 40% had minimal hair regrowth). This compares with the 7% of women reporting moderate hair regrowth after using a placebo, the liquid without minoxidil in it, for 8 months (7% had moderate regrowth; 33% had minimal regrowth).
- Store at controlled room temperature 20 °to 25 °C (68 °to 77 °F)

Alcohol, Propylene Glycol, Purified Water

Questions or comments? 1-800-641-8871 Visit website at www.rida-hair.com

#### Product label



2% MINOXIDIL TOPICAL SOLUTION, USP HAIR REGROWTH TREATMENT

#### **Drug Facts**

Active ingredient

Purpose

#### Warnings

The external use only.

Removable tasp away from the ar flame.

Benet use if your amount of hor loss is affected than that shown on the social flame is a carlon, because this product may not awar for you you have an drawn shown or have been as the social carlon, the social flame is suited an only operating your hard flow is an unique only operating the social flame of the social flame is the social flame of the social flame is the social flame in the social flame is the social flame in the social flame is the social flame is the social flame using this product of an object of the body - avoid contact with the eyes in case of accelerated contact, these eyes with large amounts of cooling water - some people have a perfected of changes is had color and/or texture - it takes that is regionable, it has many need to use this product of the social depends of the product of the social contact in the social contact is the contact of the social flame is the social flame in the social flame is the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame in the social flame in the social flame is the social flame i

- noy use and ask a douter? cheat poin, ropicineortheof, formess, or diziness occurs sucides, uneightined weight gain occurs your hands or feet aveil socip inflotion or redines occurs

- unwanted localited system occus 
- you do not see hid regarded in 4 months

May be harmful if used when pregrant or breash-feeding.

Seep out of restor of shiddern if it wouldweed, get medical help or contact 
a freision Control Center right away.

Directions - prime the sproyer by holding the bottle solight and pump the 
orange is limited to encours addison of hill sprov. Do not include misst -apply one 
mit with sprey (it apprays) zitimes a day descrip and the solight and pump the 
orange uses prime and experience of seeding the bottle solight and 
noncoosing to increase and experience or sprover, or has tools will begin again.

Other information - see had been pictures on side of this cando 
section use, sead of information common and employed back the best become noged the Systems with mist to mondated against affording the 
contact in the Systems with mist to mondated against affording the 
building response to 25 minused imposed solution and 
months (1% had incolorate agrower, 4% had not reproved in The Administration of 
months (1% had incolorate agrower, 4% had not reproved in the region of 
months (1% had incolorate agrower, 4% had regioned his region.) His 
companies with 7 solvement experting moderate had regioned in the 
surproved. This first include agreement is the 
surproved and the include agreement and in the surproved in the region of 
surproved and the include agreement and 
surproved and the previous and 
surproved and the 
surp

Inactive ingredients Abord, repplere object, nuried vic

Questions or comments? 1-800-641-8871

Remix dil

2% MINOXIDIL TOPICAL SOLUTION, USP

This Package Contains



Women's Remoidil Extra Strength (one-month supply) With child-resistant Sprayer pump applicator.

Information leaflet with complete directions on how to

9620 Towne Centre Dr., San Diego , CA 92121

abel part #





- Reactivates Hair
- Help Regrow Hair

**Topical Solution** 

Unscented

## HAIR REGROWTH TREATMENT

ONE MONTH SUPPLY

60 ml (2 fl.oz)

# Remix\_dil

## 2% MINOXIDIL TOPICAL SOLUTION, USP


HAIR REGROWTH TREATMENT 2% Minakidil Topical Solution is for women who have a general thinning of hoir on the top of the scalp as shown. 2% Minakidil Topical Solution has been shown to regrow hoir in women with the following degrees of thinning hoir or hair loss.











#### REMIXDIL HAIR REGROWTH TREATMENT FOR WOMEN

minoxidil 2% solution

<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83004-009
Route of Administration	TOPICAL		

# Active Ingredient/Active Moiety Ingredient Name Basis of Strength MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1) MINOXIDIL 2 g in 100 mL

Inactive Ingredients	
Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
NDC-03004 000			

1 01 01	4 in 1 BOX	11/22/2023			
1	60 mL in 1 BOTTLE; Type 0: Not a Combination Product				
Marketing Information					
Marketing	Information				
Marketing Marketing Category	Information  Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
Marketing	Application Number or Monograph	_	_		

### **Labeler -** Rida LLC (004425803)

Revised: 2/2025 Rida LLC