

EQUATE NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate solution
Wal-Mart Stores Inc

Wal-Mart Nighttime Cold & Flu Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 38 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow no. 10, FD&C green no. 3, FD&C yellow no. 6, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions?

1-888-287-1915

Package/Label Principal Display Panel

equate™

Compare to Vicks® NyQuil® Cold & Flu active ingredients

NIGHTTIME Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine

Nighttime Relief

- Acetaminophen – Aches, fever,
- Dextromethorphan HBr – Cough
- Doxylamine Succinate – Sneezing, runny nose

12 FL OZ (355mL)

10% ALCOHOL

NDC 49035-609-40

equate™

Compare to
Vicks® NyQuil®
Cold & Flu active
ingredients*

NIGHTTIME

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine

Nighttime Relief

- **Acetaminophen** - Aches, fever
- **Dextromethorphan HBr** - Cough
- **Doxylamine Succinate** - Sneezing, runny nose

12 FL OZ
(355mL)

10%
ALCOHOL

: 33540 2E FL

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED
NECKBAND IS BROKEN
OR MISSING 164152

6
8 1131 03677

1

Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 30 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine

Uses temporarily relieves common cold/flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

PEEL BACK AT
CORNER FOR MORE
INFORMATION

: 33540 2E B1

Drug Facts (continued)

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
■ liver disease ■ glaucoma
■ cough that occurs with too much phlegm (mucus) ■ a breathing problem such as emphysema or

ADHESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)

■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.


If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
------------------------------------	-------------------

ADHESIVE AREA
NO VARNISH • NO TYPE



EQUATE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-609
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR601S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

Product Characteristics

Color	GREEN (clear, bright green)	Score	
Shape		Size	
Flavor	FRUIT (anise / cooling menthol aroma)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-609-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/21/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/21/2019	

Labeler - Wal-Mart Stores Inc (051957769)

Revised: 10/2024

Wal-Mart Stores Inc