

**PRESSY TATTOO NUMBING CREAM- numbing cream cream**  
**Changsha Beiliang E-commerce Co., Ltd**

-----  
**83887-001**

**Active Ingredient**

Lidocaine 5%

**Purpose**

Local Anesthetic

**Use**

For temporary relief of local and anorectal itching, discomfort, and inflammation associated with anorectal disorders.

For temporary relief of pain, burning, and soreness.

**Warnings**

If symptoms worsen or do not improve within 7 days, seek medical advice.

Do not exceed the recommended daily dosage without doctor's instruction.

Consult a doctor immediately if bleeding occurs.

Do not insert this product into the rectum using fingers or any applicator.

Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.

Note: Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients

**Do not use**

Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.

**When Using**

If symptoms worsen or do not improve within 7 days, seek medical advice. Do not exceed the recommended daily dosage without doctor's instruction. Consult a doctor immediately if bleeding occurs.

Do not insert this product into the rectum using fingers or any applicator.

Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.

### **Stop Use**

Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients.

### **Ask Doctor**

Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients

### **Keep Out Of Reach Of Children**

This package is child-resistant. Keep out of reach of children

### **Directions**

Adults: When practical, cleanse the affected area with mild soap and warm water, and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.

### **Other information**

Keep away from direct sunlight or heat.

Store at room temperature between 15°C and 30°C (59°F-86°F).

This package is child-resistant. Keep out of reach of children. In case of accidental overdose or ingestion, call a doctor or poison control center immediately. Do not use this product if the seal is broken or missing.

### **Inactive ingredients**

Vitamin E Allantoin, Mineral Oil, Dimethicone, Methylparaben, Propylparaben, Petrolatum, IPBC, Diazolidinyl Urea, Glycerin, Cetostearyl Alcohol, Isopropyl Myristate, Propylene Glycol, Ammonium Methacrylate, Purified Water, Laurocapram, Steareth-2, Steareth-21, Rose Oil

### **PRINCIPAL DISPLAY PANEL**

### USAGE METHOD

- Clean and dry the area of skin where the tattoo will be applied.
- Apply a generous amount of cream onto the skin.
- Cover the cream-coated area with plastic wrap for approximately 40-60 minutes.
- Remove plastic wrap, gently pat to help cream absorb. Wipe off residue, wait 15-20 mins for maximum numbness before proceeding.

### Distributed by:

Opiyse care LLC  
1942 Broadway St Ste 314C  
Boulder, CO 80302  
Tel: 1-719-201-4680

### Manufactured by:

Changsha Beilang E-commerce Co., Ltd  
Changsha, Hunan, China 410000  
www.pressycare.com

SHELF LIFE: 3 YEARS



Pressy



6 HOURS  
MAXIMUM STRENGTH

EXTERNAL

NET: 30g/1.0oz

### Warnings (For external use only)

- If symptoms worsen or do not improve within 7 days, seek medical advice.
- Do not exceed the recommended daily dosage without doctor's instruction.
- Consult a doctor immediately if bleeding occurs.
- Do not insert this product into the rectum using fingers or any applicator.
- Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.
- Note:** Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling or pain due to possible allergies to product ingredients.

### Inactive Ingredients

Vitamin E, Allantoin, Mineral Oil, Dimethicone, Methylparaben, Propylparaben, Petrolatum, IPBC, Diazolidinyl Urea, Glycerin, Cetearyl Alcohol, Isopropyl Myristate, Propylene Glycol, Ammonium Methacrylate, Purified Water, Laureagaram, Steareth-2, Steareth-21, Rose Oil.

### DRUG FACTS

Active Ingredient	Purpose
Lidocaine 5%	Local Anesthetic

### Uses

- For temporary relief of local and anorectal itching, discomfort and inflammation associated with anorectal disorders.
- For temporary relief of pain, burning, and soreness.


### Directions

- Adults: When practical, cleanse the affected area with mild soap and warm water, and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product.
- Children under 12 years of age: Consult a doctor. Keep out of reach of children.
- Apply externally to the affected area up to 6 times daily.

### Other information

- Keep away from direct sunlight or heat.
- Store at room temperature between 15°C and 30°C (59°F - 86°F).
- This package is child-resistant. Keep out of reach of children.
- In case of accidental overdose or ingestion, call a doctor or poison control center immediately.
- Do not use this product if the seal is broken or missing.

MFG: 12/11/2023  
EXP: 12/10/2026  
LOT: 20231211-01



**DRUG FACTS**

**Active Ingredient**  
Lidocaine 5%

**Purpose**  
Local Anesthetic

**Uses**

- For temporary relief of local and procedural itching, discomfort, and inflammation associated with associated disorders.
- For temporary relief of pain, burning, and stinging.

**Warnings (For external use only)**

- If symptoms persist or do not improve within 7 days, seek medical advice.
- Do not exceed the recommended daily dosage without a doctor's instruction.
- Consult a doctor immediately if a burning occurs.
- Do not break the sealed unit the instant using fingers or any applicator.
- Some people may be allergic to the ingredients. Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain.
- Discontinue use and consult a doctor if symptoms persist or worsen, including redness, irritation, swelling, or pain due to possible allergies to product ingredients.

**DRUG FACTS(Continued)**

**Directions**

- Before first use, peel, clean the affected area with mild soap and warm water, and rinse thoroughly. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of the product.
- Use only under 12 years of age. Consult a doctor.
- Keep out of reach of children.
- Apply evenly to the affected area up to 3 times daily.

**Other Information**

- Keep away from heat, sunlight, or fire.
- Store at room temperature between 20°C and 25°C (68°F - 77°F).
- The package is child-resistant. Keep out of reach of children.
- In case of accidental overdose or ingestion, call a doctor or poison control center immediately.
- Do not use the product if the seal is broken or missing.

**Inactive Ingredients**

Vanillin, L-Alanine, Mineral Oil, Dimethylsiloxane, Methylparaben, Propylparaben, Propylparaben, PVP, Isoeucalyptol, Citrus, Citronellal, Citronellol, Geranylacetate, Myristic, Propylene Glycol, Ammonium Methacrylate, Purified Water, Laureth-9, Stearic Acid, Benzylalcohol, Benzoin Oil.

**USAGE METHOD:**

- Open and apply the unit of use within the cream will be applied.
- Apply a generous amount of cream onto the skin.
- Cover the open-ended area with plastic wrap for approximately 40-60 minutes.
- Remove plastic wrap, gently pat to help cream absorb. Wipe off residue, rest 15-20 mins for maximum numbness before proceeding.

**Distributed by:**

Pressy Inc.  
1942 Broadway St. Ste 214C  
Boulder, CO 80501  
Tel: 303-239-4100

**Manufactured by:**

Daoguang (Beijing) Pharmaceutical Co., Ltd.  
Daoguang Hanger, China 102201  
www.pressy.com

74115 1188710 11 g  
Net Wt. 1.75oz (50g)

**Pressy** EXTERNAL

**Tattoo**

**Numbing Cream**


**MAXIMUM STRENGTH PAIN RELIEF**

**NET: 50g/1.75oz**

MFG: 12/11/2023  
EXP: 12/10/2026  
LOT: 20231211-01

尺寸: 61x61x48mm

 PANTONE 485 C





# PRESSY TATTOO NUMBING CREAM

numbing cream cream

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83887-001
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	5 g in 100 g

## Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARETH-2 (UNII: V56DFE46J5)	
STEARETH-21 (UNII: 53J3F32P58)	

<b>CETOSTEARYL ALCOHOL</b> (UNII: 2DMT128M1S)
<b>AMMONIUM METHACRYLATE</b> (UNII: J2243103QO)
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)
<b>.ALPHA.-TOCOPHEROL</b> (UNII: H4N855PNZ1)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)
<b>IODOPROPYNYL BUTYLCARBAMATE</b> (UNII: 603P14DHEB)
<b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>LAUROCAPRAM</b> (UNII: 1F3X9DRV9X)
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)
<b>ISOPROPYL MYRISTATE</b> (UNII: 0RE8K4LNJS)
<b>ALLANTOIN</b> (UNII: 344S277G0Z)
<b>ROSE OIL</b> (UNII: WUB68Y35M7)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83887-001-01	30 g in 1 TUBE; Type 0: Not a Combination Product	12/22/2023	
2	NDC:83887-001-02	50 g in 1 BOTTLE; Type 0: Not a Combination Product	12/22/2023	
3	NDC:83887-001-03	60 g in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	12/22/2023	
4	NDC:83887-001-04	100 g in 1 BOTTLE; Type 0: Not a Combination Product	12/22/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	12/22/2023	

**Labeler** - Changsha Beiliang E-commerce Co., Ltd (632138945)

### Establishment

Name	Address	ID/FEI	Business Operations
Changsha Beiliang E-commerce Co., Ltd		632138945	manufacture(83887-001) , label(83887-001)

Revised: 12/2023

Changsha Beiliang E-commerce Co., Ltd