

# **BLOWFISH- aspirin, caffeine tablet, effervescent**

## **Rally Labs LLC**

-----

### **Drug Facts**

#### **Active ingredients**

Aspirin 500 mg (NSAID\*)  
Caffeine 60 mg

\*Nonsteroidal anti-inflammatory drug

#### **Purpose**

Aspirin.....Pain reliever

Caffeine...Alertness aid/Pain reliever aid

#### **Uses**

- for the temporary relief of minor aches and pains associated with a hangover
- helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover
- also for the temporary relief of headaches or body aches and pains alone
  
- for the temporary relief of headache pain
- also temporarily relieves minor aches and pains due to: backache muscle aches arthritis menstrual cramps hangover
  
- for the temporary relief of muscle aches and pain
- helps restore mental alertness when experiencing fatigue

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product. If changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reyes syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

#### **Stomach bleeding warning:**

This product contains a nonsteroidal anti-inflammatory drug (NSAID) which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever inducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- diabetes
- gout
- arthritis

### **Caffeine warning**

- the recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heart beat.
- for occasional use only. Do not use for more than 2 days for a hangover unless directed by a doctor. Not intended for use as a substitute for sleep. If fatigue or drowsiness persists or continues to occur, consult a doctor.

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain lasts for more than 10 days or gets worse
- new symptoms occur

- redness or swelling is present
- ringing in the ears or a loss of hearing occurs

### **If pregnant or breastfeeding**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- upon waking, fully dissolve 2 tablets in 16 oz. of water and drink
- do not exceed recommended dosage

<b>Adults and children 12 years and over</b> (up to 60 years of age)	2 tablets every 6 hours, as needed, or as directed by a doctor. Do not exceed 8 tablets in 24 hours
<b>Adults 60 years and over</b>	2 tablets every 6 hours, as needed, or as directed by a doctor. Do not exceed 4 tablets in 24 hours
<b>Children under 12 years</b>	Do not use

### **Other information**

- **each tablet contains:** sodium 406 mg
- phenylketonurics: contains phenylalanine 12.6 mg per tablet
- store at room temperature (59–86°F). Protect from excessive heat.

### **Inactive ingredients**

acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

### **Questions or Comments?**

**1-800-970-1793 (weekdays)**

**Blowfish for Hangovers**

Blowfish for Hangovers

Alertness Aid - Caffeine

Pain Reliever - Aspirin **(NSAID)**

12 effervescent tablets

Lemon Flavor





PN: 6052B R: 10/16

Barcode FPO

forhangovers.com  
1-800-970-1793  
New York, NY 10038  
Distributed by Rally Labs LLC

**WE GOT THIS.**

This is probably the best decision you've made in the last 24 hours. Blowfish has one job: fix your hangover. It's backed by science and guaranteed to work. Just drop in water and drink. We'll take it from there.

**blowfish**  
FOR HANGOVERS

**Lemon Flavor**  
Alertness Aid - Caffeine  
Pain Reliever - Aspirin (NSAID)



12 effervescent tablets

**Drug Facts (continued)**  
Other information:  
• each tablet contains sodium 406mg  
• phenylephrine 12.5 mg per tablet  
• store at room temperature (59-86°F).  
Protect from excessive heat.  
**Residue Ingredients**  
aspartame, potassium, erythrosorbate, sodium citrate, sodium gluconate, sodium hydroxide, sodium lauryl sulfate, sodium phosphate dibasic, sodium phosphate monobasic, sodium benzoate, sodium bicarbonate.  
**Directions or Components?**  
1-800-970-1793 (toll-free)

DO NOT USE IF PRINTED PACKETS ARE TORN OR PUNCTURED

forhangovers.com

**Drug Facts**  
**Active ingredients**      **Purpose**  
**(In each tablet)**  
Aspirin 500 mg (NSAID)\*..... Pain reliever  
Alertness Aid/..... Alertness aid/  
Caffeine 60 mg..... Pain reliever aid  
\*Nonsteroidal anti-inflammatory drug

**Uses**  
• for the temporary relief of minor aches and pains associated with a hangover  
• helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover  
• also for the temporary relief of headaches or body aches and pains alone

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock  
Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing

**Drug Facts (continued)**  
prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use** if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**  
• stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for • diabetes • gout • arthritis


**Caffeine warning**  
• the recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat. • for occasional use only.

**Drug Facts (continued)**  
Stop use and ask a doctor if • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain lasts for more than 10 days or gets worse • new symptoms occur • redness or swelling is present • ringing in the ears or a loss of hearing occurs

If pregnant or breastfeeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
• Upon waking, fully dissolve 1 or 2 tablets in 16 oz. of water and drink.  
• Adults and children 12 years and over: 1 or 2 tablets every 6 hours, as needed. Do not exceed 8 tablets in 24 hours. • Children under 12 years: do not use





**Drug Facts**

Active ingredients (in each tablet)	Purpose
Aspirin 500 mg (NSAID*)	Pain reliever
Caffeine 60 mg	Pain reliever aid / Alertness aid

\*Nonsteroidal anti-inflammatory drug

**Uses**

- For the temporary relief of headache pain
- Also temporarily relieves minor aches and pains due to:
  - backache • muscle aches • arthritis • menstrual cramps • hangover

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat. For occasional use only.

**Do not use** if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer

**Drug Facts (continued)**

**Ask a doctor before use if** • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for • diabetes • gout • arthritis

**Stop use and ask a doctor if** • an allergic reaction occurs. Seek medical help right away • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain lasts for more than 10 days or gets worse • new symptoms occur • redness or swelling is present • ringing in the ears or a loss of hearing occurs.

**If pregnant or breastfeeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- Fully dissolve 1 or 2 tablets in 16 oz. of water and drink
- Adults and children 12 years and older: 1 or 2 tablets every 6 hours as needed. Do not exceed 8 tablets in 24 hours.
- Children under 12 years: do not use

**Other Information** • each tablet contains: sodium 406mg • phenylketonurics: contains phenylalanine 12.6 mg per tablet • store at room temperature (59-86°F). Protect from excessive heat.

**Inactive Ingredients** acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Blowfish Sport**

Blowfish Sport

Caffeinated Pain Relief

20 effervescent tablets

Lemon Flavor

Alertness Aid - Caffeine

Pain Reliever - Aspirin (NSAID)



<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each tablet)</b> <b>Purpose</b>  Aspirin 500 mg (NSAID*) .....Pain reliever  Caffeine 60 mg .....Pain reliever aid / Alertness aid  *Nonsteroidal anti-inflammatory drug</p> <p><b>Uses</b>  • For the temporary relief of muscle aches and pain  • Helps restore mental alertness when experiencing fatigue</p> <p><b>Warnings</b>  <b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  <b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock  <b>Stomach bleeding warning:</b> This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed  <b>Caffeine warning</b> The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat. For occasional use only.  <b>Do not use</b> if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Ask a doctor before use if</b> • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet</p> <p><b>Ask a doctor or pharmacist before use if you are</b> taking a prescription drug for • diabetes • gout • arthritis</p> <p><b>Stop use and ask a doctor if</b> • an allergic reaction occurs. Seek medical help right away • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain lasts for more than 10 days or gets worse • new symptoms occur • redness or swelling is present • ringing in the ears or a loss of hearing occurs.</p> <p><b>If pregnant or breastfeeding,</b> ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b>  • Fully dissolve 1 or 2 tablets in 16 oz. of water and drink  • Adults and children 12 years and older: 1 or 2 tablets every 6 hours as needed. Do not exceed 8 tablets in 24 hours.  • Children under 12 years: do not use</p> <p><b>Other Information</b> • each tablet contains: sodium 406mg • phenylketonurics: contains phenylalanine 12.6 mg per tablet • store at room temperature (59-86° F). Protect from excessive heat.</p> <p><b>Inactive Ingredients</b> acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate</p>
--	--

## BLOWFISH

aspirin, caffeine tablet, effervescent

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:75920-0464
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	60 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>ASPARTAME</b> (UNII: Z0H242BBR1)	
<b>DOCUSATE SODIUM</b> (UNII: F05Q2T2JA0)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
--------------	-------	--------------	----------

<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>	LEMON	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75920-0464-1	2 in 1 BOX; Type 0: Not a Combination Product	07/23/2011	
2	NDC:75920-0464-8	12 in 1 BOX; Type 0: Not a Combination Product	07/23/2011	
3	NDC:75920-0464-2	20 in 1 BAG; Type 0: Not a Combination Product	11/30/2017	
4	NDC:75920-0464-4	40 in 1 BAG; Type 0: Not a Combination Product	11/30/2017	
5	NDC:75920-0464-6	4 in 1 BOX; Type 0: Not a Combination Product	02/01/2020	
6	NDC:75920-0464-5	20 in 1 BOX; Type 0: Not a Combination Product	12/28/2020	
7	NDC:75920-0464-7	20 in 1 BOX; Type 0: Not a Combination Product	12/28/2020	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/23/2011	

**Labeler** - Rally Labs LLC (965453108)

### Establishment

Name	Address	ID/FEI	Business Operations
Tower Laboratories Ltd		001587203	manufacture(75920-0464)

Revised: 12/2025

Rally Labs LLC