BIOTIC-PLUS- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, lidocaine ointment Provision Medical

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

TAMPER EVIDENT. DO NOT USE IF PACKET IS TORN OR CUT.

advanced Stops Pain and Prevents Infection for Cuts and Scrapes

BIOTIC-PLUS 0.5g

Active ingredients

(in each gram) Bacitracin zinc (400 units Bacitracin) Neomycin sulfate (3.5 mg neomycin) Polymyxin B sulfate (5000 units) Lidocaine .40 mg

Purpose

First Aid Antibiotic First Aid Antibiotic First Aid Antibiotic Anesthetic

Uses

First aid to help prevent infection and provide temporary relief in minor cuts, scrapes, and burns

The Provision First Aid Line

Manufactured for Provision Medical Products Palm Desert, CA 92211

Warnings

For external use only.

Keep out of reach of children

If ingested, contact a Poison Control Center directly.

Do not use

- in eyes or over large areas of the body
- more than 1 week unless directed by a doctor
- if allergic to ingredients.

Stop use, ask a doctor

- in case of deep or puncture wounds, animal bites or serious burns
- if conditions worsen or last more than 7 days
- if rash or allergic reaction occurs

Directions

- adults & children 2 years & older: clean affected area, apply amount (equal to surface area of a tip of finger) on area 1-3 times daily. May be covered with a sterile bandage.
- children under 2: do not use, ask a doctor

Inactive ingredients

white petrolatum



EXP.	<i>Inact</i> white	<i>Warning</i> For extern Keep out of If ingested, Control Cent use ■ in ey areas of the than 1 weel by a doctor ingredients. a doctor ■ or puncture bites or seri conditions v more than 7 or allergic r <i>Direction</i> . children 2 ye affected are (equal to sur of finger) on a May be cove bandage. 2: do not us
LOT	Inactive ingredients white petrolatum	<i>ngs</i> ernal use enter directh of reach of enter directh newes or ov the body ■ eek unless tor ■ if allents serious burn serious burn serious burn sworsen o ic reaction of area, apply surface area on area 1-3 ti overed with
	S	only. children a Poison y. Do not y. Do not r large directed argic to r last r last r last if rash occurs boccurs boccurs anount a of a tip mes daily. a sterile h under

Product Informat	ion						
Product T ype		HUMAN OTC DRUG	Item Co	Item Code (Source) NDC		DC:69103-2400	
Route of Administrat	tion	TOPICAL					
Active Ingredient	/Active Moi	ety					
	Basis of	Strength	Strength				
bacitracin zinc (UNII: 8	89Y4M234ES) (bacitracin - UNII:58H6RW0	O52I)	bacitracin		400 [iU] in 1 g	
neomycin sulfate (UNI	I: 057Y626693)	(neomycin - UNII:I16QD7	X297)	neomycin		3.5 mg in 1 g	
polymyxin b sulfate (U	JNII: 19371312D4	po lymyxin l	b	5000 [iU] in 1 g			
lidocaine (UNII: 98PI20	0987) (lidocair	ne - UNII:98PI200987)		lido caine		.40 mg in 1 g	
petrolatum (UNII: 4T6F		Ingredient Name				trength	
0 0					36.1		
Packaging # Item Code		kage Description	Marketin	ng Start Date	Market	ing End Date	
00	Pac 0.5 g in 1		Marketin	ng Start Date	Market	ting End Date	
 <i>Item Code</i> NDC:69103-2400-1 Marketing Info 	0.5 g in 1 ormation		Marketin	ig Start Date	Market	ing End Date	
# Item Code	0.5 g in 1 ormation			ng Start Date Marketing Start		ting End Date rketing End Date	

Labeler - Provision Medical (036936831)

Registrant - Safetec of America, Inc. (874965262)

Establishment									
Name	Address	ID/FEI	Business Operations						
Safetec of America, Inc.		874965262	MANUFACTURE(69103-2400)						

Revised: 7/2014

Provision Medical