#### IVERMECTIN- ivermectin lotion Belmora LLC

-----

#### Ivermectin

Drug Facts

#### Active ingredient

Ivermectin 0.5%

#### Purpose

Lice treatment

#### Use

treats head lice

#### Warnings

#### For external use only. Use only on the scalp.

#### Do not use

- on children under 6 months of age
- near the eyes
- inside the nose, ear, mouth, or vagina
- on lice in eyebrows or eyelashes. See a doctor if lice are present in these areas.

#### Ask a doctor before use if you

- have any skin conditions or sensitivities
- are pregnant or plan to become pregnant. It is not known if product can harm your unborn baby
- are currently or planning to breastfeed. Avoid getting product on your breast to help prevent contact by your baby.

#### When using this product

- keep eyes tightly closed and protect eyes with a washcloth or towel
- if product gets into eyes, gently flush with water
- eye redness, soreness, or irritation can occur
- dandruff, dry skin, or burning sensation of the skin can occur.

#### Stop use and ask a doctor if

- breathing difficulty occurs
- eye irritation occurs
- skin or scalp irritation continues or infection occurs
- rash develops.

If pregnant or breastfeeding, ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## Directions

### Important: Read warnings on carton before use.

- children 6 months of age to under 12 years of age: an adult should supervise use.
- adults and children 6 months of age and over:

## Inspect

- all household members should be checked by another person for lice and/or nits (eggs)
- use a magnifying glass in bright light to help you see the lice and/or nits (eggs)
- use a tool, such as a comb or two unsharpened pencils, to lift and part the hair
- look for tiny nits near the scalp, beginning at the back of the neck and behind the ears
- examine small sections of the hair (1-2 inches wide) at a time
- unlike dandruff, nits stick to the hair. Dandruff should move when lightly touched.
- If either lice or nits (eggs) are found, treat with product.

## Treat

- your hair and scalp must be **DRY** before applying product
- use the top of cap to break the tamper seal on the tube
- apply product directly to dry hair and scalp
- completely cover your scalp and hair closest to the scalp first, and then apply outwards towards the ends of your hair
- use only amount needed to completely cover hair and scalp, up to 1 entire tube, and discard the remaining
- rub product throughout your hair
- it is important to completely cover your entire head so that all lice and eggs are exposed to the lotion. Be sure that each hair is coated from the scalp to the tip.

## Wait and rinse

- allow product to stay on your hair and scalp for 10 minutes after it has been applied. Use a timer or clock. Start timing after you have completely covered your hair and scalp with product.
- After 10 minutes, rinse product completely from your hair and scalp using only water
- after rinsing, dry and style as usual. Wait 24 hours before applying shampoo.

# After treatment

## wash your hands after applying product

- nit combing is not necessary when treating with product for it to work, but if desired, a fine-tooth comb or special nit comb may be used to remove dead lice and nits
- this is a single use product. Discard tube after use.
- Do not use again on the same person and same lice infestation without talking to a

healthcare provider first

- if infestation continues, see a doctor for other treatments
- machine wash any bedding and clothing used by anyone having lice. Machine wash at high temperatures (150°F) and tumble in a hot dryer for 20 minutes.
- After finishing treatment with lice medicine, check everyone in your family for lice after one week. Consider treatment for those who have lice.

## Other information

- TAMPER EVIDENT: DO NOT USE IF THE TAMPER SEAL ON THE TUBE HAS BEEN BROKEN.
- store at room temperature 68°-77°F (20°-25°C)
- do not freeze
- keep carton and see Consumer Information Leaflet and warnings before use.

## Inactive ingredients

butylated hydroxyanisole, castor oil, cetyl alcohol, citric acid anhydrous, crodalan AWS, cyclomethicone, glycerin, imidurea, lanolin alcohols, methylparaben, oleyl alcohol, olive oil, propylene glycol, propylparaben, purified water, shea butter, sodium citrate anhydrous, sorbitan tristearate, and stearyl alcohol.

## Questions?

1-888-779-2877

## Distributed by:

Belmora LLC, 3033 Wilson Blvd. Suite 700, Arlington, VA 22201

## **PRINCIPAL DISPLAY PANEL - 117 g Tube Carton**

NDC 27854-334-01

Belmora LLC

SHEA IVERMECTIN

NET WT. 4 oz (117 g)

%5'0 'NOLLOT

**LICE TREATMENT** SHEA IVERMECTIN

m.

4

12

NDC 27854-334-01

# Ŷ SHEA IVERMECTIN LICE TREATMENT LOTION, 0.5%

#### **NO SECOND APPLICATION REQUIRED**

• NO NIT COMBING REQUIRED

• NOW AVAILABLE WITHOUT PRESCRIPTION

• IVERMECTIN LOTION KILLS LICE

#### 626 2245848-1024-01

#### SHEA IVERMECTIN LICE TREATMENT LOTION. 0.5%

#### **Drug Facts**

Active ingredient Purpose .Lice trea **Use** treats head lice

#### *Warnings* For external use only. Use only on the scalp.

Do not use a on children under 6 months of age a near the eyes a inside the nose. ear, mouth, or vagina ■ on lice in eyebrows or eyelashes. See a doctor if lice e present in these areas.

Ask a doctor before use if you a have any skin conditions or sensitivities are pregnant or plan to become pregnant. It is not known if product can harm your unborn baby are currently or planning to breastfeed. Avoid getting product on your breast to help prevent contact by

your baby When using this product the keep eyes tightly closed and protect eyes with a washcloth or towel the if product gets into eyes, gently flush with water the eye redness, soreness, or irritation can occur a dandruff, dry skin, or burning ensation of the skin can occur

Stop use and ask a doctor if m breathing difficulty occurs ■ eye irritation occurs skin or scalp irritation continues or infection occurs a rash develops.

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children. If swallow ed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

#### SHEA IVERMECTIN LICE TREATMENT" LOTION, 0.5%

#### Drug Facts (continued)

Directions: Important: Read warnings on carton before use

children 6 months of age to under 12 years of age: an adult should supervise use

adults and children 6 months of age and over:

Inspect all household members should be checked by Inspect all household memoers should be checked by another person for loce and/or risk (eggs) muse at magnifying glass in bright light to help you see the lice and/or risk (eggs) muse a tool, such as a comb or two unsharpened pencils, to lift and part the hair mlock for thy rits near the scalp, beginning at the back of the neck and behind the ears examine small sections of the hair (1-2 inches wide) at a time munike dandruf, rits stick to the hair. Dandruff should rown whole lidek to take de if the lice to are briese faund

move when lightly touched I feither lice or nits are found. treat with product.

Treat ■ your hair and scalp must be DRY before applying product 
use the top of cap to break the tamper seal on the

produce use rhe top of cap to dreak thereating is sea in the tube mapply roduct directly to dry hair and scale mompletely cover your scalp and hair closest to the scalp first, and then apply outwards to wards the ends of your hair muse only amount needed to completely cover hair and scalp, up to 1 entire theb, and discard the remaining much product throughout your hair mit is important to completely cover your prices head to get that the is and scame renewneed to the lefter entire head so that all lice and eggs are exposed to the lotion. Be sure that each hair is coated from the scalp to the tip.

Wait and rinse allow product to stay on your hair and scalp for 10 minutes after it has been applied. Use a timer or clock Start timing after you have completely covered your hair and scalp with product **a** After 10 minutes, rinse product completely from your hair and scalp using only water **a** after rinsing, dry and style as usual. Wait 24 hours before applying shampoo

After treatment 🔳 wash your hands after applying produc nit combing is not necessary when treating with product for it to work, but if desired, a fine-tooth comb or special nit comb may be used to remove dead lice and nits = this is a single use rray be used to remove dead lice and nice minism in its a single use product. Discard tube after use ID on out sea again on the same person and same lice infestation without talking to a healthcare provider first in infestation continues, see a doctor for other treatments in machine wash any bedding and dofting used by anyone having lice. Machine wash at high temperatures (150°) and tumble in a hot dyre for 20 minutes infinite finishing treatment with lice medicine, check everyone lowurg finishing the single doctor. in your family for lice after one week. Consider treatment for those who have lice.

#### SHEA IVERMECTIN LICE TREATMENT LOTION, 0.5%

NO SECOND APPLICATION REQUIRED

#### Drug Facts (continued)

Other Information TAMPER EVIDENT: DO NOT USE IF THE

TAMPER SEAL ON THE TUBE HAS BEEN BROKEN store at mom temperature 68°-77°F

store at room temperature 68"-77"-(20"-25°C) and onot freeze an keep carbon and see Consumer information Leaflet and warnings before use.

Inactive ingredients butylated hydroxyanisole, castor oil, cetyl alcohol, citric acid anhydrous, crodalan AWS, cyclomethicone, glycerin, imidurea, lanolin alcohols, methyloaraben, olevi alcohol, olive oil, propylene glycol, propylparaben, purfied water, shea butter, sodium citrate anhydrous, sorbitan tristearate, and stearyl alcohol.

 $\circ$ 

-

Questions? 1-888-779-2877

\*All trademarks are property of their respective owners.

Distributed by: Belmora LLC, 3033 Wilson Blvd. Suite 700, Arlington, VA 22201

Made in Israel



NET WT. 4 oz (117 g)

NET WT. 4 oz (117 g)

IVERMECTIN						
ivermectin lotion						
Product Information						
Product Type	HUMAN OTC DRUG	Item Code	(Source)	NDC:2	7854-334	
Route of Administration	TOPICAL					
Active Ingredient/Active Moiety						
Ingr	edient Name		Basis of Stre	ngth	Strength	
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)		IVERMECTIN		5 mg in 1 g		

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
CASTOR OIL (UNII: D5340Y2I9G)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CYCLOMETHICONE (UNII: NMQ347994Z)	
GLYCERIN (UNII: PDC6A3C0OX)	
IMIDUREA (UNII: M629807ATL)	
LANOLIN ALCOHOLS (UNII: 884C3FA9HE)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
OLEYL ALCOHOL (UNII: 172F2WN8DV)	
OLIVE OIL (UNII: 6UYK2W1W1E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC10H)	
WATER (UNII: 059QF0KO0R)	
SHEA BUTTER (UNII: K49155WL9Y)	
ANHYDROUS TRISODIUM CITRATE (UNII: RS7A450LGA)	
SORBITAN TRISTEARATE (UNII: 6LUM696811)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	

# **Product Characteristics**

Color	white (off-white;tan)	Score
Shape		Size
Flavor		Imprint Code
Contains		

# Packaging

#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:27854-334- 01	1 in 1 CARTON	08/12/2024	
1		117 g in 1 TUBE; Type 0: Not a Combination Product		
Μ	arketing	nformation		
M	arketing   Marketing Category	nformation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

# Labeler - Belmora LLC (112753244)

Establishment						
Name	Address	ID/FEI	<b>Business Operations</b>			

Revised: 10/2024

Belmora LLC