CURODONT REPAIR FLUORIDE PLUS- sodium fluoride sponge CREDENTIS AG

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

CURODONT REPAIR FLUORIDE PLUS - DISTRIBUTED BY STRAUMANN

ACTIVE INGREDIENTS

SODIUM FLUORIDE 0.05% (0.02% W/V FLUORIDE ION)

PURPOSE

ANTICAVITY

USES

AIDS IN THE PREVENTION OF DENTAL CAVITIES

- ANTICAVITY DENTAL RINSE
- RESTORES ENAMEL
- FOR THE TREATMENT OF WHITE SPOTS

WARNING

FOR TOPICAL INTRAORAL USE ONLY. FOR PROFESSIONAL OFFICE USE ONLY. THIS PRODUCT IS NOT INTENDED FOR HOME OR UNSUPERVISED CONSUMER USE.

KEEP OUT OF REACH OF CHILDREN.

DIRECTIONS

Usual dosage is one treatment per tooth. After prophylaxis treatment, remove the pellicle of the concerned tooth by using 2% sodium hypochlorite for 10 seconds. rinse the tooth surface with water. Remove inorganic deposits by pushing together the two cylinders and apply product with sponge on treatment area.

After treatment time of five minutes have patient expectorate residues. For maximum benefits for the prevention of caries, instruct patient not to rinse, eat or drink for 30 minutes. For patients with rampant or active caries, repeat treatment once or twice every six months.

OTHER INFORMATION

DO NOT USE IF SAFETY PACHAGING IS BROKEN OR MISSING.

SINGLE USE ONLY.

STORE IN A DRY AND COOL PLACE.

INACTIVE INGREDIENTS

WATER, CHLORHEXIDINE DIGLUCONATE, TROMETHAMINE, TREHALOSE DIHYDRATE, OLIGOPEPTIDE-104, HYDROXYPROPYL METHYLCELLULOSE

UNAMUARTE BY STRAUMANN

applicator with 0.1 ml rinse each. Contains 10 single packed single use Anticavity dental treatment rinse.

СПВОГОХ⊗



Water, Chlorhexidine digluconate, Tromethamine, Trehalose Dihydrate, Oligopeptide-104, Hydroxypropyl

Inactive ingredients

Store in a dry and cool place.

Do not use it safety packaging is broken or missing. Single use only.

Other information

35% phosphoric acid for 20 seconds. Rinse the tooth surface with water & dry the tooth surface. Activate product by pushing together the two cylinders and apply product with seconds on treatment laces.
After the stiment into dit two intuites have patient expectorate residues. For maximum benefits for the prevention of caries, instruct patient not to tinse, eat or drink for 30 minutes. For patients with rampant or active cates, repeat treatment once or twice every six months. Usual dosage is one treatment per tooth. After prophylaxis treatment, remove the pellidle of the concerned tooth by using 2% sodium hypochloitle for 20 seconds. After prophylaxis treatment, remove the pellidle of the concerned tooth by

Directions

Drug Facts Continued

or nome or unsupervised consumer use. Keep out or reach of children. For topical intracial use only. For Professional Office Use Only. This product is not intended Warning

- Helps prevent decay and white spots.
- Aort euswel strong.

 Helps prevent decay and cavities, which can keep
 - Anticavity dental rinse. Aids in the prevention of dental cavifies.

Sesn

Active ingredient Sodium fluoride ion) Anticavity Sodium fluoride 0.05% (0.02% w/v fluoride ion)

Drug Facts



CUROLOX®

Anticavity dental treatment rinse.

Contains 10 single packed single use applicator with 0.1 ml rinse each.

DISTRIBUTED BY STRAUMANN



10 × 0.1 ml (10 × 0.033 FL OZ)

Distributed by

Straumann USA, LLC 60 Minuteman Road Andover, MA 01810 United States info.usa@straumann.com Phone +1 978 747 2500

Manufactured by

credentis ag Dorfstrasse 69 CH-5210 Windisch Switzerland product@curodont.us Phone +41 56 560 20 44

How to use

The general principles for treatment and patient medication must be followed when using CURODONT™ REPAIR Fluoride Plus.

Preparation

- · Professional dental cleaning.
- Removal of the pellicle using 2% sodium hypochlorite for 20 seconds.
- Removal of inorganic deposits using 35% phosphoric acid for 20 seconds.
- · Rinse the tooth surface with water.
- . Dry the tooth surface.



1. Remove safety clip (black).



 Activate the CURODONT™ REPAIR Fluoride Plus applicator by pushing the two cylinders together.



 Remove the CURODONT™ REPAIR Fluoride Plus applicator from the protective cover.



above the lesion/proximal gap.

5. Allow the solution to diffuse for



5 minutes.
Discard the applicator unit after use.
The applicator is for single use only!

 Apply CURODONT™ REPAIR Fluoride Plus by squeezing out the sponge

Instruct patient not to rinse, eat or drink for 30 minutes.

Additional information

- Repeat treatment at least once or twice a year.
 For patients with rampant or active caries, repeat treatment once or twice every six months.
- · If contents have become frozen, warm to room temperature.
- · For topical intraoral use only.
- For Professional Office Use Only. This product is not intended for home or unsupervised consumer use.

P560.SU

REF

Active Ingredient/Active Moiety

Ingredient Name

Basis of

Strength

Product Information

sodium fluoride sponge

CURODONT REPAIR FLUORIDE PLUS

NDC:72247-105

ORAL

HUMAN OTC DRUG

Route of Administration

Active ingredient

Other information

Distributed by

Straumann USA, LLC

60 Minuteman Road

Andover, MA 01810

info.usa@straumann.com

Phone +1 978 747 2500

United States

Read directions for proper use.

Important

Warning

Sodium fluoride 0.05% (0.02% w/v fluoride ion).

Product Type

Item Code (Source)

Manufactured by

credentisaq

Switzerland

Dorfstrasse 69

CH-5210 Windisch

product@curodont.us

Phone +41 56 560 20 44

0.05 «

DISTRIBUTED BY STRAUMANN

Made in Switzerland

CUROLOX®

Do not use if safety packaging is broken or missing. Single use only. Store in a dry and cool place.

Office Use Only. This product is not intended for home or unsupervised consumer use.

Keep out of reach of children. If more than amount used for rinsing is accidentally swallowed, get medical help or contact a Poison Control Center right away. For topical intraoral use only. For Professional

MADDONT

Anticavity dental treatment rinse.

 Anticavity dental rinse. Aids in the prevention of dental cavities. Contains 1 single packed single use applicator with 0.1 ml (0.033 FL OZ) rinse.

Helps prevent decay and white spots.

Helps prevent decay and cavities, which can keep your ename! strong.

ì	SUDIUM MUNUFLUUKUPHUSPHALE (UNII: Cŏ IUJCZSQŲ) (FLUUKIDE IUN -	FLUORIDE ION	U.U5 g
ı	UNII:Q80 VPU408O)	FLUORIDE ION	in 100 mL

Inactive Ingredients		
Ingredient Name	Strength	
WATER (UNII: 059QF0KO0R)		
CHLORHEXIDINE GLUCONATE (UNII: MOR8 4MUD8 E)		
TROMETHAMINE (UNII: 023C2WHX2V)		
TREHALOSE DIHYDRATE (UNII: 7YIN7J07X4)		
OLIGOPEPTIDE-10 (UNII: Q46328TRNK)		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)		

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:72247-105-12	10 in 1 BOX	0 1/11/20 19	
1 NDC:72247-105-11	0.1 mL in 1 CYLINDER; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part355	0 1/11/20 19	

Labeler - CREDENTIS AG (485450345)

Registrant - CREDENTIS AG (485450345)

Establishment			
Name	Address	ID/FEI	Business Operations
System Kosmetik Produktionsgesellschaft fur kosmetische Erzeugnisse mbH		328825757	manufacture(72247-105)

Establishment			
Name	Address	ID/FEI	Business Operations
VALIDA		480120305	pack(72247-105)

Revised: 1/2020 CREDENTIS AG