

**MAX RELIEF JUNIOR- acetaminophen liquid**  
**Akron Pharma Inc.**

-----  
**MAX relief junior**  
**Dye-free Children's Pain Reliever and fever reducer**  
**Acetaminophen 160 mg per 5 mL Liquid**  
**Alcohol Free, Aspirin Free**  
**For Ages 2 to 11 years**

**Active ingredient (in each 5 mL)**

Acetaminophen 160 mg

**Purpose**

Pain Reliever/Fever Reducer

**Uses**

- temporarily
- reduces fever
- relieves minor aches and pains due to:
  - the common cold
  - flu
  - headache
  - sore throat
  - toothache

**Warnings**

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions.

**Soar throat warning:**if sore throat is severe, persists or more than 2 days, is accompanied or followed by fever headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if your child has** liver disease.

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin.

**When using this product: Do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**If pregnant or breast-feeding,** ask a health professional before use

**Overdose warning.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain direction or complete warnings for adult use**
- **Shake well before using**
- ml = milliliter
- find right dose on chart below
- if possible, use weight to dose; otherwise use age
- use only the enclosed dosing cup designed for use with this product. Do not use any other dosing device.
- if needed, repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give more than 5 days unless directed by doctor.

<b>Weight (lbs.)</b>	<b>Age (yrs.)</b>	<b>Dose (tsp or mL)</b>
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

Other Information store at room temperature 15 °-30 °C (59 °-86 °F). Protect from Freezing. Protect from light.

**Inactive ingredients:** amydrrous citric acid, bubble gum flavor, glycerin, polyethylene glycol400, punfied water, saccharin sodium, sodium benzoate, sodium citrate, sorbtol solution, sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

**Manufactured for:**

Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com

This product is not manufactured by or distributed by McNeil Consumer Healthcare,

owner of the registered trademark Tylenol Elixir.

NDC 71399-0021-4

**MAXrelief™ junior**

**Children's PAIN RELIEVER**  
and Fever Reducer

**Acetaminophen Oral Liquid**  
160 mg per 5 mL

- Alcohol Free
- Aspirin Free

For Ages 2 to 11 years

**Bubble Gum Flavor**

**Akron Pharma**



4 FL. OZ. (118 mL)

---

**Drug Facts**

**Active ingredient (in each 5 mL)**  
Acetaminophen 160 mg

**Purpose**  
Pain reliever / fever reducer

**Uses** temporarily ■ reduces fever ■ relieves minor aches and pains due to ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions** ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed (see overdose warning) ■ shake well before using ■ mL = milliliter ■ first right dose on chart below. If possible, use weight to dose; otherwise, use age. ■ remove the child protective cap and squeeze your child's dose into the dosing cup ■ repeat dose every 4 hours while symptoms last. ■ do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)	Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor	48 to 59	6 to 8	2 tsp or 10 mL
24 to 35	2 to 3	1 tsp or 5 mL	60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL	72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other information** store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Akron Pharma**  
Manufactured for:  
Akron Pharma, Inc.  
Fairfield, NJ 07704  
www.akronpharma.com



71399-0021-4

4 FL. OZ. (118 mL)

MAXrelief<sup>™</sup> junior  
Children's  
PAIN RELIEVER  
and Fever Reducer

NDC 71399-0021-4

MAXrelief<sup>™</sup> junior

Children's  
PAIN RELIEVER  
and Fever Reducer

Pain+Fever  
Acetaminophen  
Oral Liquid  
160 mg per 5 ml

- Alcohol Free
- Aspirin Free

For Ages 2 to 11 years

Bubble Gum  
Flavor

Akron Pharma

4 FL. OZ. (118 mL)

**Drug Facts**

**Active ingredient (in each 5 mL)** Purpose  
Acetaminophen 160 mg Pain reliever / fever reducer

**Uses temporarily:** reduces fever, relieves minor aches and pains due to: the common cold, flu, headache, sore throat, toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen.

**Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product,

ask a doctor before use if your child has liver disease.

ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if:** pain gets worse or lasts for more than 5 days; fever gets worse or lasts for more than 3 days; new symptoms occur; redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-422-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

this product does not contain directions or complete warnings for adult use

do not give more than directed (see overdose warning)

shake well before using

mL = milliliter

find right dose on chart below. If possible, use weight to dose; otherwise, use age.

remove the child protective cap and squeeze your child's dose into the dose cup

NDC 71399-0021-4

MAXrelief<sup>™</sup> junior

Children's  
PAIN RELIEVER  
and Fever Reducer

Pain+Fever  
Acetaminophen  
Oral Liquid  
160 mg per 5 ml

- Alcohol Free
- Aspirin Free

For Ages 2 to 11 years

Bubble Gum  
Flavor

Akron Pharma

4 FL. OZ. (118 mL)

**Drug Facts (continued)**

repeat dose every 4 hours while symptoms last  
do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
35 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other Information**

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients:** Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?**

call toll-free 1-877-225-6996.

\*\*This product is not manufactured or distributed by McNeil Consumer Products, owner of the registered trademark Tylenol®.

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Akron Pharma

Manufactured for:  
Akron Pharma, Inc.

Fairfield, NJ 07004

www.akronpharma.com

PNV0725



N 71399-0021-4 3

NDC 71399-0021-8  
**MAX**relief<sup>TM</sup>  
*junior*

**Children's  
PAIN RELIEVER**  
and Fever Reducer

**Pain+Fever**  
**Acetaminophen  
Oral Liquid**  
**160 mg per 5 mL**

- Alcohol Free
- Aspirin Free

For Ages 2 to 11 years

**Bubble Gum  
Flavor**

**Akron Pharma**

8 FL.OZ (236 mL)



**Drug Facts**

**Active ingredient (in each 5 mL)** Purpose

Acetaminophen 160 mg Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

■ this product does not contain directions or complete warnings for adult use

■ do not give more than directed (see overdose warning)

■ shake well before using ■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ remove the child protective cap and squeeze your child's dose into the dosing cup

**Drug Facts (continued)**

- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other Information**

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Akron Pharma**

Manufactured for:  
Akron Pharma, Inc.  
Fairfield, NJ 07004  
www.akronpharma.com

REU07/25



and Fever Reducer  
**PAIN RELIEVER**  
**Children's**  
**MAX**relief<sup>TM</sup>  
*junior*

NDC 71399-0021-8

**MAX**relief<sup>TM</sup>  
*junior*

**Children's  
PAIN RELIEVER**  
and Fever Reducer

**Pain+Fever**  
**Acetaminophen  
Oral Liquid**  
**160 mg per 5 mL**

- Alcohol Free
- Aspirin Free

For Ages 2 to 11 years

**Bubble Gum  
Flavor**

**Akron Pharma**

8 FL.OZ (236 mL)



**Drug Facts**

**Active ingredient (in each 5 mL)** Purpose

Acetaminophen 160 mg Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to: ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dose (see overdose warning)**

**Stop use and ask a doctor if** ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

■ this product does not contain directions or complete warnings for adult use

■ do not give more than directed (see overdose warning)

■ shake well before using ■ mL = milliliter

■ find right dose on chart below. If possible, use weight to dose; otherwise, use age.

■ remove the child protective cap and squeeze your child's dose into the dosing cup

NDC 71399-0021-8

**MAX**relief<sup>TM</sup>  
*junior*

**Children's  
PAIN RELIEVER**  
and Fever Reducer

**Pain+Fever**  
**Acetaminophen  
Oral Liquid**  
**160 mg per 5 mL**

- Alcohol Free
- Aspirin Free

For Ages 2 to 11 years

**Bubble Gum  
Flavor**

**Akron Pharma**

8 FL.OZ (236 mL)



**Drug Facts (continued)**

- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other Information**

store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir

**TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Akron Pharma**

Manufactured for:  
Akron Pharma, Inc.  
Fairfield, NJ 07004  
www.akronpharma.com

REU07/25



**Drug Facts**

**Active ingredient (in each 5 mL)** **Purpose**  
 Acetaminophen 160 mg .....Pain reliever / fever reducer

**Uses** temporarily: ■ reduces fever ■ relieves minor aches and pains due to:  
 ■ the common cold ■ flu ■ headache ■ sore throat ■ toothache

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: ■ more than 5 doses in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen.  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If your child is allergic to acetaminophen or any of the inactive ingredients in this product.  
**Ask a doctor before use if your child has liver disease.**  
**Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.**  
**When using this product, do not exceed recommended dose (see overdose warning)**  
**Stop use and ask a doctor if**  
 ■ pain gets worse or lasts for more than 5 days ■ fever gets worse or lasts for more than 3 days. ■ new symptoms occur redness or swelling is present These could be signs of a serious condition.  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.  
**Directions**  
 ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed (see overdose warning)  
 ■ shake well before using ■ mL = milliliter ■ find right dose on chart below. If possible, use weight to dose; otherwise, use age. ▶

NDC 71399-0021-6

# MAXrelief™ junior

## Children's PAIN RELIEVER and Fever Reducer

### Acetaminophen Oral Liquid

**160 mg per 5 mL**

● Alcohol Free  
● Aspirin Free

For Ages 2 to 11 years

**Bubble Gum Flavor**



**Akron Pharma** 16 FL.OZ (473 mL)

**Drug Facts (continued)**

- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours.

Weight (lbs.)	Age (yrs.)	Dose (tsp or mL)*
under 24	under 2	ask a doctor
24 to 35	2 to 3	1 tsp or 5 mL
36 to 47	4 to 5	1 1/2 tsp or 7.5 mL
48 to 59	6 to 8	2 tsp or 10 mL
60 to 71	9 to 10	2 1/2 tsp or 12.5 mL
72 to 95	11	3 tsp or 15 mL

\*or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

**Other Information**  
 store at room temperature 20°-25°C (68°-77°F). Protect from Freezing. Protect from Light.

**Inactive Ingredients** Anhydrous Citric Acid, Bubble gum flavor, F D & C Red No 40, Glycerin, Polyethylene Glycol 400, Purified Water, Sodium Benzoate, Sodium Citrate, Sorbitol Solution, Sucralose.

**Questions or comments?** call toll-free 1-877-225-6999.

\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Elixir®

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Akron Pharma**

Manufactured for:  
Akron Pharma, Inc.  
Fairfield, NJ 07004  
www.akronpharma.com

REL0725



N 3 71399 00216 7

MAX RELIEF JUNIOR			
acetaminophen liquid			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71399-0021
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
Inactive Ingredients			
	<b>Ingredient Name</b>		<b>Strength</b>
	ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
	GLYCERIN (UNII: PDC6A3C0OX)		
	POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)		
	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
	WATER (UNII: 059QF0KO0R)		
	SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
	SODIUM BENZOATE (UNII: OJ245FE5EU)		
	SODIUM CITRATE (UNII: 1Q73Q2JULR)		

<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399-0021-4	1 in 1 PACKAGE	05/14/2023	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:71399-0021-8	1 in 1 PACKAGE	05/14/2023	
2		236 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:71399-0021-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/14/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/14/2023	

**Labeler** - Akron Pharma Inc. (067878881)

Revised: 10/2025

Akron Pharma Inc.