

**MUCUS RELIEF SINUS SEVERE CONGESTION RELIEF- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated**  
**Chain Drug Marketing Association**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**QCH - 1167 - 2019-1007**

***Drug Facts***

<b><i>Active ingredients (in each caplet)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves:
  - nasal congestion
  - headache
  - minor aches and pains
  - sinus congestion and pressure
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product do not use more than directed**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or persistent headache

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information**

- store between 20-25°C (68-77°F) in a dry place
- retain outer package for complete product information

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

**PRINCIPAL DISPLAY PANEL**

Quality Choice®

NDC 63868-722-20

†Compare to Active Ingredients in Mucinex® Sinus-Max™ Severe Congestion Relief

Maximum Strength\*

Mucus Relief Sinus

Severe Congestion Relief

Pain Reliever, Expectorant, Nasal Decongestant

Acetaminophen, Guaifenesin, Phenylephrine HCl

For Relief of:

Sinus Congestion

Headache

Thins & Loosens Mucus

For Ages 12+

20 Caplets

NC

NC

NC

**Drug Facts**

**Active ingredients (in each caplet)**

**Purpose**

Acetaminophen 325 mg..... Pain reliever  
Guaifenesin 200 mg..... Expectorant  
Phenylephrine HCl 5 mg..... Nasal decongestant

**Uses** ■ temporarily relieves: ■ headache  
■ nasal congestion ■ minor aches and pains  
■ sinus congestion and pressure  
■ promotes nasal and/or sinus drainage

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 12 caplets in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks daily while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug combines an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ diabetes ■ liver disease ■ high blood pressure ■ thyroid disease

■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ▶

**Drug Facts (continued)**

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

**Stop use and ask a doctor if** ■ nervousness, dizziness, or sleeplessness occur

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

■ cough comes back or occurs with rash or persistent headache

These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

■ do not take more than directed (see Overdose warning)

■ do not take more than 12 caplets in any 24-hour period

■ adults and children 12 years and older, take 2 caplets every 4 hours

■ children under 12 years of age: do not use

**Other Information**

■ store between 20-25°C (68-77°F) in a dry place

■ retain outer package for complete product information

**Inactive ingredients**

colloidal silicon dioxide, croscarmellose sodium, croscopdone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, mannitol, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

<sup>†</sup>Compare to Active Ingredients in MUCINEX<sup>®</sup> Sinus-Max<sup>™</sup> Severe Congestion Relief

**Maximum Strength\* Mucus Relief Sinus Severe Congestion Relief**

Pain Reliever, Expectorant, Nasal Decongestant

NDC 63868-722-20

<sup>†</sup>Compare to Active Ingredients in MUCINEX<sup>®</sup> Sinus-Max<sup>™</sup> Severe Congestion Relief

**Maximum Strength\* Mucus Relief Sinus Severe Congestion Relief**

Pain Reliever, Expectorant, Nasal Decongestant

Acetaminophen, Guaifenesin, Phenylephrine HCl

**For Relief of:**  
Sinus Congestion  
Headache  
Thins & Loosens Mucus

For Ages 12+  
**Actual Size**



**20** Caplets

Maximum Strength\*  
**Mucus Relief Sinus Severe Congestion Relief**  
Pain Reliever, Expectorant, Nasal Decongestant



Distributed by C.D.M.A., Inc.©  
43157 W. Nine Mile  
Novi, MI 48376-0995  
www.qualitychoice.com  
Questions: 248-449-9300

†This product is not manufactured or distributed by Reckitt Benckiser, distributor of Mucinex<sup>®</sup> Sinus-Max<sup>™</sup> Severe Congestion Relief. F116702QCH\_R0

**DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN**

\*Per 4-hour dose, dose every 4 hours

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-722
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	AAA;1166
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-722-20	2 in 1 CARTON	06/10/2015	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	06/10/2015	

**Labeler** - Chain Drug Marketing Association (011920774)

Revised: 10/2019

Chain Drug Marketing Association