

SANAR NATURALS CHILDRENS PAIN RELIEVER- acetaminophen liquid
Sanvall Enterprises, Inc.

Sanar Naturals Childrens Pain Reliever

Active Ingredient

Acetaminophen 160mg

Purpose

Pain reliever/fever reducer

Uses

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not Use

With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a Doctor

your child has liver disease.

Ask a doctor or pharmacist before use

your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose

Stop use and ask a doctor

if pain gets worse or lasts more than 5 days, fever gets worse or lasts more than 3 days, new symptoms occur, redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children

Keep out of reach of children

in case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222- 1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

This product does not contain directions or complete warnings for adult use. Do not give more than directed (see overdose warning). Shake well before using. ml=milliliter. Find right dose on chart below. If possible, use weight to dose; otherwise, use age.

Remove the child protective cap and squeeze your child's dose into the dosing cup. Repeat dose every 4 hours while symptoms last, do not give more than 5 times in 24 hours.

Weight (lb)	Age (yr)	Dose (ml)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 ml
36-47	4-5 years	7.5 ml
48-59	6-8 years	10 ml
60-71	9-10 years	12.5 ml
72-95	11 years	15 ml

*or as directed by a doctor

Attention use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

Inactive Ingredients

Flavor, glycerin, methylparaben, propylene glycol, propylparaben, purified water, sodium carboxymethylcellulose and sucralose.

Other Information


www.sanarnaturals.com

Sanar Naturals Children's Pain + Fever

CONSOLIDATED LABEL CO. Service Rep : Danielle Dombroski
 (407) 339-2626 ext. 8161
 Pre-Press : Tim Biggins EP #: 293480

Changes or revisions, other than to correct errors or omissions by Consolidated Label, may result in additional charges.
THIS PROOF IS FOR COPY APPROVAL ONLY. ALL COLORS SHOWN ARE APPROXIMATE.

Item #: 416936-NJ Proof #: 4 2/2/2024 9:28:57 AM


Sanvall Enterprises, Inc. Sanar Naturals Childrens Acetaminophen 4fl oz - Grape Rev 2/24 Unwind Direction 

Customer Part #: LS324

Size: 2.2500 X 5.5000

Material: *Semi-Gloss FSC® Paper

Top Coating: Gloss Self Wound Lamination 404/B9077/ 100

Color Palette:  Cyan Magenta Yellow Black



Drug Facts
 (After reading the rest of label)

Warnings: Do not give to children under 2 years of age. Do not give to children 2 to 3 years of age unless directed by a doctor. Do not give to children 4 to 6 years of age unless directed by a doctor. Do not give to children 6 to 11 years of age unless directed by a doctor. Do not give to children 12 years of age and older unless directed by a doctor. Do not give to children with a fever, rash, or other symptoms of infection. Do not give to children with a history of liver disease. Do not give to children with a history of kidney disease. Do not give to children with a history of stomach or intestinal problems. Do not give to children with a history of bleeding or bruising. Do not give to children with a history of allergic reactions. Do not give to children with a history of asthma. Do not give to children with a history of diabetes. Do not give to children with a history of heart disease. Do not give to children with a history of high blood pressure. Do not give to children with a history of thyroid disease. Do not give to children with a history of seizures. Do not give to children with a history of diabetes. Do not give to children with a history of kidney disease. Do not give to children with a history of stomach or intestinal problems. Do not give to children with a history of bleeding or bruising. Do not give to children with a history of allergic reactions. Do not give to children with a history of asthma. Do not give to children with a history of diabetes. Do not give to children with a history of heart disease. Do not give to children with a history of high blood pressure. Do not give to children with a history of thyroid disease. Do not give to children with a history of seizures.

Directions: See back of box for directions. Do not give to children under 2 years of age. Do not give to children 2 to 3 years of age unless directed by a doctor. Do not give to children 4 to 6 years of age unless directed by a doctor. Do not give to children 6 to 11 years of age unless directed by a doctor. Do not give to children 12 years of age and older unless directed by a doctor.

Other information: See back of box for other information. Do not give to children under 2 years of age. Do not give to children 2 to 3 years of age unless directed by a doctor. Do not give to children 4 to 6 years of age unless directed by a doctor. Do not give to children 6 to 11 years of age unless directed by a doctor. Do not give to children 12 years of age and older unless directed by a doctor.

Sanar NATURALS
CHILDREN'S
 Acetaminophen (160 mg per 5 mL)
 Pain Reliever-Fever Reducer
PAIN + FEVER
 Ages 2-11 Years
 4 fl oz (120 mL)
 160 mg per 5 mL
 Grape Flavor

Weight (kg)	Age (yr)	Dose (mL)
10-15	2-3 years	2.5 mL
16-22	3-5 years	5 mL
23-30	6-7 years	7.5 mL
31-40	8-9 years	10 mL
41-50	10-11 years	12.5 mL
51-60	12 years	15 mL

Sanar NATURALS
 Sanar, FL 32979 (888)
 sanarnaturals.com

2.25 in

DI LINE
LINE
POST

SANAR NATURALS CHILDRENS PAIN RELIEVER

acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73386-324
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K679OBS311)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73386-324-04	120 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/05/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/05/2024	

Labeler - Sanvall Enterprises, Inc. (966905507)

Registrant - Sanvall Enterprises, Inc. (966905507)

