

IBUPROFEN- ibuprofen tablet, film coated
L.N.K. International, Inc.

Ibuprofen Tablets USP, 200 mg

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - toothache
 - the common cold
 - backache
 - headache
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- shock
- skin reddening
- asthma (wheezing)
- rash
- facial swelling
- hives
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - slurred speech
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen at 20

weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-844-428-2538

Principal Display Panel

24/7 life

BY **7-ELEVEN™**

Ibuprofen

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (**NSAID**)

compare to

ADVIL® TABLETS

active ingredient**

50

TABLETS

**Actual
Size**

QUALITY

• GUARANTEED •

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by
PF Consumer Healthcare 1 LLC, owner of the
registered trademark Advil® Tablets.

50844 REV1221A29115

DISTRIBUTED BY 7-ELEVEN, INC.
IRVING, TX 75063 WWW.7-ELEVEN.COM

Satisfaction Guaranteed 1-800-255-0711

No Print / No Varnish
Lot no. & Exp. date

0 52548 68997 1

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Ibuprofen

Ibuprofen Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

compare to
ADVI[®] TABLETS
active ingredient**

50

TABLETS

Actual Size

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients camauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate.

B-1205-291-15
REV1221A29115

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID) *..... Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - menstrual cramps
 - toothache
 - headache
 - muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- rash
- skin redness
- asthma (wheezing)
- hives
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older

• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk, if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have symptoms of heart problems or stroke:
 - chest pain
 - leg swelling
 - slurred speech
- weakness in one part or side of body

Convenience Valet 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50844-257 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength | | |
|---|--|---|----------------------|--------------------|
| IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM) | IBUPROFEN | 200 mg | | |
| Inactive Ingredients | | | | |
| Ingredient Name | Strength | | | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | | | | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | | | | |
| STARCH, CORN (UNII: O8232NY3SJ) | | | | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | | | | |
| ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK) | | | | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | | | | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | | | | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | | | | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | | | | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | | | | |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) | | | | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | | | | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | | | | |
| Product Characteristics | | | | |
| Color | brown | Score | no score | |
| Shape | ROUND | Size | 10mm | |
| Flavor | | Imprint Code | 44;291 | |
| Contains | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:50844-257-15 | 1 in 1 CARTON | 07/22/2019 | |
| 1 | | 50 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| ANDA | ANDA075010 | 07/22/2019 | | |

Labeler - L.N.K. International, Inc. (038154464)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 038154464 | pack(50844-257) |

| Establishment | | | |
|-------------------------|----------------|---------------|----------------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 832867837 | manufacture(50844-257) |

| Establishment | | | |
|-------------------------|----------------|---------------|----------------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 832867894 | manufacture(50844-257) |

| Establishment | | | |
|-------------------------|----------------|---------------|--|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 868734088 | manufacture(50844-257) , pack(50844-257) |

| Establishment | | | |
|-------------------------|----------------|---------------|----------------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 967626305 | pack(50844-257) |

| Establishment | | | |
|-------------------------|----------------|---------------|----------------------------|
| Name | Address | ID/FEI | Business Operations |
| LNK International, Inc. | | 117025878 | manufacture(50844-257) |

Revised: 4/2025

L.N.K. International, Inc.