

# PITAVASTATIN- pitavastatin calcium tablet, film coated AvKARE

## HIGHLIGHTS OF PRESCRIBING INFORMATION

### Pitavastatin Tablets

Rx only

These highlights do not include all the information needed to use PITAVASTATIN TABLETS safely and effectively. See full prescribing information for PITAVASTATIN TABLETS.

Initial U.S. Approval: 2009

## RECENT MAJOR CHANGES

Contraindications, Pregnancy and Lactation ( 4) Removed	03/2023
Warnings and Precautions, Immune-Mediated Necrotizing Myopathy ( 5.2)	03/2023

## INDICATIONS AND USAGE

Pitavastatin is a HMG-CoA reductase inhibitor (statin) indicated as an adjunct to diet to reduce low-density lipoprotein cholesterol (LDL-C) in: ( 1)

- Adult patients with primary hyperlipidemia.
- Adults with heterozygous familial hypercholesterolemia (HeFH).

## DOSAGE AND ADMINISTRATION

- Take orally once daily with or without food at the same time each day ( 2.1).
- For patients requiring a high-intensity statin or are unable to achieve their LDL-C goal receiving pitavastatin tablets 4 mg daily, prescribe alternative LDL-C-lowering treatment. ( 2.1).
- Assess LDL-C when clinically appropriate, as early as 4 weeks after initiation of pitavastatin tablets, and adjust the dosage if necessary. ( 2.1).
- Recommended dosage is 2 mg to 4 mg once daily. Maximum recommended dosage is 4 mg once daily. ( 2.2)
- Recommended starting dosage for patients with moderate and severe renal impairment and end-stage renal disease on hemodialysis is 1 mg once daily. Maximum recommended dosage is 2 mg once daily. ( 2.3)
- See full prescribing information for pitavastatin tablets dosage modifications due to drug interactions. ( 2.4)

## DOSAGE FORMS AND STRENGTHS

- Tablets: 1 mg, 2 mg, and 4 mg ( 3)

## CONTRAINDICATIONS

- Cyclosporine ( 4, 7)
- Active liver failure or decompensated cirrhosis ( 4, 5.3)
- Hypersensitivity to pitavastatin or any excipients in pitavastatin tablets ( 4)

## WARNINGS AND PRECAUTIONS

- *Myopathy and Rhabdomyolysis*: Risk factors include age 65 or greater, renal impairment, uncontrolled hypothyroidism, concomitant use with certain other drugs, and higher doses of pitavastatin tablets. Discontinue pitavastatin tablets if markedly elevated CK levels occur or myopathy is diagnosed or suspected. Temporarily discontinue pitavastatin tablets in patients experiencing an acute or serious condition at high risk of developing renal failure secondary to rhabdomyolysis. Inform patients of the risk of myopathy and rhabdomyolysis when starting or increasing the pitavastatin tablets dosage. Instruct patients to promptly report any unexplained muscle pain, tenderness or weakness particularly if accompanied by malaise or fever ( 5.1, 7, 8.5, 8.6).
- *Immune-Mediated Necrotizing Myopathy (IMNM)*: Rare reports of IMNM, an autoimmune myopathy, have been reported. Discontinue pitavastatin tablets if IMNM is suspected ( 5.2).
- *Hepatic Dysfunction*: Increases in serum transaminases have occurred, some persistent. Rare reports of fatal and non-fatal hepatic failure have occurred. Consider testing liver enzyme before initiating

therapy and as clinically indicated thereafter. If serious hepatic injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs, promptly discontinue pitavastatin tablets ( 5.3)

#### ADVERSE REACTIONS

- The most frequent adverse reactions (rate  $\geq 2\%$ ) were myalgia, constipation, back pain, diarrhea, and pain in extremity. ( 6)

To report SUSPECTED ADVERSE REACTIONS, contact AvKARE at 1-855-361-3993 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

#### DRUG INTERACTIONS

See full prescribing information for details regarding concomitant use of pitavastatin tablets with other drugs that increase the risk of myopathy and rhabdomyolysis. (2.4,7)

#### USE IN SPECIFIC POPULATIONS

- *Pregnancy*: May cause fetal harm. ( 8.1)
- *Lactation*: Breastfeeding not recommended during treatment with pitavastatin tablets

#### See 17 for PATIENT COUNSELING INFORMATION.

*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

**Revised: 12/2023**

---

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Important Dosage and Administration Information

#### 2.2 Recommended Dosage for Adults

#### 2.3 Recommended Dosage in Patients with Renal Impairment

#### 2.4 Pitavastatin Tablets Dosage Adjustments Due to Drug Interactions

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Myopathy and Rhabdomyolysis

#### 5.2 Immune-Mediated Necrotizing Myopathy

#### 5.3 Hepatic Dysfunction

#### 5.4 Increases in HbA1c and Fasting Serum Glucose Levels

### 6 ADVERSE REACTIONS

#### 6.1 Clinical Studies Experience

#### 6.2 Postmarketing Experience

### 7 DRUG INTERACTIONS

### 8 USE IN SPECIFIC POPULATIONS

#### 8.1 Pregnancy

#### 8.2 Lactation

#### 8.4 Pediatric Use

#### 8.5 Geriatric Use

#### 8.6 Renal Impairment

#### 8.7 Hepatic Impairment

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

#### 12.1 Mechanism of Action

**12.2 Pharmacodynamics**  
**12.3 Pharmacokinetics**  
**13 NONCLINICAL TOXICOLOGY**  
**13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**  
**14 CLINICAL STUDIES**  
**14.1 Primary Hyperlipidemia in Adult**  
**16 HOW SUPPLIED/STORAGE AND HANDLING**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Pitavastatin tablets are indicated as an adjunct to diet to reduce low-density lipoprotein cholesterol (LDL-C) in:

- Adult with primary hyperlipidemia.
- Adults with heterozygous familial hypercholesterolemia (HeFH).

*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Important Dosage and Administration Information**

- Take pitavastatin tablets orally once daily with or without food at the same time each day.
- For patients requiring a high-intensity statin or are unable to achieve their LDL-C goal receiving pitavastatin tablets 4 mg daily, prescribe alternative LDL-C-lowering treatment.
- Assess LDL-C when clinically appropriate, as early as 4 weeks after initiation of pitavastatin tablets, and adjust the dosage if necessary.

#### **2.2 Recommended Dosage for Adults**

- The recommended dosage range of pitavastatin tablets is 2 mg to 4 mg daily.
- The maximum recommended dosage is pitavastatin tablets 4 mg once daily.

#### **2.3 Recommended Dosage in Patients with Renal Impairment**

- The recommended starting dose for patients with moderate and severe renal impairment (estimated glomerular filtration rate 30 – 59 mL/minute/1.73 m<sup>2</sup> and 15 – 29 mL/minute/1.73 m<sup>2</sup>, respectively) and patients with end-stage renal disease receiving hemodialysis is pitavastatin tablets 1 mg once daily. The maximum recommended dose for these patients is pitavastatin tablets 2 mg once daily [see *Use in Specific Populations* (8.6)].
- There are no dosage adjustment recommendation for patients with mild renal

impairment.

## 2.4 Pitavastatin Tablets Dosage Adjustments Due to Drug Interactions

- In patients taking erythromycin, do not exceed pitavastatin tablets 1 mg once daily [*see Drug Interactions (7)*].
- In patients taking rifampin, do not exceed pitavastatin tablets 2 mg once daily [*see Drug Interactions (7)*].

*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

## 3 DOSAGE FORMS AND STRENGTHS

Tablets:

- 1 mg: White to off-white circular biconvex film-coated tablets. Debossed with the logo "OP" on one side and "57" on the other side of the tablet.
- 2 mg: White to off-white circular biconvex film-coated tablets. Debossed with the logo "OP" on one side and "58" on the other side of the tablet.
- 4 mg: White to off-white circular biconvex film-coated tablets. Debossed with the logo "OP" on one side and "59" on the other side of the tablet.

## 4 CONTRAINDICATIONS

Pitavastatin tablets are contraindicated in the following conditions:

- Concomitant use of cyclosporine [*see Drug Interactions (7)*].
- Acute liver failure or decompensated cirrhosis [*see Warning and Precautions (5.3)*].
- Hypersensitivity to pitavastatin or any excipients in pitavastatin tablets. Hypersensitivity reactions including angioedema, rash, pruritus, and urticaria have been reported with pitavastatin tablets [*see Adverse Reactions (6)*].

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Myopathy and Rhabdomyolysis

Pitavastatin tablets may cause myopathy (muscle pain, tenderness, or weakness with elevated creatine kinase [CK]) and rhabdomyolysis. Acute kidney injury secondary to myoglobinuria and rare fatalities have occurred as a result of rhabdomyolysis in patients treated with statin, including pitavastatin tablets.

#### Risk Factors for Myopathy

Risk factors for myopathy include age 65 years or greater, uncontrolled hypothyroidism, renal impairment, concomitant use of certain drugs (including other lipidlowering therapies), and higher pitavastatin tablets dosage [*see Dosage and Administration (2.2), Drug Interactions (7), and Use in Specific Populations (8.5,8.6)*]. Dosages of pitavastatin tablets greater than 4 mg once daily were associated with an increased risk for severe myopathy in premarketing clinical studies. The maximum recommended dose of pitavastatin tablets is 4 mg once daily .

## Steps to Prevent or Reduce the Risk of Myopathy and Rhabdomyolysis

Pitavastatin tablets are contraindicated in patients taking cyclosporine and not recommended in patients taking gemfibrozil [see *Contraindications (4) and Drug Interactions (7)*]. There are pitavastatin tablets dosage restrictions for patients taking erythromycin or rifampin [see *Dosage and Administration (2.4)*]. The following drugs when used concomitantly with pitavastatin tablets may also increase the risk of myopathy and rhabdomyolysis: lipid-modifying dosages of niacin (>1grams/day), fibrates, and colchicine [see *Drug Interactions (7)*].

Discontinue pitavastatin tablets if markedly elevated CK levels occur or if myopathy is either diagnosed or suspected. Muscle symptoms and CK elevations may resolve if pitavastatin tablets are discontinued. Temporarily discontinue pitavastatin tablets in patients experiencing an acute or serious condition at high risk of developing renal failure secondary to rhabdomyolysis, (e.g., sepsis; shock; severe hypovolemia; major surgery; trauma; severe metabolic, endocrine, or electrolyte disorders; or uncontrolled epilepsy).

Inform patients of the risk of myopathy and rhabdomyolysis when starting or increasing the pitavastatin tablets dosage. Instruct patients to promptly report any unexplained muscle pain, tenderness or weakness, particularly if accompanied by malaise or fever.

### **5.2 Immune-Mediated Necrotizing Myopathy**

There have been rare reports of immune-mediated necrotizing myopathy (IMNM), an autoimmune myopathy, associated with statin use, including reports of recurrence when the same or a different statin was administered. IMNM is characterized by proximal muscle weakness and elevated serum creatine kinase that persist despite discontinuation of statin treatment; positive anti-HMG CoA reductase antibody; muscle biopsy showing necrotizing myopathy; and improvement with immunosuppressive agents. Additional neuromuscular and serologic testing may be necessary. Treatment with immunosuppressive agents may be required. Discontinue pitavastatin tablets if IMNM is suspected.

### **5.3 Hepatic Dysfunction**

Increases in serum transaminases have been reported with pitavastatin tablets [see *Adverse Reactions (6)*]. In most cases, these changes appeared soon after initiation, were transient, were not accompanied by symptoms, and resolved or improved on continued therapy or after a brief interruption in therapy. There have been rare postmarketing reports of fatal and non-fatal hepatic failure in patients taking statins, including pitavastatin tablets.

Patients who consume substantial quantities of alcohol and/or have a history of liver disease may be at increased risk for hepatic injury.

Consider liver enzyme testing before the initiation of pitavastatin tablets and when clinically indicated thereafter. Pitavastatin tablets are contraindicated in patients with acute liver failure or decompensated cirrhosis [see *Contraindications (4)*]. If serious hepatic injury with clinical symptoms and/or hyperbilirubinemia or jaundice occurs, promptly discontinue pitavastatin tablets.

### **5.4 Increases in HbA1c and Fasting Serum Glucose Levels**

Increases in HbA1c and fasting serum glucose levels have been reported with statins, including pitavastatin tablets. Optimize lifestyle measures, including regular exercise, maintaining a healthy body weight, and making healthy food choices.

## 6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in other sections of the labeling:

- Myopathy and Rhabdomyolysis [see Warnings and Precautions (5.1)]
- Immune-Mediated Necrotizing Myopathy [see Warning and Precautions (5.2)]
- Hepatic Dysfunction [see Warning and Precautions (5.3)]
- Increases in HbA1c and Fasting Serum Glucose Levels [see Warning and Precautions (5.4)].

### 6.1 Clinical Studies Experience

Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of one drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice.

#### Adverse Reactions in Adults with Primary Hyperlipidemia

In 10 controlled clinical studies and 4 subsequent open-label extension studies, 3,291 adult patients with primary hyperlipidemia were administered pitavastatin tablets 1 mg to 4 mg daily. The mean continuous exposure of pitavastatin (1 mg to 4 mg) was 36.7 weeks (median 51.1 weeks). The mean age of the patients was 60.9 years (range; 18 years – 89 years) and 52% were females.

Approximately 93% of the patients were White, 7% were Asian/Indian, 0.2% were African American and 0.3% were Hispanic and other.

In controlled clinical studies and their open-label extensions, 3.9% (1 mg), 3.3% (2 mg), and 3.7% (4 mg) of pitavastatin tablets -treated patients were discontinued due to adverse reactions. The most common adverse reactions that led to treatment discontinuation were: elevated creatine phosphokinase (0.6% on 4 mg) and myalgia (0.5% on 4 mg).

Adverse reactions reported in  $\geq 2\%$  of patients in controlled clinical studies and at a rate greater than or equal to placebo are shown in Table 1. These studies had treatment duration of up to 12 weeks.

**Table 1. Adverse Reactions ( $\geq 2\%$  and  $\geq$  placebo) in Adults with Primary Hyperlipidemia in Studies up to 12 Weeks**

Adverse Reactions	Placebo (n= 208) %	Pitavastatin Tablets 1 mg (n=309) %	Pitavastatin Tablets 2 mg (n=951) %	Pitavastatin Tablets 4 mg (n=1540) %
Myalgia	1.4	1.9	2.8	3.1
Constipation	1.9	3.6	1.5	2.2
Diarrhea	1.9	2.6	1.5	1.9
Back Pain	2.9	3.9	1.8	1.4

Pain in extremity	1.9	2.3	0.6	0.9
-------------------	-----	-----	-----	-----

Other adverse reactions reported from clinical studies were arthralgia, headache, influenza, and nasopharyngitis.

Hypersensitivity reactions including rash, pruritus, and urticaria have been reported with Pitavastatin tablets.

The following laboratory abnormalities have been reported: elevated creatine phosphokinase, transaminases, alkaline phosphatase, bilirubin, and glucose.

#### Adverse Reactions in Adult HIV-Infected Patients with Dyslipidemia

In a double-blind, randomized, controlled, 52-week trial, 252 HIV-infected patients with dyslipidemia were treated with either pitavastatin tablets 4mg once daily (n=126) or another statin (n=126). All patients were taking antiretroviral therapy (excluding darunavir) and had HIV-1 RNA less than 200 copies/mL and CD4 count greater than 200 cell/ $\mu$ L for at least 3 months prior to randomization. The safety profile of pitavastatin tablets was generally consistent with that observed in the clinical trials described above. One patient (0.8%) treated with pitavastatin tablets had a peak creatine phosphokinase value exceeding 10 times the upper limit of normal (ULN), which resolved spontaneously. Four patients (3%) treated with pitavastatin tablets had at least one ALT value exceeding 3 times but less than 5 times the ULN, none of which led to drug discontinuation. Virologic failure was reported for four patients (3%) treated with pitavastatin tablets, defined as a confirmed measurement of HIV-1 RNA exceeding 200 copies/mL that was also more than a 2-fold increase from baseline.

*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

## **6.2 Postmarketing Experience**

The following adverse reactions have been identified during postapproval use of pitavastatin tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Gastrointestinal disorders:* abdominal discomfort, abdominal pain, dyspepsia, nausea

*General disorders:* asthenia, fatigue, malaise, dizziness

*Hepatobiliary disorders:* hepatitis, jaundice, fatal and non-fatal hepatic failure

*Immune system disorders:* angioedema, immune-mediated necrotizing myopathy associated with statin use.

*Metabolism and nutrition disorders:* increases in HbA1c, fasting serum glucose levels

*Musculoskeletal and connective tissue disorders:* muscle spasms, myopathy, rhabdomyolysis

*Nervous system disorders:* hypoesthesia, peripheral neuropathy

*Psychiatric disorders:* insomnia, depression. Rare reports of cognitive impairment (e.g., memory loss, forgetfulness, amnesia, memory impairment, confusion) associated with statin use. Cognitive impairment was generally nonserious, and reversible upon statin discontinuation, with variable times to symptom onset (1 day to years) and symptom resolution (median of 3 weeks).

Reproductive system and breast disorders:erectile dysfunction  
 Respiratory, thoracic and mediastinal disorders:interstitial lung disease  
 Skin and subcutaneous tissue disorders:lichen planus

To report SUSPECTED ADVERSE REACTIONS, contact AvKARE at 1-855-361-3993 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

## 7 DRUG INTERACTIONS

Table 2 includes a list of drugs that increase the risk of myopathy and rhabdomyolysis when administered concomitantly with pitavastatin tablets and instructions for preventing or managing drug interactions [see *Warnings and Precautions (5.1)*, *Clinical Pharmacology (12.3)*].

**Table 2. Drug Interactions that Increase the Risk of Myopathy and Rhabdomyolysis with pitavastatin tablets**

<b>Cyclosporine</b>	
<i>Clinical Impact:</i>	Cyclosporine significantly increases pitavastatin exposure and increases the risk of myopathy and rhabdomyolysis.
<i>Intervention:</i>	Concomitant use of cyclosporine with pitavastatin tablets is contraindicated [see <i>Contraindications (4)</i> ].
<b>Gemfibrozil</b>	
<i>Clinical Impact:</i>	Gemfibrozil may cause myopathy when given alone. The risk of myopathy and rhabdomyolysis is increased with concomitant use of gemfibrozil with statins, including pitavastatin tablets.
<i>Intervention:</i>	Avoid concomitant use of gemfibrozil with pitavastatin tablets.
<b>Erythromycin</b>	
<i>Clinical Impact:</i>	Erythromycin significantly increases pitavastatin exposure and increases the risk of myopathy and rhabdomyolysis.
<i>Intervention:</i>	In patients taking erythromycin, do not exceed pitavastatin tablets 1 mg once daily [see <i>Dosage and Administration (2.4)</i> ].
<b>Rifampin</b>	
<i>Clinical Impact:</i>	Rifampin significantly increases peak pitavastatin exposure and increases the risk of myopathy and rhabdomyolysis.
<i>Intervention:</i>	In patients taking rifampin, do not exceed pitavastatin tablets 2 mg once daily [see <i>Dosage and Administration (2.4)</i> ].
<b>Fibrates</b>	
<i>Clinical Impact:</i>	Fibrates may cause myopathy when given alone. The risk of myopathy and rhabdomyolysis is increased with concomitant use of fibrates with statins, including pitavastatin tablets.
<i>Intervention:</i>	Consider if the benefit of using fibrates concomitantly with pitavastatin tablets outweighs the increased risk of myopathy and rhabdomyolysis.
<b>Niacin</b>	
<i>Clinical Impact:</i>	The risk of myopathy and rhabdomyolysis may be increased with concomitant use of lipid-modifying doses ( $\geq 1$ g/day) of niacin with pitavastatin tablets.

<i>Intervention:</i>	Consider if the benefit of using lipid-modifying doses ( $\geq 1$ g/day) of niacin concomitantly with pitavastatin tablets outweighs the increased risk of myopathy and rhabdomyolysis.
<b>Colchicine</b>	
<i>Clinical Impact:</i>	Cases of myopathy and rhabdomyolysis have been reported with concomitant use of colchicine with statins, including pitavastatin tablets.
<i>Intervention:</i>	Consider the risk/benefit of concomitant use of colchicine with pitavastatin tablets.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Discontinue pitavastatin tablets when pregnancy is recognized. Alternatively, consider the ongoing therapeutic needs of the individual patient.

Pitavastatin tablets decreases synthesis of cholesterol and possibly other biologically active substances derived from cholesterol; therefore, pitavastatin may cause fetal harm when administered to pregnant patients based on the mechanism of action [see *Clinical Pharmacology* (12.1)].

In addition, treatment of hyperlipidemia is not generally necessary during pregnancy. Atherosclerosis is a chronic process and the discontinuation of lipid-lowering drugs during pregnancy should have little impact on the outcome of long-term therapy of primary hyperlipidemia for most patients.

Available data from case series and prospective and retrospective observational cohort studies over decades of use with statins in pregnant women have not identified a drug-associated risk of major congenital malformations. Published data from prospective and retrospective observational cohort studies with statin use in pregnant women are insufficient to determine if there is a drug associated risk of miscarriage [see *Data*].

In animal reproduction studies, no embryo-fetal toxicity or congenital malformations were observed in pregnant rats and rabbits orally administered pitavastatin during the period of organogenesis at doses which were 22 and 4 times, respectively, the human exposure at the maximum recommended human dosage (MRHD) of 4 mg, based on AUC [see *Data*].

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. Adverse outcomes in pregnancy occur regardless of the health of the mother or the use of medications. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

#### Data

##### *Human Data*

A Medicaid cohort linkage study of 1152 statin-exposed pregnant women compared to 886,996 controls did not find a significant teratogenic effect from maternal use of statins in the first trimester of pregnancy, after adjusting for potential confounders – including maternal age, diabetes mellitus, hypertension, obesity, and alcohol and tobacco

use – using propensity score-based methods. The relative risk of congenital malformations between the group with statin use and the group with no statin use in the first trimester was 1.07 (95% confidence interval 0.85 to 1.37) after controlling for confounders, particularly pre-existing diabetes mellitus. There were also no statistically significant increases in any of the organ-specific malformations assessed after accounting for confounders. In the majority of pregnancies, statin treatment was initiated prior to pregnancy and was discontinued at some point in the first trimester when pregnancy was identified.

Study limitations include reliance on physician coding to define the presence of a malformation, lack of control for certain confounders such as body mass index, use of prescription dispensing as verification for the use of a statin, and lack of information on non-live births.

### *Animal Data*

Embryo-fetal developmental studies were conducted in pregnant rats administered 3, 10, 30 mg/kg/day pitavastatin by oral gavage during organogenesis (gestation days 7-17). No adverse effects were observed at 3 mg/kg/day, systemic exposures 22 times human systemic exposure at 4 mg/day based on AUC.

Embryo-fetal developmental studies were conducted in pregnant rabbits administered 0.1, 0.3, 1 mg/kg/day pitavastatin by oral gavage during the period of fetal organogenesis (gestation days 6-18). Maternal toxicity consisting of reduced body weight and abortion was observed at all doses tested (4 times human systemic exposure at 4 mg/day based on AUC).

In perinatal/postnatal studies in pregnant rats given oral gavage doses of pitavastatin at 0.1, 0.3, 1, 3, 10, 30 mg/kg/day from organogenesis through weaning (gestation day 17 to lactation day 21), maternal toxicity consisting of mortality at  $\geq 0.3$  mg/kg/day and impaired lactation at all doses contributed to the decreased survival of neonates in all dose groups (0.1 mg/kg/day represents approximately 1 time human systemic exposure at 4 mg/day dose based on AUC).

Reproductive toxicity studies have shown that pitavastatin crosses the placenta in rats and is found in fetal tissues at  $\leq 36\%$  of maternal plasma concentrations following a single dose of 1 mg/kg/day during gestation (at the end of organogenesis).

## **8.2 Lactation**

### Risk Summary

There is no available information about the presence of pitavastatin in human or animal milk, the effects of the drug on the breastfed infant or the effects of the drug on milk production. However, it has been shown that another drug in this class passes into human milk. Statins, including pitavastatin tablets, decrease cholesterol synthesis and possibly the synthesis of other biologically active substances derived from cholesterol and may cause harm to the breastfed infant.

Because of the potential for serious adverse reactions in a breastfed infant, based upon the mechanism of action, advise patients that breastfeeding is not recommended during treatment with pitavastatin tablets. [see *Use in Specific Populations* (8.1), *Clinical Pharmacology* (12.1)]

## **8.4 Pediatric Use**

The safety and effectiveness of pitavastatin tablets have not been established in pediatric patients younger than 8 years of age with HeFH or in pediatric patients with other types of hyperlipidemia (other than HeFH).

*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

## **8.5 Geriatric Use**

In controlled clinical studies, 1,209 (43%) patients were 65 years and older. No significant differences in efficacy or safety were observed between geriatric patients and younger patients.

Advanced age ( $\geq 65$  years) is a risk factor for pitavastatin tablets-associated myopathy and rhabdomyolysis. Dose selection for a geriatric patient should be cautious, recognizing the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy and the higher risk of myopathy. Monitor geriatric patients receiving pitavastatin tablets for the increased risk of myopathy [see *Warnings and Precautions* (5.1)].

## **8.6 Renal Impairment**

Renal impairment is a risk factor for myopathy and rhabdomyolysis. Monitor all patients with renal impairment for development of myopathy. Due to the risk of myopathy, a dosage modification of pitavastatin tablets is recommended for patients with moderate and severe renal impairment (estimated glomerular filtration rate 30 – 59 mL/min/1.73 m<sup>2</sup> and 15 – 29 mL/min/1.73 m<sup>2</sup>, respectively), as well as end-stage renal disease receiving hemodialysis. [see *Dosage and Administration* (2.3), *Warnings and Precautions* (5.1), *Clinical Pharmacology* (12.3)].

## **8.7 Hepatic Impairment**

Pitavastatin is contraindicated in patients with active liver failure or decompensated cirrhosis [see *Contraindications* (4), *Warnings and Precautions* (5.3)].


## **10 OVERDOSAGE**

No specific treatment for pitavastatin tablets overdose is known. Contact Poison Control (1-800-222-1222) for latest recommendations. Hemodialysis is unlikely to be of benefit due to high protein binding ratio of pitavastatin tablets.

## **11 DESCRIPTION**

Pitavastatin tablets for oral use is an HMG-CoA reductase inhibitor.

The chemical name for pitavastatin is (+)monocalcium *bis*{(3R, 5S, 6 E)-7-[2-cyclopropyl-4-(4-fluorophenyl)-3-quinoly]-3,5-dihydroxy-6-heptenoate}. The structural formula is:



The empirical formula for pitavastatin is  $C_{50}H_{46}CaF_2N_2O_8$  and the molecular weight is 880.98. Pitavastatin is odorless and occurs as white to pale-yellow powder. It is freely soluble in pyridine, chloroform, dilute hydrochloric acid, and tetrahydrofuran, soluble in ethylene glycol, sparingly soluble in octanol, slightly soluble in methanol, very slightly soluble in water or ethanol, and practically insoluble in acetonitrile or diethyl ether. Pitavastatin is hygroscopic and slightly unstable in light.

Each film-coated tablet of pitavastatin tablets contains 1 mg, 2 mg, or 4 mg of pitavastatin, which is equivalent to 1.045 mg, 2.09 mg, or 4.18 mg, respectively, of pitavastatin calcium and the following inactive ingredients: crospovidone, lactose monohydrate, microcrystalline cellulose, magnesium oxide, magnesium stearate, povidone, sodium starch glycolate, and film coating containing the following inactive ingredients: hypromellose, lactose monohydrate, titanium dioxide, and triacetin.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Pitavastatin is an inhibitor of 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase, the enzyme that catalyzes the conversion of HMG-CoA to mevalonate, a rate-limiting step in the biosynthetic pathway for cholesterol. As a result, the expression of LDL-receptors followed by the uptake of LDL from blood to liver is accelerated and then the plasma TC decreases. Sustained inhibition of cholesterol synthesis in the liver also decreases levels of very low density lipoproteins.

### 12.2 Pharmacodynamics

#### Cardiac Electrophysiology

In a randomized, double-blind, placebo-controlled, 4-way parallel, active-comparator study with moxifloxacin in 174 healthy participants, pitavastatin tablets were not associated with clinically meaningful prolongation of the QTc interval or heart rate at daily doses up to 16 mg (4 times the recommended maximum dose of 4 mg daily).

### 12.3 Pharmacokinetics

#### Absorption

Pitavastatin peak plasma concentrations are achieved about 1 hour after oral administration. Both  $C_{max}$  and  $AUC_{0-inf}$  increased in an approximately dose-proportional manner for single pitavastatin tablets doses from 1 mg to 24 mg once daily. The

absolute bioavailability of pitavastatin oral solution is 51%. The  $C_{max}$  and AUC of pitavastatin did not differ following evening or morning drug administration. In healthy volunteers receiving 4 mg pitavastatin, the percent change from baseline for LDL-C following evening dosing was slightly greater than that following morning dosing. Pitavastatin was absorbed in the small intestine but very little in the colon.

#### *Effect of Food*

Administration of pitavastatin tablets with a high fat meal (50% fat content) decreases pitavastatin  $C_{max}$  by 43% but does not significantly reduce pitavastatin AUC.

#### Distribution

Pitavastatin is more than 99% protein bound in human plasma, mainly to albumin and alpha 1-acid glycoprotein, and the mean volume of distribution is approximately 148 L.

#### Elimination

##### *Metabolism*

The principal route of pitavastatin metabolism is glucuronidation via liver uridine 5'-diphosphate glucuronosyltransferase (UGT) with subsequent formation of pitavastatin lactone. There is only minimal metabolism by the cytochrome P450 system. Pitavastatin is marginally metabolized by CYP2C9 and to a lesser extent by CYP2C8. The major metabolite in human plasma is the lactone, which is formed via an ester-type pitavastatin glucuronide conjugate by UGTs (UGT1A3 and UGT2B7).

##### *Excretion*

A mean of 15% of radioactivity of orally administered, single 32 mg  $^{14}C$ -labeled pitavastatin dose was excreted in urine, whereas a mean of 79% of the dose was excreted in feces within 7 days. The mean plasma elimination half-life is approximately 12 hours.

#### Specific Populations

##### *Geriatric Patients*

In a pharmacokinetic study which compared healthy young and geriatric ( $\geq 65$  years) volunteers, pitavastatin  $C_{max}$  and AUC were 10 and 30% higher, respectively, in the geriatric patients [see *Use in Specific Populations* (8.5)]

##### *Pediatric Patients*

*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

##### *Male and Female Patients*

In a pharmacokinetic study, which compared healthy male and female volunteers, pitavastatin  $C_{max}$  and AUC were 60 and 54% higher, respectively in females.

##### *Racial or Ethnic Groups*

In pharmacokinetic studies pitavastatin  $C_{max}$  and AUC were 21 and 5% lower, respectively in Black or African American healthy volunteers compared with those of White healthy volunteers. In pharmacokinetic comparison between White volunteers and Japanese volunteers, there were no significant differences in  $C_{max}$  and AUC.

##### *Patients with Renal Impairment*

In patients with moderate renal impairment (estimated glomerular filtration rate of 30 –

59 mL/min/1.73 m<sup>2</sup>) and end stage renal disease receiving hemodialysis, pitavastatin AUC<sub>0-inf</sub> is 102% and 86% higher than those of healthy volunteers, respectively, while pitavastatin C<sub>max</sub> is 60% and 40% higher than those of healthy volunteers, respectively. Patients received hemodialysis immediately before pitavastatin dosing and did not undergo hemodialysis during the pharmacokinetic study. Hemodialysis patients have 33% and 36% increases in the mean unbound fraction of pitavastatin as compared to healthy volunteers and patients with moderate renal impairment, respectively [see Use in Specific Populations (8.6)].

In another pharmacokinetic study, patients with severe renal impairment (estimated glomerular filtration rate 15 – 29 mL/min/1.73 m<sup>2</sup>) not receiving hemodialysis were administered a single dose of pitavastatin tablets 4 mg. The AUC<sub>0-inf</sub> and the C<sub>max</sub> were 36% and 18% higher, respectively, compared with those of healthy volunteers. For both patients with severe renal impairment and healthy volunteers, the mean percentage of protein-unbound pitavastatin was approximately 0.6% [see Use in Specific Populations (8.6)].

The effect of mild renal impairment on pitavastatin exposure has not been studied.

#### *Patients with Hepatic Impairment*

The disposition of pitavastatin was compared in healthy volunteers and patients with various degrees of hepatic impairment. Pitavastatin C<sub>max</sub> and AUC<sub>inf</sub> in patients with moderate hepatic impairment (Child-Pugh B disease) was 2.7-fold and 3.8-fold higher, respectively as compared to healthy volunteers. In patients with mild hepatic impairment (Child-Pugh A disease), pitavastatin C<sub>max</sub> and AUC<sub>inf</sub> were 30% and 60% higher as compared to healthy volunteers. Mean pitavastatin half-life for moderate hepatic impairment, mild hepatic impairment, and healthy volunteers were 15, 10, and 8 hours, respectively [see Contraindications (4), Warnings and Precautions (5.3)].

#### Drug Interaction Studies

##### *Warfarin*

The steady-state pharmacodynamics (international normalized ratio [INR] and prothrombin time [PT]) and pharmacokinetics of warfarin in healthy volunteers were unaffected by the coadministration of pitavastatin tablets 4 mg daily.

Table 3 presents the effect of coadministered drugs on pitavastatin systemic exposure:

**Table 3. Effect of Coadministered Drugs on Pitavastatin Systemic Exposure**

<b>Coadministered drug</b>	<b>Dosage regimen</b>	<b>Change in AUC*</b>	<b>Change in C<sub>max</sub>*</b>
Cyclosporine	Pitavastatin 2 mg QD for 6 days + cyclosporine 2 mg/kg on Day 6	↑ 4.6 fold †	↑ 6.6 fold †
Erythromycin	Pitavastatin 4 mg single dose on Day 4 + erythromycin 500 mg 4 times daily for 6 days	↑ 2.8 fold †	↑ 3.6 fold †
Rifampin	Pitavastatin 4 mg QD + rifampin 600 mg QD for 5 days	↑ 29%	↑ 2.0 fold †
Atazanavir	Pitavastatin 4 mg QD + atazanavir 300 mg daily for 5 days	↑ 31%	↑ 60%
Darunavir/Ritonavir	Pitavastatin 4 mg QD on Days 1-5 and 12-16 + darunavir/ritonavir 800 mg/100 mg QD on Days 6-	↓ 26%	↓ 4%

	16		
Lopinavir/Ritonavir	Pitavastatin 4 mg QD on Days 1-5 and 20-24 + lopinavir/ritonavir 400 mg/100 mg BID on Days 9-24	↓ 20%	↓ 4 %
Gemfibrozil	Pitavastatin 4 mg QD + gemfibrozil 600 mg BID for 7 days	↑ 45%	↑ 31%
Fenofibrate	Pitavastatin 4 mg QD + fenofibrate 160 mg QD for 7 days	↑ 18%	↑ 11%
Ezetimibe	Pitavastatin 2 mg QD + ezetimibe 10 mg for 7 days	↓ 2%	↓ 0.2%
Enalapril	Pitavastatin 4 mg QD + enalapril 20 mg daily for 5 days	↑ 6%	↓ 7%
Digoxin	Pitavastatin 4 mg QD + digoxin 0.25 mg for 7 days	↑ 4%	↓ 9%
Diltiazem LA	Pitavastatin 4 mg QD on Days 1-5 and 11-15 and diltiazem LA 240 mg on Days 6-15	↑ 10%	↑ 15%
Grapefruit Juice	Pitavastatin 2 mg single dose on Day 3 + grapefruit juice for 4 days	↑ 15%	↓ 12%
Itraconazole	Pitavastatin 4 mg single dose on Day 4 + itraconazole 200 mg daily for 5 days	↓ 23%	↓ 22%

\* Data presented as x-fold change represent the ratio between coadministration and pitavastatin alone (i.e., 1-fold = no change). Data presented as % change represent % difference relative to pitavastatin alone (i.e., 0% = no change).

† Considered clinically significant [see *Dosage and Administration (2.4), Drug Interactions (7)*]

BID = twice daily; QD = once daily; LA = Long Acting

Table 4 presents the effect of pitavastatin coadministration on systemic exposure of other drugs:

**Table 4. Effect of Pitavastatin Coadministration on Systemic Exposure to Other Drugs**

<b>Coadministered drug</b>	<b>Dosage regimen</b>	<b>Change in AUC*</b>	<b>Change in C<sub>max</sub>*</b>
Atazanavir	Pitavastatin 4 mg QD + atazanavir 300 mg daily for 5 days	↑ 6%	↑ 13%
Darunavir	Pitavastatin 4 mg QD on Days 1-5 and 12-16 + darunavir/ritonavir 800 mg/100 mg QD on Days 6-16	↑ 3%	↑ 6%
Lopinavir	Pitavastatin 4 mg QD on Days 1-5 and 20-24 + lopinavir/ritonavir 400 mg/100 mg BID on Days 9-24	↓ 9%	↓ 7%
Ritonavir	Pitavastatin 4 mg QD on Days 1-5 and 20-24 + lopinavir/ritonavir 400 mg/100 mg BID on Days 9-24	↓ 11%	↓ 11%
	Pitavastatin 4 mg QD on Days 1-5 and 12-16 +		

Ritonavir	darunavir/ritonavir 800 mg/100 mg QD on Days 6-16		↑ 8%	↑ 2%
Enalapril	Pitavastatin 4 mg QD + enalapril 20 mg daily for 5 days	Enalapril	↑ 12%	↑ 12%
		Enalaprilat	↓ 1%	↓ 1%
Warfarin	Individualized maintenance dose of warfarin (2-7 mg) for 8 days + pitavastatin 4 mg QD for 9 days	R-warfarin	↑ 7%	↑ 3%
		S-warfarin	↑ 6%	↑ 3%
Ezetimibe	Pitavastatin 2 mg QD + ezetimibe 10 mg for 7 days		↑ 9%	↑ 2%
Digoxin	Pitavastatin 4 mg QD + digoxin 0.25 mg for 7 days		↓ 3%	↓ 4%
Diltiazem LA	Pitavastatin 4 mg QD on Days 1-5 and 11-15 and diltiazem LA 240 mg on Days 6-15		↓ 2%	↓ 7%
Rifampin	Pitavastatin 4 mg QD + rifampin 600 mg QD for 5 days		↓ 15%	↓ 18%

\*Data presented as % change represent % difference relative to the investigated drug alone (i.e., 0% = no change).

BID = twice daily; QD = once daily; LA = Long Acting

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 92-week carcinogenicity study in mice given pitavastatin, at the maximum tolerated dose of 75 mg/kg/day with systemic maximum exposures (AUC) 26 times the clinical maximum exposure at 4 mg daily, there was an absence of drug-related tumors.

In a 92-week carcinogenicity study in rats given pitavastatin at 1, 5, 25 mg/kg/day by oral gavage there was a significant increase in the incidence of thyroid follicular cell tumors at 25 mg/kg/day, which represents 295 times human systemic exposures based on AUC at the 4 mg daily maximum human dose.

In a 26-week transgenic mouse (Tg rasH2) carcinogenicity study where animals were given pitavastatin at 30, 75, and 150 mg/kg/day by oral gavage, no clinically significant tumors were observed.

Pitavastatin was not mutagenic in the Ames test with *Salmonella typhimurium* and *Escherichia coli* with and without metabolic activation, the micronucleus test following a single administration in mice and multiple administrations in rats, the unscheduled DNA synthesis test in rats, and a Comet assay in mice. In the chromosomal aberration test, clastogenicity was observed at the highest doses tested, which also elicited high levels of cytotoxicity.

Pitavastatin had no adverse effects on male and female rat fertility at oral doses of 10 and 30 mg/kg/day, respectively, at systemic exposures 56- and 354-times clinical exposure at 4 mg daily based on AUC.

Pitavastatin treatment in rabbits resulted in mortality in males and females given 1 mg/kg/day (30-times clinical systemic exposure at 4 mg daily based on AUC) and higher during a fertility study. Although the cause of death was not determined, rabbits had

gross signs of renal toxicity (kidneys whitened) indicative of possible ischemia. Lower doses (15-times human systemic exposure) did not show significant toxicity in adult males and females. However, decreased implantations, increased resorptions, and decreased viability of fetuses were observed.

## 14 CLINICAL STUDIES

### 14.1 Primary Hyperlipidemia in Adult

Study with Atorvastatin (Study 301)

Pitavastatin tablets were compared with Atorvastatin Calcium Tablets (referred to as atorvastatin) in a randomized, multicenter, double-blind, double-dummy, active-controlled, non-inferiority study of 817 adult patients with primary hyperlipidemia or mixed dyslipidemia. Patients entered a 6- to 8-week wash-out/dietary lead-in period and then were randomized to a 12-week treatment with either pitavastatin tablets or atorvastatin (Table 5). Non-inferiority of pitavastatin to a given dose of atorvastatin was considered to be demonstrated if the lower bound of the 95% CI for the mean treatment difference was greater than -6% for the mean percent change in LDL-C.

Lipid results are shown in Table 5. For the percent change from baseline to endpoint in LDL-C, pitavastatin tablets were non-inferior to atorvastatin for the two pairwise comparisons: pitavastatin tablets 2 mg vs. atorvastatin 10 mg and pitavastatin tablets 4 mg vs. atorvastatin 20 mg. Mean treatment differences (95% CI) were 0% (-3%, 3%) and 1% (-2%, 4%), respectively.

**Table 5. Lipid Response by Dose of Pitavastatin Tablets and Atorvastatin in Adult Patients with Primary Hyperlipidemia or Mixed Dyslipidemia in Study 301 (Mean % Change from Baseline at Week 12)**

Treatment	N	LDL-C	Apo-B	TC	TG	HDL-C	non-HDL-C
Pitavastatin Tablets 2 mg daily	315	-38	-30	-28	-14	4	-35
Pitavastatin Tablets 4 mg daily	298	-45	-35	-32	-19	5	-41
Atorvastatin 10 mg daily	102	-38	-39	-28	-18	3	-35
Atorvastatin 20 mg daily	102	-44	-36	-33	-22	2	-41

Study with Simvastatin (Study 302)

Pitavastatin tablets were compared with Simvastatin Tablets (referred to as simvastatin) in a randomized, multicenter, double-blind, double-dummy, active-controlled, non-inferiority study of 843 adult patients with primary hyperlipidemia or mixed dyslipidemia. Patients entered a 6- to 8-week wash-out/dietary lead-in period and then were randomized to a 12 week treatment with either pitavastatin tablets or simvastatin (Table 6). Non-inferiority of pitavastatin to a given dose of simvastatin was considered to be demonstrated if the lower bound of the 95% CI for the mean treatment difference was greater than -6% for the mean percent change in LDL-C.

Lipid results are shown in Table 6. For the percent change from baseline to endpoint in LDL-C, pitavastatin tablets were non-inferior to simvastatin for the two pairwise comparisons: pitavastatin tablets 2 mg vs. simvastatin 20 mg and pitavastatin tablets 4 mg vs. simvastatin 40 mg. Mean treatment differences (95% CI) were 4% (1%, 7%) and 1% (-2%, 4%), respectively.

**Table 6. Lipid Response by Dose of Pitavastatin Tablets and Simvastatin in Adult Patients with Primary Hyperlipidemia or Mixed Dyslipidemia in Study 302 (Mean % Change from Baseline at Week 12)**

Treatment	N	LDL-C	Apo-B	TC	TG	HDL-C	non-HDL-C
Pitavastatin Tablets 2 mg daily	307	-39	-30	-28	-16	6	-36
Pitavastatin Tablets 4 mg daily	319	-44	-35	-42	-17	6	-41
Simvastatin 20 mg daily	107	-35	-27	-25	-16	6	-32
Simvastatin 40 mg daily	110	-43	-34	-31	-16	7	-39

#### Study with Pravastatin in Geriatric Patients (Study 306)

Pitavastatin tablets were compared with Pravastatin Sodium Tablets (referred to as pravastatin) in a randomized, multicenter, double-blind, double-dummy, parallel group, active-controlled non-inferiority study of 942 geriatric patients ( $\geq 65$  years) with primary hyperlipidemia or mixed dyslipidemia. Patients entered a 6- to 8-week wash-out/dietary lead-in period, and then were randomized to a once daily dose of pitavastatin tablets or pravastatin for 12 weeks (Table 7). Non-inferiority of pitavastatin tablets to a given dose of pravastatin was assumed if the lower bound of the 95% CI for the treatment difference was greater than -6% for the mean percent change in LDL-C.

Lipid results are shown in Table 7. Pitavastatin tablets significantly reduced LDL-C compared to pravastatin as demonstrated by the following pairwise dose comparisons: pitavastatin tablets 1 mg vs. pravastatin 10 mg, pitavastatin tablets 2 mg vs. pravastatin 20 mg and pitavastatin tablets 4 mg vs. pravastatin 40 mg. Mean treatment differences (95% CI) were 9% (6%, 12%), 10% (7%, 13%) and 10% (7%, 13%), respectively.

**Table 7. Lipid Response by Dose of Pitavastatin Tablets and Pravastatin in Geriatric Patients with Primary Hyperlipidemia or Mixed Dyslipidemia in Study 306 (Mean % Change from Baseline at Week 12)**

Treatment	N	LDL-C	Apo-B	TC	TG	HDL-C	non-HDL-C
Pitavastatin Tablets 1 mg daily	207	-31	-25	-22	-13	1	-29
Pitavastatin Tablets 2 mg daily	224	-39	-31	-27	-15	2	-36
Pitavastatin Tablets 4 mg daily	210	-44	-37	-31	-22	4	-41
Pravastatin	102	-22	-17	-15	-5	0	-20

10 mg daily	103	-22	-17	-15	-5	0	-20
Pravastatin 20 mg daily	96	-29	-22	-21	-11	-1	-27
Pravastatin 40 mg daily	102	-34	-28	-24	-15	1	-32

#### Study with Simvastatin in Patients with $\geq 2$ Risk Factors for Coronary Heart Disease (Study 304)

Pitavastatin tablets were compared with Simvastatin Tablets (referred to as simvastatin) in a randomized, multicenter, double-blind, double-dummy, active-controlled, non-inferiority study of 351 adult patients with primary hyperlipidemia or mixed dyslipidemia with  $\geq 2$  risk factors for coronary heart disease. After a 6- to 8-week wash-out/dietary lead-in period, patients were randomized to a 12-week treatment with either pitavastatin tablets or simvastatin (Table 8). Non-inferiority of pitavastatin tablets to simvastatin was considered to be demonstrated if the lower bound of the 95% CI for the mean treatment difference was greater than -6% for the mean percent change in LDL-C.

Lipid results are shown in Table 8. Pitavastatin tablets 4 mg was non-inferior to simvastatin 40 mg for percent change from baseline to endpoint in LDL-C. The mean treatment difference (95% CI) was 0% (-2%, 3%).

**Table 8. Lipid Response by Dose of Pitavastatin Tablets and Simvastatin in Adult Patients with Primary Hyperlipidemia or Mixed Dyslipidemia with  $\geq 2$  Risk Factors for Coronary Heart Disease in Study 304 (Mean % Change from Baseline at Week 12)**

Treatment	N	LDL-C	Apo-B	TC	TG	HDL-C	non-HDL-C
Pitavastatin Tablets 4 mg daily	233	-44	-34	-31	-20	7	-40
Simvastatin 40 mg daily	118	-44	-34	-31	-15	5	-39

#### Study with Atorvastatin in Patients with Type 2 Diabetes Mellitus (Study 305)

Pitavastatin tablets were compared with Atorvastatin Calcium Tablets (referred to as atorvastatin) in a randomized, multicenter, double-blind, double-dummy, parallel group, active-controlled, non-inferiority study of 410 adult patients with type 2 diabetes mellitus and mixed dyslipidemia. Patients entered a 6- to 8-week washout/dietary lead-in period and were randomized to a once daily dose of pitavastatin tablets or atorvastatin for 12 weeks. Non-inferiority of pitavastatin tablets was considered to be demonstrated if the lower bound of the 95% CI for the mean treatment difference was greater than -6% for the mean percent change in LDL-C.


Lipid results are shown in Table 9. The treatment difference (95% CI) for LDL-C percent change from baseline was -2% (-6.2%, 1.5%). The two treatment groups were not statistically different on LDL-C. However, the lower limit of the CI was -6.2%, slightly exceeding the -6% non-inferiority limit. The study failed to demonstrate that pitavastatin tablets were not significantly different than atorvastatin in lowering LDL-C in patients with type 2 diabetes mellitus and mixed dyslipidemia.

**Table 9. Lipid Response by Dose of Pitavastatin Tablets and Atorvastatin in Adult Patients with Type 2 Diabetes Mellitus and Mixed Dyslipidemia in Study 305 (Mean % Change from Baseline at Week 12)**

Treatment	N	LDL-C	Apo-B	TC	TG	HDL-C	non-HDL-C
Pitavastatin Tablets 4 mg daily	274	-41	-32	-28	-20	7	-36
Atorvastatin 20 mg daily	136	-43	-34	-32	-27	8	-40

The treatment differences in efficacy in LDL-C change from baseline between pitavastatin tablets and active controls (i.e., atorvastatin, simvastatin, or pravastatin) in the active-controlled studies described above are summarized in Figure 1.

**Figure 1. Treatment Difference in Adjusted Mean Percent Change in LDL-C between Pitavastatin Tablets and the Comparator (Atorvastatin, Simvastatin, or Pravastatin)**



*Pediatric use information is approved for Kowa Co Ltd's LIVALO (pitavastatin) tablets. However, due to Kowa Co Ltd's marketing exclusivity rights, this drug product is not labeled with that information.*

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Pitavastatin tablets for oral administration are provided as white to off-white circular biconvex film-coated tablets that contain 1 mg, 2 mg, or 4 mg of pitavastatin. Each tablet has "OP" debossed on one side and a code number specific to the tablet strength on the other.

### **Packaging**

Pitavastatin tablets are supplied as:

- 1 mg: White to off-white circular biconvex film-coated tablets, debossed with the logo "OP" on one side and "57" on the other side.

☐ Supplied as follows:

☐☐ NDC 42291-905-90; bottles of 90 tablets

- 2 mg: White to off-white circular biconvex film-coated tablets, debossed with the logo "OP" on one side and "58" on the other side.

☐ Supplied as follows:

☐☐ NDC 42291-906-90; bottles of 90 tablets

- 4 mg: White to off-white circular biconvex film-coated tablets, debossed with the logo "OP" on one side and "59" on the other side.

☐ Supplied as follows:

☐☐ NDC 42291-907-90; bottles of 90 tablets

### **Storage**

Store at 25°C (77° F), excursions permitted from 15°C-30°C (59°F-86°F). [See USP Controlled Room Temperature]. Protect from light.

## **17 PATIENT COUNSELING INFORMATION**

### *Myopathy and Rhabdomyolysis*

Advise patients that pitavastatin tablets may cause myopathy and rhabdomyolysis. Inform patients that the risk is increased when taking certain types of medication and they should discuss all medication, both prescription and over the counter, with their healthcare provider. Instruct patients to promptly report any unexplained muscle pain, tenderness or weakness particularly if accompanied by malaise or fever [see *Warnings and Precautions* (5.1)].

### *Hepatic Dysfunction*

Inform patients that pitavastatin tablets may cause liver enzyme elevations and possibly liver failure. Advise patients to promptly report fatigue, anorexia, right upper abdominal discomfort, dark urine or jaundice [see *Warnings and Precautions* (5.3)].

### *Increases in HbA1c and Fasting Serum Glucose Levels*

Inform patients that increases in HbA1c and fasting serum glucose levels may occur with pitavastatin tablets. Encourage patients to optimize lifestyle measures, including regular exercise, maintaining a healthy body weight, and making healthy food choices [see *Warnings and Precautions* (5.4)].

### *Pregnancy*

Advise pregnant patients and patients who become pregnant of the potential risk to a fetus. Advise patients to inform their healthcare provider of a known or suspected pregnancy to discuss if pitavastatin tablets should be discontinued [see *Use in Specific Populations* (8.1)].

### *Lactation*

Advise patients that breastfeeding is not recommended during treatment with pitavastatin tablets [see *Use in Specific Populations* (8.2)].

Manufactured for:

AvKARE

Pulaski, TN 38478

Mfg. Rev. 12/23

AV 12/23 (M)

To request additional information or if you have questions concerning pitavastatin tablets please call AvKARE at 1-855-361-3993.

## **PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**AVKARE.**  
NDC 42291-905-90

**Pitavastatin  
Tablets**

**1 mg**

**90 Tablets Rx Only**

Dosage and use: See Package Insert for Full Prescribing information.  
Each film-coated tablet contains 1.045 mg pitavastatin calcium equivalent to 1 mg pitavastatin.  
Store at 25°C (77°F); excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature].  
Protect from moisture and light. Dispense in an appropriate tight, light-resistant, child-resistant container.

**Keep this and all drugs out of the reach of children.**

Manufactured for:  
**AVKARE**  
Pulaski, TN 38478  
www.avkare.com

Mfg. Rev. 08/22 AV 11/23 (W)

3 42291 90590 0



**AVKARE.**  
NDC 42291-906-90

# Pitavastatin Tablets

**2 mg**

90 Tablets Rx Only

Dosage and use: See Package Insert for Full Prescribing information.  
Each film-coated tablet contains 2.09 mg pitavastatin calcium equivalent to 2 mg pitavastatin.  
Store at 25°C (77°F), excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature].  
Protect from moisture and light. Dispense in an appropriate tight, light-resistant, child-resistant container.

**Keep this and all drugs out of the reach of children.**

Manufactured for:  
**AvKARE**  
Pulaski, TN 38478  
www.avkare.com

Mfg. Rev. 08/22 AV 11/23 (W)





**AVKARE.**  
NDC 42291-907-90

# Pitavastatin Tablets

**4 mg**

90 Tablets Rx Only

Dosage and use: See Package Insert for Full Prescribing information.  
Each film-coated tablet contains 4.18 mg pitavastatin calcium equivalent to 4 mg pitavastatin.  
Store at 25°C (77°F), excursions permitted from 15°C to 30°C (59°F to 86°F). [See USP Controlled Room Temperature].  
Protect from moisture and light. Dispense in an appropriate tight, light-resistant, child-resistant container.

**Keep this and all drugs out of the reach of children.**

Manufactured for:  
**AvKARE**  
Pulaski, TN 38478  
www.avkare.com

Mfg. Rev. 08/22 AV 11/23 (W)




## PITAVASTATIN

pitavastatin calcium tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42291-906(NDC:0480-3632)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PITAVASTATIN CALCIUM (UNII: IYD54XEG3W) (PITAVASTATIN - UNII:M5681Q5F9P)	PITAVASTATIN	2.09 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

<b>MAGNESIUM OXIDE</b> (UNII: 3A3U0GI71G)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>CROSPROVIDONE</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRACETIN</b> (UNII: XHX3C3X673)	

### Product Characteristics

<b>Color</b>	white (White)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Round)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	58;OP
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-906-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	01/16/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205932	01/16/2024	

## PITAVASTATIN

pitavastatin calcium tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42291-907(NDC:0480-3633)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PITAVASTATIN CALCIUM</b> (UNII: IYD54XEG3W) (PITAVASTATIN - UNII:M5681Q5F9P)	PITAVASTATIN	4.18 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	

MAGNESIUM OXIDE (UNII: 3A3U0GI71G)
POVIDONE K30 (UNII: U725QWY32X)
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)
CROSPROVIDONE (UNII: 2S7830E561)
MAGNESIUM STEARATE (UNII: 70097M6I30)
HYPROMELLOSES (UNII: 3NXW29V3WO)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIACETIN (UNII: XHX3C3X673)

### Product Characteristics

Color	white (White)	Score	no score
Shape	ROUND (Round)	Size	9mm
Flavor		Imprint Code	59;OP
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-907-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	01/16/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205932	01/16/2024	

## PITAVASTATIN

pitavastatin calcium tablet, film coated

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42291-905(NDC:0480-3631)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PITAVASTATIN CALCIUM (UNII: IYD54XEG3W) (PITAVASTATIN - UNII:M5681Q5F9P)	PITAVASTATIN	1.045 mg

### Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

<b>MAGNESIUM OXIDE</b> (UNII: 3A3U0GI71G)	
<b>CROSPOVIDONE</b> (UNII: 2S7830E561)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	

### Product Characteristics

<b>Color</b>	white (White)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Round)	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	57;OP
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-905-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	01/16/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA205932	01/16/2024	

**Labeler - AvKARE (796560394)**