

**RIGHT REMEDIES STOMACH RELIEF BISMUTH SUBSALICYLATE 262 MG
SOFTGEL- bismuth subsalicylate capsule, liquid filled
Strive Pharmaceuticals Inc.**

RIGHT REMEDIES Stomach Relief BISMUTH SUBSALICYLATE 262 mg softgels

Drug Facts

***Active ingredient
(in each softgel)***

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea • diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn • indigestion • nausea • gas • belching • fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have

- fever • mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days

- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
 - 2 softgels every 1/2 hour or 4 softgels every hour as needed for diarrhea
 - 2 softgels every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 16 softgels in 24 hours
- use until diarrhea stops but no more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each softgel contains: **salicylate 102 mg**
- store at controlled room temperature 20–25°C (68–77°F)
- protect from light, heat and moisture

Inactive ingredients

beeswax, colloidal silicon dioxide, D&C Red No. 33, FD&C Yellow No. 6, gelatin, glycerin, light liquid paraffin, povidone, purified water, sodium carboxy methyl cellulose, sorbitol sorbitan solution, soy lecithin, titanium dioxide

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

Compare to the active ingredient in **Pepto-Bismol® Liquicaps™***

Stomach Relief

For relief of:

Heartburn, Indigestion

Upset Stomach,

Nausea, Diarrhea

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by The Procter & Gamble company, owner of the registered trademarks Pepto-Bismol® Liquicaps™.

Distributed by: Strive Pharmaceuticals Inc.,
East Brunswick, NJ 08816

PRODUCT OF INDIA

Packaging



Compare to the active ingredient in
Pepto-Bismol® Liquicaps™
NDC 70692-829-84

Stomach Relief

BISMUTH SUBSALICYLATE 262 mg
Upset stomach reliever/Antidiarrheal

For relief of:

- Heartburn, Indigestion
- Upset Stomach,
- Nausea, Diarrhea



Actual Size

48
softgels

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	Purpose
Active ingredient (in each softgel) Bismuth subsalicylate 262 mg.....	Upset stomach relieverand antidiarrheal
Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness	
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	

Drug Facts (continued under label)

(*) This product is not manufactured or distributed by The Procter & Gamble company, owner of the registered trademarks Pepto-Bismol® Liquicaps™.



3 70692 00124 2

LOT: NO VARNISH
EXP: 

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts


Active ingredient (in each softgel)	Purpose
Bismuth subsalicylate 262 mg.....	Upset stomach relieverand antidiarrheal

Uses relieves
■ travelers' diarrhea ■ diarrhea
■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness


Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Drug Facts (continued under label)

(*) This product is not manufactured or distributed by The Procter & Gamble company, owner of the registered trademarks Pepto-Bismol® Liquicaps™.



3 70692 00124 2

LOT: NO VARNISH
EXP: 

Drug Facts (continued)

Allergy alert: Contains salicylate. Do not take if you are
■ allergic to salicylates (including aspirin)
■ taking other salicylate products

Do not use if you have
■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have
■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood)
■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if
■ symptoms get worse or last more than 2 days
■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
■ 2 softgels every 1/2 hour or 4 softgels every hour as needed for diarrhea
- 2 softgels every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 16 softgels in 24 hours
- use until diarrhea stops but no more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- each softgel contains: salicylate 102 mg
- store at controlled room temperature 20–25°C (68–77°F)
- protect from light, heat and moisture

Inactive ingredients

beeswax, colloidal silicon dioxide, D&C Red No. 33, FD&C Yellow No. 6, gelatin, glycerin, light liquid paraffin, povidone, purified water, sodium carboxy methyl cellulose, sorbitol sorbitan solution, soy lecithin, titanium dioxide

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

Distributed by: Strive Pharmaceuticals Inc.,
East Brunswick, NJ 08816 LB7069282984
PRODUCT OF INDIA REV.00-092025/

RIGHT REMEDIES STOMACH RELIEF BISMUTH SUBSALICYLATE 262 MG SOFTGEL

bismuth subsalicylate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70692-829
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
YELLOW WAX (UNII: 2ZA36H0S2V)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	
SOYBEAN LECITHIN (UNII: 1DI56QDM62)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
-------	------	-------	----------

Shape	capsule (Oblong)		Size	22mm	
Flavor			Imprint Code	829	
Contains					
Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:70692-829-84	48 in 1 BOTTLE; Type 0: Not a Combination Product	09/23/2025		
Marketing Information					
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug		M008	09/23/2025		

Labeler - Strive Pharmaceuticals Inc. (080028013)