

**BISACODYL LAXATIVE ENTERIC COATED- bisacodyl tablet, coated  
Good Sense (Geiss, Destin & Dunn, Inc.)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

Bisacodyl USP 5 mg

**Purpose**

Stimulant laxative

**Use**

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

**Warnings**

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- take with a glass of water

---

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

---

### **Other information**

- store between 20°-25C° (68°-77°F)

### **Inactive ingredients**

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow # 6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

### **Questions or comments?**

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

### **Principal Display Panel**

#### **Overnight Relief**

Bisacodyl Laxative

Gentle, Predictable

Overnight Relief

Enteric Coated Tablets (5 mg Each)

\*Compare to active ingredient in Dulcolax®

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

Distributed by:

Geiss, Destin & Dunn, Inc.

Peachtree City, GA 30269

www.valuelabels.com

Product of U.S.A.

### Product Label

**GOODSENSE®**

# **Bisacodyl Laxative**

**GOODSENSE®** NDC 50804-190-03

**Overnight Relief**

# **Bisacodyl Laxative**

*Gentle, Predictable Overnight Relief*



**\*Compare to active ingredient in Dulcolax\***



**25 Enteric Coated Tablets (5 mg Each)**

\*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.  
Distributed by:  
Geiss, Destin & Dunn, Inc.  
Peachtree City, GA 30269  
www.valuelabels.com  
Product of U.S.A  
PLD-C23J FC002038



8 46036 00354 3

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

Lot No.:  
Exp. Date:

if pregnant or breast-feeding, ask a health professional before use.  
Monday-Friday 9AM-5PM EST

<b>Drug Facts</b>		<b>Drug Facts</b>	
<b>Active ingredient</b> (in each tablet) Bisacodyl USP 5 mg.....Stimulant laxative		<b>Purpose</b>	
<b>Use</b> ■ for temporary relief of occasional constipation and irregularly ■ this product generally produces a bowel movement in 6 to 12 hours		<b>Warnings</b> Do not use if you cannot swallow without chewing. ■ Ask a doctor before use if you have stomach pain, nausea or vomiting ■ a sudden change in bowel habits that lasts more than 2 weeks	
<b>When using this product</b> ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk ■ it may cause stomach discomfort, flatness and cramps		<b>Stop use and ask a doctor if</b> ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week	
<b>Questions or comments?</b> Call 1-877-753-3935		<b>Questions or comments?</b> Call 1-877-753-3935	
<b>Other information</b> ■ store between 20°-25°C (68°-77°F)		<b>Other information</b> ■ store between 20°-25°C (68°-77°F)	
<b>Directions</b> ■ take with a glass of water		<b>Directions</b> ■ take with a glass of water	
adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose	children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor	children under 6 years of age	ask a doctor
<b>Inactive ingredients</b> acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide		<b>Inactive ingredients</b> acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide	

## Good Sense Laxative Tablets

### BISACODYL LAXATIVE ENTERIC COATED

bisacodyl tablet, coated

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50804-190
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

#### Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	

<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)
<b>SHELLAC</b> (UNII: 46N107B71O)
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>Polyvinyl Acetate Phthalate</b> (UNII: 58QVG85GW3)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)
<b>POVIDONE</b> (UNII: FZ989GH94E)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	TCL003
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50804-190-03	1 in 1 CARTON	07/16/2010	01/31/2025
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	07/16/2010	01/31/2025

**Labeler** - Good Sense (Geiss, Destin & Dunn, Inc.) (076059836)