

**SAFETUSSIN PM- dextromethorphan doxylamine succinate liquid**  
**Kramer Laboratories**

-----  
**Drug Facts**

<b>Active ingredients (in each 5ml tsp)</b>	<b>Purpose</b>
Dextromethorphan HBr, USP 7.5 mg	Cough Suppressant
Doxylamine Succinate 3.125 mg	Antihistamine

**Uses**

- temporarily relieves cough
- runny nose and sneezing

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleep.

**Ask a doctor before use if you have**

- asthma
- emphysema
- glaucoma
- excessive phlegm (mucus)
- breathing problem
- chronic bronchitis
- persistent or chronic cough
- cough associated with smoking
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if** you are taking sedatives or tranquilizers.

**When using this product**

- do not use more than directed
- excitability may occur, especially in children

- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not take more than 4 doses in any 24-hour period

age	dose
adults and children 12 years and over	4 teaspoons every 6 hours
children 4 to 12 years	ask a doctor
children under 4 years	do not use

**Other information:**

- Protect from excessive heat and freezing. Store at 15° - 30° C (59° - 86° F)
- Phenylketonurics: each teaspoon contains phenylalanine 4.2 mg

**Save this package for complete information**

**Inactive ingredients:** aspartame, benzoic acid, citric acid, glycerin, menthol, methylparaben, natural peppermint flavor, propylene glycol, propylparaben, purified water

**PRINCIPAL DISPLAY PANEL**

**Safetussin® PM**

**NIGHT TIME**

COUGH RELIEF

*Cough Suppressant / Antihistamine*

Specially formulated for adults with

High Blood Pressure

Diabetes

Controls Cough  
Relieves Sneezing  
& Runny Nose

**No** Sugar  
**No** Dyes  
**No** Gluten  
**No** Alcohol

4.0 FL OZ (118mL)

**Safetussin® PM**

ALIVIA LA TOS

**DURANTE LA NOCHE**

Antiusivo / Antihistaminíco

SEGURO para los que padecen de

Presión alta  
Diabetes

Calma la tos,  
alivia estornudos  
y goteo de la nariz

**Sin** Azúcar  
**Sin** Colorantes  
**Sin** Gluten  
**Sin** Alcohol

4 FL. OZ (118 mL)

**Safetussin® PM**

**NIGHT TIME**

COUGH RELIEF

*Cough Suppressant / Antihistamine*

Up To

**8 HOURS OF COUGH RELIEF**

To Help You Rest

**Pharmacist Recommended**

Developed by a pharmacist who recognized the need for simple, safe cough relief for adults afflicted with high blood pressure, diabetes or the common cold.

Specially-formulated **Safetussin®** is free of alcohol, sugar, dyes, decongestants and other ingredients that are unnecessary for safe, adult-strength cough treatment. Pure and simple.

**Questions? Comments?**  
**Call 1-800-824-4894 ● [safetussin.com](http://safetussin.com)**

© 2019 Kramer Laboratories, Inc.  
Bridgewater, NJ 08807 ● [kramerlabs.com](http://kramerlabs.com)

k0519

Cough Suppressant / Antihistamine  
NIGHT TIME  
COUGH RELIEF



Safe  
tussin.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON BOTTLES BROKEN OR MISSING.

### Drug Facts

**Active Ingredients**  
(In each 5mL teaspoon) **Purpose**  
Dextromethorphan HBr USP 7.5mg Cough Suppressant  
Doxylamine Succinate 3.125mg Antihistamine

**Uses**  
• temporarily relieves cough  
• runny nose and sneezing

### Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product, to make a child sleep.

Ask a doctor before use if you have:  
• asthma • emphysema • glaucoma • excessive phlegm (mucus) • breathing problem • chronic bronchitis • persistent or chronic cough • cough associated with smoking • trouble urinating due to enlarged prostate gland • a sodium-restricted diet.

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives and tranquilizers may increase drowsiness.

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

• do not take more than 4 doses in any 24-hour period

age	dose
adults and children 12 years and over	4 teaspoons every 6 hours
children 4 to 12 years	ask a doctor
children under 4 years	do not use

### Other information

• Protect from excessive heat and freezing. Store at 15° - 30° C (59° - 86° F).  
• Phenylephrine: each teaspoon contains phenylephrine 4.2 mg.  
Save this package for complete information.

**Inactive ingredients** aspartame, benzoic acid, citric acid, glycerin, menthyl, methylparaben, natural peppermint flavor, propylene glycol, propylparaben, purified water.

Safe  
tussin.



**NIGHT TIME  
COUGH RELIEF**  
Cough Suppressant / Antihistamine

UP TO  
**8 HOURS OF  
COUGH RELIEF**  
TO HELP YOU REST

Pharmacist Recommended

Developed by a pharmacist who recognized the need for simple, safe cough relief for adults afflicted with high blood pressure, diabetes or the common cold.

Specialty-formulated **Safetussin**<sup>®</sup> is free of alcohol, sugar, dyes, decongestants and other ingredients that are unnecessary for safe, adult-strength cough treatment. Pure and simple.

Questions? Comments?  
Call 1-800-824-4894 • safetussin.com  
© 2019 Kramer Laboratories, Inc.  
Bridgewater, NJ 08807 • kramerlabs.com

Safe  
tussin.



**NIGHT TIME  
COUGH RELIEF**  
Cough Suppressant / Antihistamine

Specially formulated for adults with

**High Blood Pressure**  
 **Diabetes**

Controls Cough  
Relieves Sneezing  
& Runny Nose

✓ No Sugar ✓ No Gluten  
✓ No Dyes ✓ No Alcohol

4.0 FL OZ (118mL)

Safe  
tussin.



ALIVIA LA TOS  
**DURANTE  
LA NOCHE**  
Antitusivo / Antihistamínico

Seguro para los que padecen de

**Presión Alta**  
 **Diabetes**

Calma la tos,  
alivia estornudos  
y goteo de la nariz

✓ Sin Azúcar ✓ Sin Gluten  
✓ Sin Colorantes ✓ Sin Alcohol

4.0 FL OZ (118mL)

K0919



0 53076 17333 8

**Safetussin® PM**  
**NIGHT TIME**  
 COUGH RELIEF  
 Cough Suppressant / Antihistamine

Specially formulated for adults with

High Blood Pressure  
 Diabetes

Controls Cough  
 Relieves Sneezing & Runny Nose

4 FL. OZ. (118 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL ON BOTTLE IS BROKEN OR MISSING.

K0519

www.kramerlabs.com ● www.safetussin.com  
 © 2019 Kramer Laboratories, Inc. Bridgewater, NJ 08807

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>								
<b>Active ingredients</b> (in each 5mL teaspoon) Dextromethorphan HBr, USP 7.5mg ..... Cough Suppressant Doxylamine Succinate 3.125mg ..... Antihistamine	<b>Purpose</b> Cough Suppressant Antihistamine	<b>Stop use and ask a doctor</b> if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.								
<b>Uses</b> • temporarily relieves cough • runny nose and sneezing		<b>If pregnant or breast-feeding</b> , ask a health professional before use.								
<b>Warnings</b> <b>Do not use</b> if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleep.		<b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.								
<b>Ask a doctor before use if you have</b> • asthma • emphysema • glaucoma • excessive phlegm (mucus) • breathing problem • chronic bronchitis • persistent or chronic cough • cough associated with smoking • trouble urinating due to enlarged prostate gland • a sodium-restricted diet		<b>Directions</b> • do not take more than 4 doses in any 24-hour period								
<b>Ask a doctor or pharmacist before use</b> if you are taking sedatives or tranquilizers.		<table border="1"> <thead> <tr> <th>age</th> <th>dose</th> </tr> </thead> <tbody> <tr> <td>adults &amp; children 12 years &amp; over</td> <td>4 teaspoons every 6 hours</td> </tr> <tr> <td>children 4 to 12 years</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 years</td> <td>do not use</td> </tr> </tbody> </table>	age	dose	adults & children 12 years & over	4 teaspoons every 6 hours	children 4 to 12 years	ask a doctor	children under 4 years	do not use
age	dose									
adults & children 12 years & over	4 teaspoons every 6 hours									
children 4 to 12 years	ask a doctor									
children under 4 years	do not use									
<b>When using this product</b> • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives and tranquilizers may increase drowsiness		<b>Other information</b> • Protect from excessive heat and freezing. Store at 15° - 30° C (59° - 86° F) • Phenylethanamine: each teaspoon contains phenylethanamine 4.2mg. Save this package for complete information								
		<b>Inactive ingredients</b> aspartame, benzoic acid, citric acid, glycerin, menthol, methylparaben, natural peppermint flavor, propylene glycol, propylparaben, purified water.								
		kramerlabs.com • safetussin.com © 2019 Kramer Laboratories, Inc. Bridgewater, NJ 08807								

**SAFETUSSIN PM**  
 dextromethorphan doxylamine succinate liquid

<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55505-173
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Dextromethorphan</b> (UNII: 7355X3ROTS) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan Hydrobromide	7.5 mg in 5 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	3.125 mg in 5 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>Aspartame</b> (UNII: Z0H242BBR1)	
<b>benzoic acid</b> (UNII: 8SKN0B0MIM)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>glycerin</b> (UNII: PDC6A3C0OX)	
<b>menthol</b> (UNII: L7T10EIP3A)	
<b>methylparaben</b> (UNII: A2I8C7HI9T)	
<b>MINT</b> (UNII: FV98Z8GITP)	
<b>propylene glycol</b> (UNII: 6DC9Q167V3)	
<b>propylparaben</b> (UNII: Z8IX2SC1OH)	
<b>water</b> (UNII: 059QF0KO0R)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55505-173-33	1 in 1 CARTON	02/09/2017	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/09/2017	

**Labeler** - Kramer Laboratories (122720675)

