

VALUMEDS EXTRA STRENGTH HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet
Cabinet Health P.B.C.

ValuMeds Extra Strength Headache Relief, Acetaminophen, Aspirin and Caffeine

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Pain reliever

Pain reliever aid

Uses

temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- sinusitis
- toothache
- premenstrual & menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning:

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have high blood pressure
- you have asthma

Ask a doctor or pharmacist before use if you are

- any other drug containing an NSAID (prescription or non-prescription)
- a prescription drug for diabetes, gout or arthritis
- any other drug, or are under a doctor's care for any serious condition
- a blood thinning (anticoagulant) or steroid drug

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms appear

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see Overdose warning)**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** magnesium 0.58 mg
- store at controlled room temperature 20° - 25° C (68° - 77° F)
- read all product information before using.

Inactive ingredients

colloidal silicon dioxide, hypermellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, purified water, stearic acid, talc

Questions or comments?

contact **1-908-242-6108** Mon-Fri 8:00 AM to 5:00 PM EST

Package Labeling:

300 caplets
pain reliever/ pain reliever aid

NDC 82125-1023-1

VALUMEDS
Compare to active ingredients in
Excedrin® Extra Strength*

**EXTRA STRENGTH
HEADACHE
RELIEF**

**Acetaminophen 250 mg,
Aspirin 250 mg (NSAID)
and Caffeine 65 mg**

**TAMPER EVIDENT: DO NOT USE IF PRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses temporarily relieves minor aches and pains due to:
 ■ headache ■ a cold ■ arthritis ■ muscular aches
 ■ sinusitis ■ toothache ■ premenstrual & menstrual cramps

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you
 ■ more than 8 caplets in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Stomach bleeding warning:
 This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

(Continued on back of the label)

*This product is not manufactured or distributed by Novartis Consumer Health, Inc., owner of the registered trademark Excedrin®.

Distributed By:
Cabinet Health P.B.C
Brooklyn, NY 11222
REV 11/24
Made in India

ITEM# 10023-01



LOT:
EXP:



Drug Facts (continued)

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use
 ■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask doctor before use if ■ you have liver disease
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have heart disease, liver cirrhosis or kidney disease
 ■ you are taking a diuretic
 ■ you have high blood pressure
 ■ you have asthma

Ask a doctor or pharmacist before use if you are taking
 ■ any other drug containing an NSAID (prescription or non-prescription)
 ■ a prescription drug for diabetes, gout or arthritis
 ■ any other drug, or are under a doctor's care for any serious condition
 ■ a blood thinning (anticoagulant) or steroid drug

Stop use and ask doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding
 ■ feel faint ■ vomit blood ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ ringing in the ears or loss of hearing occurs
 ■ painful area is red or swollen
 ■ pain gets worse or lasts for more than 10 days
 ■ fever gets worse or lasts for more than 3 days
 ■ any new symptoms appear
 These could be signs of a serious condition.

If pregnant or breast-feeding ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children
 In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
 ■ do not use more than directed (see Overdose warning)
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
 ■ children under 12 years: ask a doctor

Other information
 ■ each caplet contains: magnesium 0.58 mg
 ■ store at controlled room temperature 20° - 25° C (68° - 77° F)
 ■ read all product information before using.

Inactive ingredients
 colloidal silicon dioxide, hypermellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, purified water, stearic acid, talc

Questions or comments? contact 1-908-242-6108 Mon-Fri 8:00 AM to 5:00 PM EST

VALUMEDS EXTRA STRENGTH HEADACHE RELIEF
acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:82725-1023
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ACETYLSALICYLIC ACID - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	S53
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:82725-1023-1	300 in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	01/25/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/25/2025	

Labeler - Cabinet Health P.B.C. (117102391)