

# **COLD AND FLU RELIEF- acetaminophen dextromethorphan hbr liquid Best Choice (Valu Merchandisers Company)**

-----  
**Drug Facts**

## **Active ingredients (in each 15 mL)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

## **Purposes**

**Pain reliever/fever reducer**

Cough suppressant

## **Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs by excessive phlegm (mucus)

**Ask a doctor or pharmacist before use if the user is**

taking the blood thinning drug warfarin.

**When using this product,**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse, or lasts more more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product

- mL=milliliter

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

- **When using Daytime or Nighttime products, carefully read each label to ensure correct dosing**

### **Other information**

- **each 15 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate

### **Inactive ingredients**

citric acid, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

### **Questions or comments?**

Call

### **Principal Display Panel**

MULTI-SYMPTOM

#### **Daytime**

#### **Cold & Flu Relief**

ACETAMINOPHEN 325 mg

PAIN RELIEVER/FEVER REDUCER

DEXTROMETHORPHAN HBr 10 mg

COUGH SUPPRESSANT

For relief of:

- Aches, fever & sore throat
- Cough

FOR AGES 6 YEARS & OVER

ALCOHOL FREE

ANTIHISTAMINEFREE

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and DayQuil® are registered trademarks of The Procter & Gamble Company.

DISTRIBUTED BY:

ASSOCIATED WHOLESALE GROCERS, INC.

KANSAS CITY, KANSAS 66106

**Package Label**

**Best Choice**  
**MULTI-SYMPTOM Daytime Cold & Flu Relief**  
**ACETAMINOPHEN 325 mg**  
**PAIN RELIEVER/FEVER REDUCER**  
**DEXTROMETHORPHAN HBr 10 mg**  
**COUGH SUPPRESSANT**

For relief of:  
 • Aches, fever & sore throat  
 • Cough

ALCOHOL FREE  
 ANTIHISTAMINE FREE  
 FOR AGES 6 YEARS & OVER

**Best Choice, 100% Guaranteed**  
[www.bestchoicebrand.com](http://www.bestchoicebrand.com)  
 PROUDLY DISTRIBUTED BY:  
 ASSOCIATED WHOLESALE GROCERS, INC.  
 KANSAS CITY, KANSAS 66106  
 For product information please visit: [bestchoicebrand.com/67497](http://bestchoicebrand.com/67497)

12 FL OZ (355 mL)

PLD-A854A LB009761

0 70038 67497 9

**Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
 Dextromethorphan HBr 10 mg.....Cough suppressant

**Uses**

- temporarily relieves these common cold and flu symptoms
  - minor aches and pains
  - headache
  - sore throat
  - cough due to minor throat and bronchial irritation
  - temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if:  
 ■ adult takes more than 4 doses (30 mL each) of acetaminophen in 24 hours, which is the maximum daily amount for this product

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**



PLD-A854A LB009761

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

- child takes more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

PEEL CORNER FOR MORE DRUG FACTS

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain or cough gets worse, or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see **Overdose warning**)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children 4 to under 6 years	ask a doctor
children under 4 years	do not use

■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- each 15 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

**Drug Facts (continued)**

**Inactive ingredients** citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

# COLD AND FLU RELIEF

acetaminophen dextromethorphan hbr liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63941-291
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-291-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/04/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/04/2024	

**Labeler** - Best Choice (Valu Merchandisers Company) (868703513)

