# BURKHART NEUTRAL PH- sodium fluoride gel Burkhart Dental Supply Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

# **Active Ingredient:**

Sodium Fluoride 2%

# **Purpose:**

Fluoride Treatment Gel

# Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Use Only. This product is not intended for home or unsupervised use.

### Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

# **Dosage and Administration:**

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

- 1. After thorough prophylaxsis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
- 2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
- 3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

### Other Information:

Store at controlled room temperature 59° to 86°F (15°-30°C).

Protect from freezing.

# **Inactive Ingredients:**

Citric Acid, FD&C Red #40, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Potassium Hydroxide, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

# NEUTRAL GEL

with Xylitol

2% Neutral Sodium Fluoride for in-office topical oral application

Gluten Free

Ronly

**BUBBLE GUM** 



**Active Ingredient:** 

Purpose:

Sodium Fluoride 2%......Fluoride Treatment Gel

#### **Indications and Usage:**

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Office Use Only. This product is not intended for home or unsupervised consumer use.

### Warnings:

- · Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- · Read directions carefully before using.

### Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

- After thorough prophylaxis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
- Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
- 3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

### Other Information:

Store at controlled room temperature 59° to 86°F (15°-30°C). Protect from freezing.

### **Inactive Ingredients:**

Citric Acid, FD&C Red #40, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Potassium Hydroxide, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.



Net wt. 16 oz (454 g)

MADE IN THE USA

Manufactured for Burkhart Dental Supply 2502 S. 78th Street Tacoma, Washington 98409 [800] 562-8176 www.burkhartdental.com

99092632 Rev 9/13

# **BURKHART NEUTRAL PH**

sodium fluoride gel

<b>Product</b>	Inform	ation
rioduct	1111 01 111	ation

Product Type HUMAN PRESCRIPTION DRUG Item Code (Source) NDC:43498-151

Route of Administration DENTAL

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII: Q80VPU4080)	FLUORIDE ION	4.086 g in 454 g	

Inactive Ingredients		
Ingredient Name	Strength	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)		
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)		
PHOSPHORIC ACID (UNII: E4GA8884NN)		
POLYSORBATE 20 (UNII: 7T1F30V5YH)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
WATER (UNII: 059QF0KO0R)		
XANTHAN GUM (UNII: TTV12P4NEE)		
XYLITOL (UNII: VCQ006KQ1E)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics			
Color	pink	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

F	Packaging			
#	t Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:43498- 151-15	454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/01/2013	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/14/2012	

# **Labeler -** Burkhart Dental Supply Inc (027532357)