

**RITE-AID GLYCERIN LAXATIVE- glycerin suppository suppository**  
**RITE AID CORPORATION**

-----  
**Rite-Aid® Glycerin Suppositories**

**Active Ingredient**

Active Ingredient (in each suppository)	Purpose
Glycerin, USP 2 grams .....	Laxative

**Uses**

**Uses-** relieves occasional constipation (irregularity)

- generally produces bowel movement in 1/4 to 1 hour

**Warnings**

For rectal use only

**Do not use**

Do not use laxative products for longer than 1 week unless directed by a doctor

**Ask a doctor before use if you have**

Ask a doctor before use if you have

- stomach pain - nausea - vomiting - noticed a sudden change in bowel habits that lasts over 2 weeks

**When using this product**

When using this product you may have rectal discomfort or a burning sensation

**Stop use and ask a doctor if**

Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.**

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

## Directions

adults and children 6 years and over  
rectum and retain for 15 minutes; it need not melt to

Insert 1 suppository well into the  
produce laxative action. Do not exceed

1 suppository daily or as directed by a doctor

children 2 to under 6 years  
children under 2 years

use Child Suppositories  
ask a doctor


## Other information

- keep tightly closed - avoid excessive heat

## Inactive ingredients

purified water, sodium hydroxide, stearic acid

## Rite-Aid® Glycerin Suppositories | 50 Count in a Jar




<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active Ingredient</b> (in each suppository) Glycerin, USP 2 grams	<b>Purpose</b> Laxative	If pregnant or breast-feeding, ask a health professional before use. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Use</b> • relieves occasional constipation (irregularity) • generally produces bowel movement in ¼ to 1 hour		<b>Directions</b>	
<b>Warnings</b> For rectal use only Do not use laxative products for longer than 1 week unless directed by a doctor. Ask a doctor before use if you have • abdominal pain • nausea • vomiting • noticed a sudden change in bowel habits that lasts over 2 weeks When using this product you may have rectal discomfort or a burning sensation Stop use and ask a doctor if • you have rectal bleeding • you fail to have a bowel movement after use of a laxative These could be signs of a serious condition.		adults and children 6 years and over	1 suppository per day or as directed by a doctor
		children 2 to under 6 years	use Child Suppositories
		children under 2 years	ask a doctor
		• Insert 1 suppository well up into the rectum and retain for 15 minutes • It need not melt to produce laxative action	
		<b>Other Information</b> • keep tightly closed • avoid excessive heat	
		<b>Inactive Ingredients</b> purified water, sodium hydroxide, stearic acid	
		<b>Questions or Comments?</b> Call 1-800-748-3243 Monday-Friday 9:00AM - 5:00PM EST	

\*This product is not manufactured or distributed by C.B. Fleet Co., owner of the registered trademark Fleet®.

**DISTRIBUTED BY:**  
RITE AID  
30 HUNTER LANE,  
CAMP HILL, PA 17011  
www.riteaid.com

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.



0 11822 42019 8

## Rite-Aid® Glycerin Suppositories | 100 Count in a Jar

Compare to the active ingredient in Fleet® Glycerin Suppositories\*

# LAXATIVE GLYCERIN SUPPOSITORIES

**GLYCERIN 2g**

often works in minutes for gentle, effective relief of constipation not made with natural rubber latex

**100** ADULT SIZE SUPPOSITORIES

Drug Facts		Purpose	
<b>Active ingredient (in each suppository)</b> Glycerin, USP 2 grams		Laxative	
<b>Uses</b> • relieves occasional constipation (irregularity) • generally produces bowel movement in ¼ to 1 hour			
<b>Warnings</b> <b>For rectal use only</b> <b>Do not use</b> laxative products for longer than 1 week unless directed by a doctor <b>Ask a doctor before use if you have</b> • abdominal pain • nausea • vomiting • noticed a sudden change in bowel habits that lasts over 2 weeks <b>When using this product</b> you may have rectal discomfort or a burning sensation <b>Stop use and ask a doctor if</b> • you have rectal bleeding • you fail to have a bowel movement after use of a laxative These could be signs of a serious condition. <b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.			
<b>Directions</b>		<b>Drug Facts (continued)</b>	
adults and children 6 years and over	1 suppository per day or as directed by a doctor	<b>Directions</b>	
children 2 to under 6 years	use Rite Aid® Infants & Children suppositories	• insert 1 suppository well up into the rectum and retain for 15 minutes	
children under 2 years	ask a doctor	• it need not melt to produce laxative action	
<b>Other information</b> • keep tightly closed • avoid excessive heat			
<b>Inactive ingredients</b> purified water, sodium hydroxide, stearic acid			
<b>Questions or Comments?</b> Call 1-800-748-3243 Monday-Friday 9:00AM - 5:00PM EST			
*This product is not manufactured or distributed by C.B. Fleet Co., owner of the registered trademark Fleet®.			

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE, CAMP HILL, PA 17011 [www.riteaid.com](http://www.riteaid.com)

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

0 11822 42018 1

## RITE-AID GLYCERIN LAXATIVE

glycerin suppository suppository

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-9100
<b>Route of Administration</b>	RECTAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX)	GLYCERIN	2 g

### Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

### Product Characteristics

<b>Color</b>	white (CLEAR)	<b>Score</b>	
<b>Shape</b>	BULLET	<b>Size</b>	32mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-9100-5	50 in 1 JAR; Type 0: Not a Combination Product	01/01/2024	
2	NDC:11822-9100-0	100 in 1 JAR; Type 0: Not a Combination Product	01/01/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	01/01/2012	

**Labeler** - RITE AID CORPORATION (014578892)

**Registrant** - Unipack LLC (116015769)

## Establishment

Name	Address	ID/FEI	Business Operations
Unipack LLC		009248480	manufacture(11822-9100)

Revised: 1/2024

RITE AID CORPORATION